

Kinetics of phycocyanobilin cleavage from C-phycocyanin by methanolysis

Malwade, Chandrakant Ramkrishna; Roda Serrat, Maria Cinta; Christensen, Knud Villy; Fretté, Xavier; Christensen, Lars Porskjær

Published in:
Computer - Aided Chemical Engineering

DOI:
10.1016/B978-0-444-63428-3.50015-1

Publication date:
2016

Document version:
Other version

Citation for pulished version (APA):
Malwade, C. R., Roda Serrat, M. C., Christensen, K. V., Fretté, X., & Christensen, L. P. (2016). Kinetics of phycocyanobilin cleavage from C-phycocyanin by methanolysis. In Z. Kravanja, & M. Bogataj (Eds.), *Computer - Aided Chemical Engineering* (pp. 61-66). Elsevier Masson. <https://doi.org/10.1016/B978-0-444-63428-3.50015-1>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.
Unless otherwise specified it has been shared according to the terms for self-archiving.
If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim.
Please direct all enquiries to puresupport@bib.sdu.dk

Kinetics of Phycocyanobilin Cleavage from C-Phycocyanin by Methanolysis

Chandrakant R. Malwade*, Maria C. Roda-Serrat, Knud V. Christensen, Xavier Fretté, Lars P. Christensen


Department of Chemical Engineering, Biotechnology and Environmental Technology
University of Southern Denmark, Campusvej 55, DK-5230 Odense M, Denmark
crm@kbm.sdu.dk

ESCAPE26




INTRODUCTION

Phycocyanobilin (PCB) is a linear tetrapyrrole chromophore covalently attached to protein subunits of phycobiliproteins, C-Phycocyanin (C-PC) and Allophycocyanin (APC), present in the light harvesting complexes of the blue-green algae *Arthrospira platensis*. PCB absorbs light in the red region of the electromagnetic spectrum, thereby exhibiting a vivid blue color. Therefore, it has great significance to the food industry due to its potential as a natural blue food color. The chemical synthesis of PCB is very complex and economically not feasible. Hence, there is a demand for the development of process to obtain PCB from phycobiliproteins. PCB is attached to the protein subunits through a cysteine residue with a thioether linkage. In this work, the kinetics of the cleavage process of PCB from protein subunits by methanolysis is investigated.



KINETIC MODEL FOR CLEAVAGE OF PCB BY METHANOLYSIS



PCB attached to alpha (left) and beta (right) subunits of C-PC via thioether linkage

Cleavage of PCB can be described either as two first order reactions in parallel:



Or two first order reactions in series:



Where PCB(I) is easily accessible and PCB(II) is less accessible for cleavage, v_1 and v_2 are stoichiometric coefficients of PCB(I) and PCB(II), respectively. In a batch reactor the reactions in parallel will appear as a single first order reaction and can be represented by following set of equations:

$$\frac{dC_1}{dt} = -k_1 \cdot C_1; \quad \frac{dC_2}{dt} = k_1 \cdot C_1 - k_2 \cdot C_2$$

Analytical solutions for set of equations above is:

$$C_1(t) = C_{10} \cdot e^{-k_1 \cdot t}; \quad C_2(t) = \frac{k_1 \cdot C_{10}}{k_1 - k_2} \cdot (e^{-k_2 \cdot t} - e^{-k_1 \cdot t}) + C_{20} \cdot e^{-k_2 \cdot t}$$


Where C_{10} and C_{20} are initial concentration of PCB(I) and PCB(II), respectively

Based on stoichiometry of reaction, the concentration of PCB can be expressed as:

$$C_{\text{PCB}}(t) = v_1 \cdot C_{10} \cdot \left(1 + \frac{v_2}{v_1} + \frac{v_2}{v_1} \cdot \frac{C_{20}}{C_{10}}\right) - v_1 \cdot C_{10} \cdot \left(1 + \frac{v_2}{v_1} + \frac{v_2}{v_1} \cdot \frac{k_1}{k_1 - k_2}\right) \cdot e^{-k_1 \cdot t} + v_1 \cdot C_{10} \cdot \left(\frac{v_2}{v_1} \cdot \frac{k_1}{k_1 - k_2} - \frac{v_2}{v_1} \cdot \frac{C_{20}}{C_{10}}\right) \cdot e^{-k_2 \cdot t}$$

EXPERIMENTAL

- ❑ Linablue (Commercial extract of *Arthrospira platensis*) boiled in 400 mL methanol for 16 h at 65 °C
- ❑ Mixture samples are taken at regular interval for HPLC analysis
- ❑ Three different initial concentration of Linablue used




CONCLUSION

- ❑ Kinetic model describes the experimental data adequately
- ❑ The ratio between v_2 and v_1 is too large compared to the prior findings where a ratio 0.2 and 0.3 is more likely if all PCB is cleaved
- ❑ Although the model explains kinetic observations well, a two step model might be an over simplification

RESULTS

Table 1. Model data fitted to experimental data.

Initial protein concentration (mg/mL)	$v_1 \cdot C_{10}$ (mg/mL)	$\frac{v_2}{v_1}$	$\frac{C_{20}}{C_{10}}$	k_1 (h ⁻¹)	k_2 (h ⁻¹)
5	2.7×10^{-3}	24	1.0×10^{-4}	33	0.29
10	4.7×10^{-3}				
25	12×10^{-3}				



Cleavage of PCB as a function of time. Fully drawn lines are calculated using the model with the parameters from Table 1.

ACKNOWLEDGEMENT

This work is a part of the research within the Project Natural Blue Colorant for Food Applications financially supported by Danish Agency for Science Technology and Innovation to whom the authors are indebted