



University of Southern Denmark

Non-cirrhotic portal hypertension

A possibly benign but complicated disease

Trebicka, Jonel

Published in:
Digestive and Liver Disease

DOI:
10.1016/j.dld.2018.05.018

Publication date:
2018

Document version:
Accepted manuscript

Document license:
CC BY-NC-ND

Citation for pulished version (APA):
Trebicka, J. (2018). Non-cirrhotic portal hypertension: A possibly benign but complicated disease. *Digestive and Liver Disease*, 50(8), 845-846. <https://doi.org/10.1016/j.dld.2018.05.018>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.
Unless otherwise specified it has been shared according to the terms for self-archiving.
If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim.
Please direct all enquiries to puresupport@bib.sdu.dk

Accepted Manuscript

Title: Non-Cirrhotic Portal Hypertension: A possibly benign but complicated disease

Author: Jonel Trebicka

PII: S1590-8658(18)30762-X
DOI: <https://doi.org/10.1016/j.dld.2018.05.018>
Reference: YDLD 3764

To appear in: *Digestive and Liver Disease*

Received date: 22-5-2018

Please cite this article as: Trebicka Jonel. Non-Cirrhotic Portal Hypertension: A possibly benign but complicated disease. *Digestive and Liver Disease* (2018), <https://doi.org/10.1016/j.dld.2018.05.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Title:**Non-Cirrhotic Portal Hypertension: A possibly benign but complicated disease.****Author:****Jonel Trebicka^{1, 2, 3, 4}**

1. Department of Internal Medicine I, University Clinic Bonn, Bonn, Germany

2. European Foundation for the Study of Chronic Liver Failure, Barcelona, Spain

3. Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark

4. Institute for Bioengineering of Catalonia, Barcelona, Spain

Prof. Jonel Trebicka, MD, PhD,

Department of Internal Medicine I, University Clinic Bonn, Sigmund-Freud-Str. 25, 53105

Bonn, Germany. Email: jonel.trebicka@efclif.com

In the last Baveno Consensus Conference held in 2015 a dedicated section described the management of patients with portal vein thrombosis, in presence or absence of cirrhosis (1). In this specific collective of patients few studies have described their natural history and therefore new studies are urgently needed in order to recommend evidence-based management. So far the recommendation regarding the treatment of complications of portal hypertension are aligned with the recommendation in the presence of cirrhosis (1, 2). However, these recommendations are based less on evidence and more on eminence. There are few published studies, which describe the natural history and management procedures in these patients, such as variceal management (3, 4), anticoagulation (5) or shunting procedures (6). The comparison to the cirrhotic patients

has not been done in a systematic manner, but the conclusions drawn have been stressing the similarities of NCPH with liver cirrhosis with portal hypertension.

The paper by Gioia and colleagues (7) presents a step in the right direction to fill in this gap. They nicely describe that the outcome of patients with non-cirrhotic portal hypertension (NCPH) is not more favorable than in patients with a compensated Child A cirrhosis. This is clearly seen in these patients when these two groups are compared with their risk of variceal progression. Also, the risk of development of portal vein thrombosis and variceal bleeding was higher in NCPH patients compared to the Child A cirrhotic patients, while the development of ascites was higher in the cirrhotic patients. This study suggests that the complications of portal hypertension in the patients with NCPH are not due to the same mechanisms as in cirrhosis. Especially interesting is the fact that NCPH are more associated with thrombosis and less with ascites. This indicates that the pathophysiological mechanisms of portal hypertension seem to be more related to pure increase in the portal pressure, rather than the hyperdynamic circulation with consecutive decrease in effective arterial blood volume ultimately affecting the kidney function and leading to ascites (8, 9). These findings suggest that non-selective betablockers might not be the right treatment in these patients. Moreover, the effects one would desire in this situation would be a vasodilating drug to decrease the resistance to portal blood flow and thereby improve the flow and prevent thrombosis, rather than decreasing the portal blood flow and increasing the chance of developing portal vein thrombosis. These thoughts would ultimately change the approach in the patients with NCPH and require further investigations and research in this neglected field. In consequence, the development of treatments should also not be aligned to the treatment of cirrhotic portal hypertension, as it is currently (1, 2).

Therefore, new therapeutic strategies are required for these patients. Due to the small number of cases international initiatives such as EASL-VALDIG-registry will be helpful in the future to perform the correct comparisons and elaborate on the right procedures for these patients. Another issue is the matter of definition, which has not been unified until recently. Therefore, the definitions of NCPH, as well as the concept of advanced chronic liver disease have been introduced (1), which should be used in a consequent manner in future studies.

In summary, the recommendation for the management of NCPH patients should be elaborated independently of the cirrhotic portal hypertension, due to the different course of disease, which is ultimately demonstrated in the present study.

CONFLICT OF INTEREST

This author declares the following financial disclosures:

The author is supported by grants from the Deutsche Forschungsgemeinschaft (SFB TRR57 to P18), Cellex Foundation (PREDICT) and European Union's Horizon 2020 research and innovation program (No 668031).

References:

1. de Franchis R, Faculty BV. Expanding consensus in portal hypertension: Report of the Baveno VI Consensus Workshop: Stratifying risk and individualizing care for portal hypertension. *Journal of Hepatology* 2015;63:743-752.
2. European Association for the Study of the Liver. Electronic address eee. EASL Clinical Practice Guidelines: Vascular diseases of the liver. *J Hepatol* 2015.
3. Siramolpiwat S, Seijo S, Miquel R, Berzigotti A, Garcia-Criado A, Darnell A, Turon F, et al. Idiopathic portal hypertension: natural history and long-term outcome. *Hepatology* 2014;59:2276-2285.
4. Noronha Ferreira C, Seijo S, Plessier A, Silva-Junior G, Turon F, Rautou PE, Baiges A, et al. Natural history and management of esophagogastric varices in chronic noncirrhotic, nontumoral portal vein thrombosis. *Hepatology* 2016;63:1640-1650.
5. De Gottardi A, Trebicka J, Klinger C, Plessier A, Seijo S, Terziroli B, Magenta L, et al. Antithrombotic treatment with direct-acting oral anticoagulants in patients with splanchnic vein thrombosis and cirrhosis. *Liver Int* 2017;37:694-699.
6. Bissonnette J, Garcia-Pagan JC, Albillos A, Turon F, Ferreira C, Tellez L, Nault JC, et al. Role of the transjugular intrahepatic portosystemic shunt in the management of severe complications of portal hypertension in idiopathic noncirrhotic portal hypertension. *Hepatology* 2016;64:224-231.

7. Gioia S, Nardelli S, Pasquale C, et al. Natural history of patients with non cirrhotic portal hypertension: Comparison with patients with compensated cirrhosis. Digestive and Liver Disease. Advance online publication, 03 February 2018, DOI <https://doi.org/10.1016/j.dld.2018.01.132> [Epub ahead of print]
8. Trebicka J. Emergency TIPS in a Child-Pugh B patient: When does the window of opportunity open and close? J Hepatol 2017;66:442-450.
9. Trebicka J. Does Transjugular Intrahepatic Portosystemic Shunt Stent Differentially Improve Survival in a Subset of Cirrhotic Patients? Semin Liver Dis 2018;38:87-96.

ACCEPTED MANUSCRIPT