



University of Southern Denmark

Correction to

Impact of socioeconomic position and distance on mental health care utilization: a nationwide Danish follow-up study

Packness, Aake; Waldorff, Frans Boch; Christensen, René De Pont; Hastrup, Lene Halling; Simonsen, Erik; Vestergaard, Mogens; Halling, Anders

Published in:

Social Psychiatry and Psychiatric Epidemiology

DOI:

10.1007/s00127-018-1556-4

Publication date:

2018

Document version:

Final published version

Document license:

CC BY

Citation for pulished version (APA):

Packness, A., Waldorff, F. B., Christensen, R. D. P., Hastrup, L. H., Simonsen, E., Vestergaard, M., & Halling, A. (2018). Correction to: Impact of socioeconomic position and distance on mental health care utilization: a nationwide Danish follow-up study. *Social Psychiatry and Psychiatric Epidemiology*, 53(9), 1003–1004. <https://doi.org/10.1007/s00127-018-1556-4>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.

Unless otherwise specified it has been shared according to the terms for self-archiving.

If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim. Please direct all enquiries to puresupport@bib.sdu.dk



Correction to: Impact of socioeconomic position and distance on mental health care utilization: a nationwide Danish follow-up study

Aake Packness¹  · Frans Boch Waldorff¹ · René dePont Christensen¹ · Lene Halling Hastrup² · Erik Simonsen^{2,5} · Mogens Vestergaard³ · Anders Halling⁴

Published online: 10 July 2018
© The Author(s) 2018

Correction to:
Soc Psychiatry Psychiatr Epidemiol (2017) 52:1405–1413
<https://doi.org/10.1007/s00127-017-1437-2>

In the original publication of this article, Table 3 was published incorrectly. The corrected table is shown below.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

The original article can be found online at <https://doi.org/10.1007/s00127-017-1437-2>.

✉ Aake Packness
apackness@health.sdu.dk

- ¹ Research Unit of General Practice, Institute of Public Health, University of Southern Denmark, J.B. Winsløvs Vej 9A, 5000 Odense C, Denmark
- ² Psychiatric Research Unit, Region of Zealand, Slagelse, Denmark
- ³ Department of Public Health, Institute of General Medical Practice, Aarhus University, Aarhus, Denmark
- ⁴ Centre for Primary Health Care Research, Lund University, Lund, Sweden
- ⁵ Faculty of Health and Medical Sciences, Institute of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

Table 3 Separate analyses of correlation of income or education with mental health service used, in odds ratios and incidence rate ratios of contact

Odds ratio of contact to services ^a																
Logit	Contact out-patient psychiatrist			Contact psychologist			Contact GP-MHS			Psychiatric emergency clinic			Admission MH > 1 day			
	OR	P	CI	N	OR	P	CI	OR	P	CI	OR	P	CI	OR	P	CI
<i>Family equivalent income^b</i>																
Highest third	1			50,373	1			1			1			1		
Middle third	1.03	0.414	0.97	1.09	0.71	< 0.001	0.67	0.75	0.90	< 0.001	0.86	0.95	0.94	0.344	0.82	1.07
Lower third	1.25	< 0.001	1.17	1.34	0.49	< 0.001	0.46	0.53	0.81	< 0.001	0.77	0.86	0.82	0.007	0.71	0.95
<i>Education (years)^b</i>																
12+	1			48,183	1			1			1			1		
10–12	0.92	0.005	0.86	0.97	0.67	< 0.001	0.63	0.71	0.89	< 0.001	0.85	0.94	0.99	0.888	0.86	1.13
< 10	0.95	0.103	0.89	1.01	0.37	< 0.001	0.35	0.40	0.71	< 0.001	0.67	0.75	0.90	0.147	0.78	1.04
Incidence rate ratios of contact among patients who had one visit or more ^a																
Poisson	Out-patient psychiatrist			Psychologist			GP-MHS			Psych. emergency clinic			Admission MH > 1 day			
	IRR	P	CI	N	IRR	P	CI	IRR	P	CI	IRR	P	CI	IRR	P	CI
<i>Family equivalent income^b</i>																
Highest third	1			11,113	1			9033	1				17,638	1		
Middle third	0.90	< 0.001	0.88	0.91	0.93	< 0.001	0.91	0.95	1.00	0.639	0.98	1.02	1.02	0.734	0.91	1.15
Lower third	0.83	< 0.001	0.81	0.84	0.94	< 0.001	0.91	0.96	0.94	< 0.001	0.92	0.97	1.06	0.364	0.94	1.14
<i>Education (years)^b</i>																
12+	1			10,659	1			8869	1				17,038	1		
10–12	0.92	< 0.001	0.90	0.93	0.92	< 0.001	0.90	0.93	0.99	0.395	0.97	1.01	0.98	0.729	0.87	1.10
< 10	0.75	< 0.001	0.74	0.76	0.80	< 0.001	0.79	0.82	0.93	< 0.001	0.91	0.96	1.06	0.370	0.94	1.19

Psychiatric emergency clinic includes emergency contacts and admissions up to 1 day

Results significant within a 95% confidence interval are marked in bold

GP-MHS mental health services by general practitioner (talk therapy), Admission MH admission to mental hospital, IRR incidence rate ratio, CI confidence interval, P 0.05

^aTwo separate analyses of correlation of income or education with mental health service used

^bAdjusted for age, gender, country of origin, cohabitating status, access to vehicle, comorbidity psychiatric, comorbidity somatic