



University of Southern Denmark

Using BCG vaccine to enhance non-specific protection of health care workers during the COVID-19 pandemic

A structured summary of a study protocol for a randomised controlled trial in Denmark

Madsen, Anne Marie Rosendahl; Schaltz-Buchholzer, Frederik; Benfield, Thomas; Bjerregaard-Andersen, Morten; Dalgaard, Lars Skov; Dam, Christine; Ditlev, Sisse Bolm; Faizi, Gulia; Johansen, Isik Somuncu; Kofoed, Poul Erik; Kristensen, Gitte Schultz; Loekkegaard, Ellen Christine Leth; Mogensen, Christian Backer; Mohamed, Libin; Ostefeld, Anne; Oedegaard, Emilie Sundhaugen; Soerensen, Marcus Kjaer; Wejse, Christian; Jensen, Aksel Karl Georg; Nielsen, Sebastian; Krause, Tyra Grove; Netea, Mihai G.; Aaby, Peter; Benn, Christine Stabell

Published in:
Trials

DOI:
10.1186/s13063-020-04714-3

Publication date:
2020

Document version:
Final published version

Document license:
CC BY

Citation for pulished version (APA):

Madsen, A. M. R., Schaltz-Buchholzer, F., Benfield, T., Bjerregaard-Andersen, M., Dalgaard, L. S., Dam, C., Ditlev, S. B., Faizi, G., Johansen, I. S., Kofoed, P. E., Kristensen, G. S., Loekkegaard, E. C. L., Mogensen, C. B., Mohamed, L., Ostefeld, A., Oedegaard, E. S., Soerensen, M. K., Wejse, C., Jensen, A. K. G., ... Benn, C. S. (2020). Using BCG vaccine to enhance non-specific protection of health care workers during the COVID-19 pandemic: A structured summary of a study protocol for a randomised controlled trial in Denmark. *Trials*, 21, Article 799. <https://doi.org/10.1186/s13063-020-04714-3>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.

Unless otherwise specified it has been shared according to the terms for self-archiving.

If no other license is stated, these terms apply:


- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

LETTER

Open Access



Using BCG vaccine to enhance non-specific protection of health care workers during the COVID-19 pandemic: A structured summary of a study protocol for a randomised controlled trial in Denmark

Anne Marie Rosendahl Madsen^{1*} , Frederik Scholtz-Buchholzer¹, Thomas Benfield², Morten Bjerregaard-Andersen³, Lars Skov Dalgaard⁴, Christine Dam⁵, Sisse Bolm Ditlev⁶, Gulia Faizi³, Isik Somuncu Johansen⁷, Poul-Erik Kofoed⁸, Gitte Schultz Kristensen⁹, Ellen Christine Leth Loekkegaard¹⁰, Christian Backer Mogensen⁹, Libin Mohamed⁸, Anne Ostenfeld¹⁰, Emilie Sundhaugen Oedegaard¹, Marcus Kjaer Soerensen¹, Christian Wejse¹¹, Aksel Karl Georg Jensen¹², Sebastian Nielsen¹, Tyra Grove Krause¹³, Mihai G. Netea^{14,15}, Peter Aaby¹ and Christine Stabell Benn¹

Abstract

Objectives: The Bacille Calmette-Guérin (BCG) vaccine against tuberculosis is associated with non-specific protective effects against other infections, and significant reductions in all-cause morbidity and mortality have been reported. We aim to test whether BCG vaccination may reduce susceptibility to and/or the severity of COVID-19 and other infectious diseases in health care workers (HCW) and thus prevent work absenteeism.

The primary objective is to reduce absenteeism due to illness among HCW during the COVID-19 pandemic. The secondary objectives are to reduce the number of HCW that are infected with SARS-CoV-2, and to reduce the number of hospital admissions among HCW during the COVID-19 pandemic.

Hypothesis: BCG vaccination of HCW will reduce absenteeism by 20% over a period of 6 months.

Trial design: Placebo-controlled, single-blinded, randomised controlled trial, recruiting study participants at several geographic locations. The BCG vaccine is used in this study on a different indication than the one it has been approved for by the Danish Medicines Agency, therefore this is classified as a phase III study.

Participants: The trial will recruit 1,500 HCW at Danish hospitals.

To be eligible for participation, a subject must meet the following criteria: Adult (≥ 18 years); Hospital personnel working at a participating hospital for more than 22 hours per week.

A potential subject who meets any of the following criteria will be excluded from participation in this study:

(Continued on next page)

* Correspondence: arosendahl@health.sdu.dk

¹Bandim Health Project, OPEN, Department of Clinical Research, University of Southern Denmark/Odense University Hospital, Odense, Denmark
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

(Continued from previous page)

- Known allergy to components of the BCG vaccine or serious adverse events to prior BCG administration
- Known prior active or latent infection with *Mycobacterium tuberculosis* (*M. tuberculosis*) or other mycobacterial species
- Previous confirmed COVID-19
- Fever (>38 C) within the past 24 hours
- Suspicion of active viral or bacterial infection
- Pregnancy
- Breastfeeding
- Vaccination with other live attenuated vaccine within the last 4 weeks
- Severely immunocompromised subjects. This exclusion category comprises:
 - a) subjects with known infection by the human immunodeficiency virus (HIV-1)
 - b) subjects with solid organ transplantation
 - c) subjects with bone marrow transplantation
 - d) subjects under chemotherapy
 - e) subjects with primary immunodeficiency
 - f) subjects under treatment with any anti-cytokine therapy within the last year
 - g) subjects under treatment with oral or intravenous steroids defined as daily doses of 10 mg prednisone or equivalent for longer than 3 months
 - h) Active solid or non-solid malignancy or lymphoma within the prior two years
- Direct involvement in the design or the execution of the BCG-DENMARK-COVID trial

Intervention and comparator: Participants will be randomised to BCG vaccine (BCG-Denmark, AJ Vaccines, Copenhagen, Denmark) or placebo (saline). An adult dose of 0.1 ml of resuspended BCG vaccine (intervention) or 0.1 ml of sterile 0.9% NaCl solution (control) is administered intradermally in the upper deltoid area of the right arm. All participants will receive one injection at inclusion, and no further treatment of study participants will take place.

Main outcomes: Main study endpoint: Days of unplanned absenteeism due to illness within 180 days of randomisation.

Secondary study endpoints: The cumulative incidence of documented COVID-19 and the cumulative incidence of hospital admission for any reason within 180 days of randomisation.

Randomisation: Randomisation will be done centrally using the REDCap tool with stratification by hospital, sex and age groups (+/- 45 years of age) in random blocks of 4 and 6. The allocation ratio is 1:1.

Blinding (masking): Participants will be blinded to treatment. The participant will be asked to leave the room while the allocated treatment is prepared. Once ready for injection, vaccine and placebo will look similar, and the participant will not be able to tell the difference.

The physicians administering the treatment are not blinded.

Numbers to be randomised (sample size): Sample size: N=1,500. The 1,500 participants will be randomised 1:1 to BCG or placebo with 750 participants in each group.

Trial Status: Current protocol version 5.1, from July 6, 2020.

Recruitment of study participants started on May 18, 2020 and we anticipate having finished recruiting by the end of December 2020.

Trial registration: The trial was registered with EudraCT on April 16, 2020, EudraCT number: 2020-001888-90, and with ClinicalTrials.gov on May 1, 2020, registration number NCT04373291.

Full protocol: The full protocol is attached as an additional file, accessible from the Trials website (Additional file 1). In the interest in expediting dissemination of this material, the familiar formatting has been eliminated; this Letter serves as a summary of the key elements of the full protocol.

Keywords: COVID-19, Randomised controlled trial, Protocol, BCG vaccine, NSEs/Non-specific effects of vaccines, Heterologous effects of vaccines, Health care workers, Pandemic, Immune training.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13063-020-04714-3>.

Additional file 1.

Acknowledgements

Open Patient data Explorative Network (OPEN), Odense University Hospital, Region of Southern Denmark, provides solutions for E-data capture, randomization and data management.

Authors' contributions

CSB is the sponsor, AMRM is the principal investigator and FSB is investigator. CSB, PA, FSB and AMRM conceived and designed the trial and co-authored the protocol with help from TKG and MGN. TB, MBA, LSD, SBD, ISJ, PEK, ECLL, CBM and CW helped organising recruitment at the hospitals. AO, CD, GF, GSK, LM, ESO and MKS assisted with the recruitment of participants. AGJ and SN are the trial statisticians. The authors read and approved the final manuscript.

Funding

We have received a grant of 30,000 DKK (approx. 4,000 euros) from "Torben and Alice Frimodts Fund" (Danish tax number/CVR 11592333) to cover running costs. AJ Vaccines, Copenhagen, Denmark, have donated the BCG vaccines for the trial. Statens Serum Institut, Copenhagen, Denmark, has kindly provided rapid tests for COVID-19 antibody testing.

None of the parties in the study group have any economic interest in the company or in the production of BCG vaccines. The funders have had no influence on the design of the study and will not have influence on any aspect of the analysis, interpretation or reporting of the study results. We are currently applying for further funding from foundations that support projects aiming to mitigate the effect of the COVID-19 pandemic.

Availability of data and materials

Trial data is registered and stored via OPEN (Open Patient data Explorative Network, University of Southern Denmark and Odense University Hospital, Denmark) on secure servers in the Region of Southern Denmark. The investigators and sponsors representative will have access to the final dataset. Data will be available from the author on reasonable request and pending ethics approval (arosendahl@health.sdu.dk).

Ethics approval and consent to participate

The trial has been approved by The Regional Committee on Health Research Ethics for Southern Denmark on April 30, 2020 (reference number: S-20200062C) and by the Danish Medicines Agency on April 27, 2020 (reference number 2020041936).

Before participating, all subjects are asked to sign a consent form after receiving oral and written information about the study.

We declare that we certify that this trial has received ethical approval from the appropriate ethical committee as described above.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Bandim Health Project, OPEN, Department of Clinical Research, University of Southern Denmark/Odense University Hospital, Odense, Denmark. ²Center of Research & Disruption of Infectious Diseases (CREDID), Department of Infectious Diseases, Copenhagen University Hospital, Amager and Hvidovre Hospital, Hvidovre, Denmark. ³Department of Medicine, Sydvestjysk Hospital, Esbjerg, Denmark. ⁴Department of Medicine, Herning Hospital, Herning, Denmark. ⁵Department of Respiratory Medicine, Bispebjerg and Frederiksberg Hospital, Copenhagen, Denmark. ⁶The Copenhagen Center for Translational Research (CCTR), Bispebjerg Hospital, Copenhagen, Denmark. ⁷Department of Infectious Diseases, Odense University Hospital, Odense, Denmark. ⁸Department of Pediatrics, Lillebaelt Hospital, Kolding, Denmark. ⁹Department of Medicine, Soenderjylland Hospital, Aabenraa, Denmark.

¹⁰Department of Gynaecology and Obstetrics, Nordsjaellands Hospital, Hilleroed, Denmark. ¹¹Department of Infectious Diseases, Aarhus University Hospital, Skejby, Denmark. ¹²Section of Biostatistics, Department of Public Health, University of Copenhagen, Copenhagen, Denmark. ¹³Statens Serum Institut, Copenhagen, Denmark. ¹⁴Department of Internal Medicine, Radboud University Medical Centre, Nijmegen, the Netherlands. ¹⁵Department for Genomics & Immunoregulation, Life and Medical Sciences Institute (LIMES), University of Bonn, Bonn, Germany.

Received: 21 August 2020 Accepted: 27 August 2020

Published online: 17 September 2020

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

