



University of Southern Denmark

Postponement of cardiovascular outcomes by statin use

A systematic review and meta-analysis of randomized clinical trials

Hansen, Morten Rix; Hróbjartsson, Asbjørn; Pottegård, Anton; Damkier, Per; Madsen, Kenneth Grønkjær; Pareek, Manan; Olesen, Morten; Hallas, Jesper

Published in:

Basic & Clinical Pharmacology & Toxicology

DOI:

10.1111/bcpt.13485

Publication date:

2021

Document version:

Accepted manuscript

Citation for published version (APA):

Hansen, M. R., Hróbjartsson, A., Pottegård, A., Damkier, P., Madsen, K. G., Pareek, M., Olesen, M., & Hallas, J. (2021). Postponement of cardiovascular outcomes by statin use: A systematic review and meta-analysis of randomized clinical trials. *Basic & Clinical Pharmacology & Toxicology*, 128(2), 286-296. <https://doi.org/10.1111/bcpt.13485>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.
Unless otherwise specified it has been shared according to the terms for self-archiving.
If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim.
Please direct all enquiries to puresupport@bib.sdu.dk

DR. MORTEN RIX HANSEN (Orcid ID : 0000-0002-1582-7866)

Article type : Original Article

Postponement of Cardiovascular Outcomes by Statin Use: A Systematic review and Meta-Analysis of Randomized Clinical Trials

Morten Rix Hansen MD, PhD^{1,2}; Asbjørn Hróbjartsson MD, PhD^{3,4,5}; Anton Pottegård
MScPharm, PhD, DMSc^{1,6}; Per Damkier MD, PhD^{2,7}; Kenneth Grønkjær Madsen MD¹;
Manan Pareek MD, PhD^{8,9}; Morten Olesen¹; Jesper Hallas MD, DMSc^{1,2}

1) Clinical Pharmacology and Pharmacy, Department of Public Health,
University of Southern Denmark, Odense, Denmark

2) Department of Clinical Biochemistry and Pharmacology
Odense University Hospital, Denmark

3) Centre for Evidence-Based Medicine Odense, Odense University Hospital, Denmark

4) Department of Clinical Research, University of Southern Denmark

5) Odense Explorative Patient data Network (OPEN), Odense University Hospital,
Denmark

6) Hospital Pharmacy Funen, Odense University Hospital, Denmark

7) Department of Clinical Research, University of Southern Denmark

8) Department of Cardiology, North Zealand Hospital, Hillerød, Denmark

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/BCPT.13485](https://doi.org/10.1111/BCPT.13485)

This article is protected by copyright. All rights reserved

9) Department of Internal Medicine, Yale New Haven Hospital, New Haven, CT, USA

Abstract: 199

Main text: 2138

Correspondence

Morten Rix Hansen, MD PhD

Clinical Pharmacology and Pharmacy

University of Southern Denmark

J. B. Winsløws Vej 19, 2

5000 Odense C, Denmark

E-mail: mrix@health.sdu.dk

Keywords: Effect measures, Meta-analysis, outcome postponement, cholesterol, statins, cardiovascular endpoints.

ABSTRACT

Objective: To estimate the average outcome postponement (gain in days to an event) for cardiovascular outcomes in a meta-analysis of randomized, controlled statin trials, including any myocardial infarction, any stroke and cardiovascular death.

Design: Systematic review of large randomized, placebo-controlled trials of statin use, including a random-effects meta-analysis of all included trials.

Data Sources: We searched Medline (15 July 2019) and ClinicalTrials.gov (16 October 2019).

Eligibility criteria for selecting studies: Randomized, placebo-controlled trials of statin use that included at least 1000 participants. We identified 15 cardiovascular outcomes that were reported in more than 2 trials.

Results: We included 19 trials. The summary outcome postponements for the 15 cardiovascular outcomes varied between -1 and 38 days. For four major outcomes, the summary outcome postponement in days were: cardiovascular mortality, 9.27 days (95% CI: 3.6 to 14.91; $I^2 = 72%$; 9 trials) non-vascular

and non-cardiovascular mortality, 1.5 days (95% CI: -2.2 to 5.3; $I^2 = 0\%$; 6 trials) any myocardial infarction 18.0 days (95% CI; 12.1 to 24.1; $I^2 = 92\%$; 15 trials); and any stroke, 6.1 days (95% CI; 2.86 to 9.39; $I^2 = 66\%$; 14 trials).

Conclusion: Statin treatment provided a small, average postponement of cardiovascular outcomes during trial duration.

Strengths

- This is the first systematic review and meta-analysis on statins effect on cardiovascular outcomes that employs outcome postponement.
- Outcome postponement is an effect measure that conveys the treatment effect in terms of postponement in time e.g. days, of an outcome, such as death, achieved by taking the drug.
- Patients show much more responsiveness when being informed using outcome postponement than by using other measures, i.e., their preferences with respect to accepting or rejecting a proposed drug intervention changes when they are presented with different values for outcome postponement.

Limitations

- The main limitation of using outcome postponement is the inability of the measure to capture the benefit of continued treatment in the decades after trial termination, although this is a limitation that applies to all effect measures currently used

INTRODUCTION

Although statins have been a subject of trials for over 30 years, controversies about their effect magnitude still exist, and different trials have demonstrated substantial variations in effect size (1). The effect size

of preventive interventions, including statins, is often expressed as the relative risk reduction or number needed to treat (NNT). However, some patients may find it difficult to understand such measures (1–3). Outcome postponement, i.e., the average gain in event-free time of the study endpoint in all treated, is an alternative way to illustrate the effect of practically any preventive interventions (5,6), which has shown some promise in terms of patients' understanding (2).

Our group found an all-cause mortality postponement with statin use of 12.6 days (95% CI; 7.1-18.0) during five years of trial duration (7). The effect of statins may be larger on specific cardiovascular events, and since these drugs were designed to prevent progression of atherosclerosis, all-cause mortality might not be an entirely appropriate measure of their overall effect, especially as severe non-fatal cardiovascular outcomes would impact on patients' quality of life (8).

The aim of this meta-analysis was to estimate the average outcome postponement on fatal and non-fatal cardiovascular outcomes in randomized studies comparing statins to placebo.

METHODS

We systematically reviewed large placebo-controlled trials of statins, calculated the postponement of outcomes and subjected it to a meta-analysis.

Data Sources and Searches

We searched Medline (search index date: 15 July 2019) using the following MeSH terms: statins, placebo, and random*. We also searched ClinicalTrials.gov (search index date: 16 October 2019) using the following terms: “statins” AND “placebo” (interventional). We screened the reference lists of the included papers for relevant publications.

Study Selection

We included randomized trials of at least 1000 patients, in which a statin intervention (any drug) was compared with placebo, with a follow-up of minimum two years.

Data Extraction and Quality Assessment

Two authors (MRH and KGM) independently extracted trial characteristics and outcome data from each included trial published in a journal. Any discrepancies were resolved by consensus. The extracted trial characteristics included duration; whether it represented primary, secondary or mixed intervention; and baseline low-density lipoprotein (LDL) cholesterol level. We defined primary prevention trials as trials in which no patients had manifest cardiovascular disease at baseline, secondary prevention trials as those in which all patients had cardiovascular disease, and mixed prevention trials as trials including patients both with and without cardiovascular disease. Two physicians independently performed trial classification. The effect data included hazard ratio (HR), or relative risk (RR), and 95% confidence interval (CI), and the event rate or cumulative events in the placebo group. For trials that reported relative risk ($n = 8$) instead of hazard ratio as the outcome measure, we used relative risk as a proxy for hazard ratio. These measures are very similar when the cumulative risk is low (9). We used the five-year standardized outcome postponements as input to reduce the influence of trial duration on outcome postponement and thereby to reduce the level of heterogeneity as shown in Hansen et al. (7).

Bias of the individual trials was assessed, and outcomes were grouped after type, using the Cochrane's risk of bias tool (10). If the published papers did not contain the essential parameters for our model, e.g., mean trial follow-up, we contacted the authors. Five groups provided additional information. We followed PRISMA guidelines for systematic literature review and meta-analysis (11)

Data Synthesis and Analysis: Postponement of Outcomes in Single Trials

We calculated postponement of outcomes in each trial using two different methods, a mathematical modelling of the area between survival curves and a pixel counting of the area. We used linear regression to compare model-derived estimates with the corresponding estimates derived by pixel counting. Details of the two methods have been presented previously (7). A description is given in appendix A, B and C in the supplementary file.

Data Synthesis and Analysis: Meta-Analysis

We performed meta-analyses on postponement of cardiovascular events using inverse variance weighting and random-effects models (Stata 15, Stata Corp, Texas, USA). We used the five-year standardized outcome postponements as input. Finally, to assist in the interpretation of the postponement meta-analysis, we performed a HR-based meta-analysis of each cardiovascular event.

In order to account for heterogeneity by trial characteristics, we subgrouped according to high vs. low

overall risk of bias. In addition, we analysed trials according to the purpose of prevention, whether it be primary, secondary or mixed.

Cardiovascular outcomes

We selected categories of cardiovascular outcomes that had been investigated in three trials or more. These included cardiac death, cardiovascular death, death from coronary heart disease, any myocardial infarction, non-fatal myocardial infarction, any cardiovascular event, any coronary event, unstable angina, coronary revascularization, non-fatal stroke, fatal stroke, any stroke, death from non-vascular and non-cardiovascular causes, hospitalization for unstable angina, and hospitalization for heart failure.

We categorized outcomes as “coronary revascularization” if reported as: coronary revascularization, revascularization procedures, percutaneous coronary intervention (PCI), coronary artery bypass graft surgery (CABG), and PCI or CABG. We categorized outcomes as “any myocardial infarction” if reported as: any myocardial infarction and death from CHD or non-fatal myocardial infarction. We categorized outcomes as “non-cardiovascular death” if reported as: death from non-cardiovascular causes and death from non-vascular causes. We categorized outcomes as “any stroke” if reported as: any stroke, fatal stroke and non-fatal cerebrovascular events. Outcomes were combined in this way if they, by clinical reasoning, were thought to have large overlap.

The systematic review was registered in PROSPERO [CRD42016037507].

Patient involvement

Patients were not involved in the planning or conduct of the study.

Role of the Funding source:

This study was funded by the University of Southern Denmark. The funder played no role in the design, conduct or reporting.

RESULTS

We identified 120 potentially eligible trials of which 101 were excluded (Figure 1), leaving 19 trials that met our inclusion criteria. The basic characteristics of the trials are reported in Table 1. Of the 19 trials, four (21%) were for primary prevention, six (32%) were for secondary prevention, and nine (47%) were for mixed prevention.

Table 2 shows the results of the meta-analyses grouped by cardiovascular outcome. The modelled outcome postponement for the four major outcomes were (standardized to 5 years of trial duration); cardiovascular mortality 9.3(95% CI: 3.6 to 14.9; $I^2 = 72\%$; trials); non-vascular and non-cardiovascular mortality 1.5 days (95% CI: -2.2 to 5.3; $I^2 = 0\%$; 6 trials), any myocardial infarction 18.0 days (95% CI: 12.1 to 24.1; $I^2 = 92\%$; 15 trials), and any stroke 6.1 days (95% CI: 2.9 to 9.4; 14 trials). Forest plots for these estimates are presented in figures 2-5, the remaining forest plots included in our analysis are shown as E-Figure 3-13.

Table 3 presents the results of modelled outcome postponement from the individual trials (standardized to a 5-year trial duration). The largest gain in outcome postponement was found for any cardiovascular event in the SPARCL trial, i.e., 83 days (95% CI; 54 to 109). The least favourable outcome was found in the 4D trial for any stroke, -25.0 (-72.0 to 7.7). E-Figure 1 presents the non-standardized outcome postponements, and corresponding results obtained by use of the Pixel Counting Method are shown in E-Figure 2. We found a strong agreement between modelled outcome and corresponding estimates from pixel counting, with a slope of 0.95 with linear regression.

When subgrouping on trial duration, the postponement was substantially larger among trials with a duration above ≥ 5 years than below < 5 years, 18.6 days vs 6.3 days, respectively.

All included trials were very large, with accessible protocols, concealed allocation and blinding. All trials had a low overall risk of bias as evaluated by the Cochrane risk of bias tool. No trials were assessed as high risk of attrition bias; only two trials were assessed as unclear, the rest were assessed as low risk of bias.

Treatment switches are a concern as some trials described varying percentages of patients (range, 4.8-25.4%) assigned to the placebo group switching to statin treatment, or were assigned to the statin group, but stopped their treatment. We analysed treatment switches with respect to any myocardial infarction. Outcome postponement was 16.3 days (CI, 10.0-22.7) in the nine trials with high cumulative incidence ($>4\%$) of treatment switches, 73.4 days (CI, 54.1-92.7) in the single trial with a low degree of switching, and 16.9 days (PI, 6.5 – 27.4) in the 5 trials with an unclear degree of switching.

All trials included in this meta-analysis were analysed using the intention-to-treat principle which protects against bias due to difference in prognosis between patients switching treatment groups and not, but the procedure is conservative and may bias the effect estimate towards the null value. Some of the included

trials reported large proportions of patients switching from the placebo group to the statin group, or patients who were in the statin group, but ceased treatment (range, 4.8-32%). For any myocardial infarction, outcome postponement was 16.3 days (95% CI; 10.0-22.7) in the nine trials with high cumulative incidence (>4%) of treatment switches, 55.4 days (95% CI; 40.9-70.0) in the single trial with a low degree of switching, and 13.4 days (95% CI; 6.5 – 20.3) in the five trials with an unclear degree of switching.

DISCUSSION

In this meta-analysis of 19 large randomized, controlled trials, we demonstrated that statin treatment resulted in small average outcome postponement for cardiovascular endpoints. Since statins primarily prevent progression of atherosclerosis (8), we anticipated that the postponement for strictly cardiovascular outcomes would be larger than for all-cause death. In general, our findings were consistent with this model of understanding; postponements were slightly more pronounced for cardiac outcomes than for all-cause mortality, and strictly non-vascular outcomes were not postponed at all. As expected, the effect of statin therapy on stroke prevention was modest. For four major outcomes, i.e., cardiovascular mortality, non-cardiovascular mortality, any myocardial infarction and any stroke, the summary outcome postponement was 9.3, 1.5, 18.0 and 6.1 days, respectively, when standardized to 5 years of trial duration.

When we stratified according to prevention type, we found the largest postponement in the secondary prevention group, which was expected. Our HR-based meta-analysis showed particularly large relative risk reductions for cardiovascular mortality and non-fatal myocardial infarction. Although generated from the same source data, the magnitude of the relative risk reductions and the corresponding outcome postponement values may seem contradictory. A key factor in understanding this apparent discrepancy is that outcomes rates are generally low in these trials. Consequently, a large relative risk reduction that represents a modest absolute risk reduction translates into a small outcome postponement.

Outcome postponement has been widely appraised as a tool for conveying the effect of preventive treatment to patients. It appears to be superior to NNT in some respects [Christensen 2003] (2). Patients show much more responsiveness when being informed by use of the outcome postponement than by other measures, i.e., their preferences with respect to accepting or rejecting a proposed drug

intervention changes when they are presented with different values for the postponement. For NNT, a similar proportion of patients accepts treatment, whether they are presented with values between 10 and 400 (2). It has been demonstrated that patients are more responsive to information conveyed through outcome postponement, i.e., their chance of accepting a treatment increases when they are presented with higher values of outcome postponement. On the other hand, even extreme differences in the presented values of NNT do not lead to greater or lower rates of treatment acceptance (2). A recent study randomized practices to use either absolute risk reduction or outcome postponement, when informing patients about the effect of statins. The study found a markedly lower proportion of patients redeeming their prescriptions if outcome postponement was used, even though the presented values reflected the same effect magnitude (12). A survey examined which threshold of absolute risk reduction and outcome postponement that would be required for the patients to accept preventive drugs. The lowest effect for patients to accept a hypothetical cholesterol-lowering drug varied, depending on their risk profile, from 20% to 30% in absolute risk reduction, and from 12 to 18 months on outcome postponement (13). Few, if any, preventive drugs can demonstrate such effects. Thus, there might be a gap between reality and patient expectations pertaining to the effect of preventive drugs.

The main limitation of using outcome postponement is the inability of the measure to capture the benefit of continued treatment in the decades after trial termination, although this is a limitation that applies to all effect measures currently used (7). The outcome postponement will continue to grow after trial termination as long as the survival curves are separated. Unfortunately, estimating the effect of continued statin treatment is difficult and depends heavily on untestable assumptions. For instance, some attempts to estimate the outcome postponement after trial termination assumed that the HR achieved during the trial will remain applicable after trial termination, and that patients will continue statin treatment with near-perfect adherence, something that has proven unrealistic in most drug utilization studies (14,15).

As an indicator of what a realistic outcome postponement for all-cause mortality could be for long-term statin treatment, we used the long-term follow-up on the WOSCOPS trial by Ford et al., demonstrating a hazard ratio for all-cause mortality of 0.87 (CI 95% 0.80-0.94). We calculated outcome postponement for the 20-year follow-up by using variables available in the paper, estimating an outcome postponement of 152 days (95% CI; 70-236 days) for 20 years. The patients were kept on placebo for the duration of the trial (6 years). Information of treatment after trial termination was not

provided (16). Given the age of typical statin initiators, 20 years seem like a realistic duration for statin treatment until end of life.

A large proportion of patients receive statins as secondary prevention initiated late in life, e.g., by having their first acute myocardial infarction in their eighties. Our results indicate that with their limited life expectancy in mind, these patients are not likely to gain a substantial OP, regardless of their adherence. Størring et al. estimated that lifelong statin treatment was associated with gains in survival ranging from 3 to 11 months (17).

A second limitation in our study is the varying trial duration between the trials. We have displayed the results both standardized and non-standardized. For the short-term trials, this implicates an assumption that the effect observed during the trial running time can be extrapolated to five years.

When assessing heterogeneity among the different outcomes, we found moderate to high heterogeneity. The degree of heterogeneity should be considered when evaluating the results. Moderate to high heterogeneity could indicate grouping of trials that do not examine the same biological effect, or that the patients included varies according to baseline characteristic e.g. age, previous ischaemic heart disease or treatment type. Our aim was to estimate the overall effect of statins on cardiovascular and non-cardiovascular outcomes; therefore, we have chosen to display the results even though there is a moderate to high heterogeneity. By standardizing according to trial duration, the level of heterogeneity drops in Table 2. Furthermore, as shown in E-figure 1A in the supplementary file, the level of heterogeneity decreases after subgrouping according to prevention type.

We envision three policy implications for the modelling of outcome postponement. Firstly, we suggest it should be calculated on all new trials to facilitate comparison of effects and incorporated into new guidelines. Secondly, it should be implemented in clinical practice when communicating treatment effects to patients. And finally, use of outcome postponement could support prescribers in withholding treatment for patients with a short life expectancy or those who have proven intolerant. Future analyses of subgroups within trials might identify patients with a particularly large or small outcome postponement and thereby make it possible to target the optimal patient population better.

In conclusion, statin treatment did postpone cardiovascular events, but the effect magnitude was small when measured in outcome postponement.

Competing interests

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: Dr. Hansen and Dr. Hallas reports grants from Pfizer, paid to our employer, outside the submitted work; Dr. Hansen received speaking honoraria from Novartis outside the submitted work; Dr. Pareek reports the following relationships – advisory board and speaking honoraria: AstraZeneca. The remaining authors have nothing to declare.

Contributions

All authors have made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data. They have all participated in drafting of the paper or revising it critically for important intellectual content. All authors have read and approved the final version of the manuscript.

Acknowledgements

We would like to thank the CTT Secretariat (especially Dr. Colin Baigent and Christina Reith) who helped obtain the mean follow-up time for the following trials; 4S, ASPEN. We thank Dr. Stuart Pocock from the ASPEN trial and Dr. Terje Pedersen from the 4S trial. We also thank Dr. Helen Colhoun from the CARDS trial, Dr. Donata Lucci from the GISSI-HF trial and Dr. Frank Sacks, Dr. Brian Claggett and Dr. Marc Pfeffer from the CARE trial, all of whom helped us access the mean follow-up time for the respective trials.

TABLE 1 Characteristics of the included trials. All were randomized trials comparing a statin vs placebo, ≥ 1000 patients, with a follow-up of minimum two years.

Trial name	Statin and daily dose	Type of prevention	Number of patients		Trial duration (years)	Baseline LDL-C (mmol/L) (8,18)	LDL-C difference at 1 year, compared to baseline (mmol/L) (8,18)
			Statin	Placebo			
ASCOT-LLA(19)	Atorvastatin (10 mg)	Mixed	5168	5137	3.5	3.44	-1.07
CARDS (20) *	Atorvastatin (10 mg)	Primary	1428	1410	4.75	3.03	-1.14
ASPEN(18)	Atorvastatin (10 mg)	Mixed	1211	1199	4.25	2.93	-.99
4D(21)	Atorvastatin (20 mg)	Mixed	619	636	6	3.25	-.89
SPARCL(22)	Atorvastatin (80 mg)	Secondary	2365	2366	6	3.44	NA
WOSCOPS(23)*	Pravastatin (40 mg)	Mixed	3302	3293	6	4.96	-1.07
CARE(24)*	Pravastatin (40 mg)	Secondary	2081	2078	5.5	3.58	-1.03
LIPID(25)*	Pravastatin (40 mg)	Secondary	4512	4502	6.63	3.88	-1.03
PROSPER(26)	Pravastatin (40 mg)	Mixed	2891	2913	3.7	3.79	-1.04
CORONA(27)	Rosuvastatin (10 mg)	Secondary	2497	2514	3	NA	NA
GISSI-HF(28)	Rosuvastatin (10 mg)	Mixed	2285	2289	4.5	3.06	-.92
AURORA(29)	Rosuvastatin (10 mg)	Mixed	1389	1384	5	2.58	-.99
JUPITER(30)	Rosuvastatin (20 mg)	Primary	8901	8901	4.5	2.7	-1.09
4S(31)	Simvastatin (10-40mg)	Secondary	2221	2223	5.8	4.88	-1.77
HPS(32)*	Simvastatin (40mg)	Mixed	10269	10267	6	3.38	-1.29
ALERT(33)*	fluvastatin (40 mg)	Mixed	1050	1052	6	4.14	-.84
LIPS(34)*	fluvastatin (80mg)	Secondary	844	833	4	3.42	-.92
AFCAPS(35)*	lovastatin (20-40 mg)	Primary	3304	3301	6	3.89	-.94
HOPE (36)*	rosuvastatin (10 mg)	Primary	6361	6344	7.46	3.31	-1.02

Abbreviations: LDL-C, Low-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; NA, Not available

*RR used instead of HR.

§ The LDL-C is the difference between the statin group and the placebo group at 1 year.

Table 2
Meta-Analyses of Postponement of 15 different Cardiovascular Outcomes.

Subgroup	Number of trials	Non-standardized Outcome postponement, days (95% CI)	I ² (%)	Outcome postponement standardized to five years, days (95% CI)	I ² (%)	HR-based meta-analysis, HR (95% CI)	I ² (%)
Cardiac death	4	11.6 (-0.8 – 24.0)	65	12.2 (-0.5 – 25.0)	71	0.77 (0.62 – 0.95)	16
Cardiovascular mortality	9	11.9 (4.0 – 19.7)	80	9.3 (3.6 – 14.9)	72	0.85 (0.77 – 0.95)	64
Death from CHD	4	11.9 (5.1 – 18.7)	56	10.1 (5.0 – 5.2)	40	0.76 (0.68 – 0.85)	0
Any myocardial infarction	15	19.8 (13.7 – 25.8)	90	18.0 (12.1 – 24.0)	92	0.71 (0.66, 0.76)	38
Non-fatal myocardial infarction	10	24.6 (14.6 – 34.7)	90	19.9 (13.5 – 26.4)	82	0.69 (0.61 – 0.78)	65
Any cardiovascular event	4	39.2 (16.8 – 61.5)	84	38.0 (15.5 – 60.5)	84	0.77 (0.70 – 0.84)	26

Any coronary event	5	20.7 (8.7 – 32.64)	82	21.7 (14.3 – 29.2)	46	0.73 (0.61 – 0.87)	75
Unstable angina	4	4.4 (-1.0 – 9.8)	71	3.1 (-0.7 – 6.8)	63	0.84 (0.74 – 0.95)	0
Coronary revascularization	12	21.9 (13.3 – 30.4)	91	16.5 (9.6 – 23.4)	92	0.72 (0.66, 0.78)	43
Non-fatal stroke	5	5.1 (2.4 – 7.7)	1	6.6 (3.5 – 9.8)	0	0.89 (0.73 – 1.10)	53
Fatal stroke	3	1.4 (-10.7 – 13.4)	81	-0.6 (-11.8 – 10.6)	75	1.19 (0.52 – 2.71)	82
Any stroke	14	6.6 (3.1 – 10.0)	63	6.1 (2.9 – 9.4)	66	0.85 (0.76 – 0.96)	62
Death from non- CV causes.	6	1.2 (-3.6 – 5.9)	0	1.5 (-2.2 – 5.3)	0	0.97 (0.90 – 1.05)	0
Hospitalizations for unstable angina	3	5.3 (-2.4 – 13.0)	76	8.1 (-2.8 – 19.1)	71	0.88 (0.81 – 0.95)	0
Hospitalizations for heart failure	3	5.0 (-1.5 – 11.6)	0	8.3 (-3.4 – 20.1)	0	0.94 (0.87 – 1.01)	0

Abbreviations: 95% CI: 95% confidence interval; HR: Hazard ratio; CV: cardiovascular; CHD: Coronary Heart Disease

TABLE 3A

Results of outcome postponement calculations from the mathematical model standardized to five years of trial duration, with 95% confidence interval.

Trial name	Cardiac death	Cardiovascular Death	Death from CHD	Any myocardial infarction	Non-fatal myocardial infarction	Any cardiovascular event	Any coronary event	Unstable angina
ASCOT-LLA	-	2.2 (-5.1 – 7.5)	-	14.1 (7.0 – 19.7)	-	-	19.3 (9.3 – 27.3)	8.0 (-34.3 – 31.5)
CARDS*	-	-	-	-	-	47.9 (22.2 – 67.8)	23.3 (5.8 – 35.8)	-
ASPEN	-	-	-	18.0 (-4.0 – 32.9)	-	-	-	-
4D	43.6 (-6.7 – 83.9)	-	-	15.8 (-27.2 – 48.0)	15.8 (-27.2 – 48.0)	-	-	-
SPARCL	0.0 (-8.5 – 5.5)	8.3 (-2.2 – 15.9)	-	-	15.5 (8.2 – 20.7)	59.4 (38.5 – 78.2)	32.2 (20.6 – 41.6)	-
WOSCOPS*	-	6.5 (0.6 - 10.8)	5.6 (0.2 – 9.4)	23.0 (11.8 – 31.8)	18.0 (8.0 – 26.7)	-	-	-
CARE*	-	-	10.5 (-2.6 – 20.6)	27.8 (10.3 – 41.9)	22.7 (5.9 – 35.7)	-	-	19.1 (-1.5 – 37.0)
LIPID*	-	17.2 (8.9 – 24.1)	14.3 (7.1 – 20.9)	26.4 (16.5 – 35.4)	-	-	-	-
PROSPER	-	-	13.8 (0.6 – 24.2)	29.5 (9.3 – 48.6)	16.0 (-3.4 – 32.2)	32.4 (6.4 – 54.5)	-	-
CORONA	-	9.7 (-28.4 – 42.5)	-	-	-	-	26.7 (-13.1 – 61.0)	-
GISSI-HF	-	9.1 (-20.3 – 34.7)	-	4.3 (-10.0 – 14.4)	-	-	-	-
AURORA	-	0.0 (-41.2 – 40.0)	-	16.9 (-11.5 – 38.4)	16.9 (-11.5 – 38.4)	-	-	-
JUPITER	-	-	-	9.0 (5.0 – 11.7)	9.7 (6.3 – 11.7)	-	-	-
4S	-	27.2 (15.5 – 37.5)	-	55.4 (40.2 – 69.3)	55.4 (40.2 – 69.3)	-	-	-
HPS*	-	-	-	18.8 (14.8 – 22.8)	18.8 (14.8 – 22.8)	-	-	-
ALERT*	17.0 (1.8 – 26.9)	-	-	17.4 (0.0 – 32.7)	17.4 (0.0 – 32.7)	-	-	-
LIPS*	15.6 (-1.6 – 24.2)	-	-	-	-	-	-	-
AFCAPS*	-	-	-	10.0 (4.2 – 14.2)	-	16.2 (5.8 – 24.8)	13.8 (4.4 – 21.6)	7.3 (1.1 – 11.7)
HOPE	-	2.2 (-2.2 – 5.6)	-	2.1 (0.4 – 3.3)	-	-	-	1.0 (-1.8 – 2.9)

Abbreviations: CHD, Coronary heart disease.

*RR used instead of HR.

Author Manuscript

TABLE 3B

Results of Outcome Postponement Calculations from the Mathematical Model standardized to Five Years of Trial Duration 95% Confidence Interval.

Trial name	Coronary revascularisation	Non-fatal stroke	Fatal stroke	Any stroke	Death from non- CV cause	Hospitalizations for unstable angina	Hospitalizations for heart failure
ASCOT-LLA	-	-	-	8.9 (1.3 – 14.6)	-	-	-
CARDS*	6.6 (-3.4 – 12.5)	-	-	16.1 (3.7 – 23.2)	-	-	-
ASPEN	-	-	-	-	-	-	-
4D	-	-2.3 (-38.4 – 20.5)	-24.0 (-67.2 – -1.2)	-25.0 (-72.0 – 7.7)	11.9 (-41.7 – 58.0)	-	-

SPARCL	27.9 (17.3 – 35.4)	13.3 (-3.0 – 27.8)	6.9 (0.8 – 10.4)	18.1 (1.1 – 32.9)	-	-	-
WOSCOPS*	8.3 (2.5 – 12.5)	-	-	1.6 (-4.7 – 5.7)	1.9 (-4.8 – 6.6)	-	-
CARE*	43.2 (23.8 – 59.6)	-	-	10.8 (1.0 – 18.2)	-	-	-
LIPID*	21.8 (10.9 – 30.6)	-	-	6.3 (0.0 – 11.3)	-	19.5 (6.5 – 31.0)	-
PROSPER	4.2 (-6.0 – 10.7)	1.1 (-14.4 – 13.5)	-3.9 (-14.1 – 1.4)	-1.8 (-18.8 – 11.7)	-7.6 (-26.0 – 7.6)	-	5.2 (-10.3 – 16.7)
CORONA	-	-	-	-	-	4.7 (-14.1 – 18.0)	34.5 (-7.5 – 70.2)
GISSI-HF	-	-	-	-8.4 (-25.4 – 4.0)	-	-	8.4 (-24.8 – 36.8)
AURORA	2.9 (-33.1 – 32.5)	-	-	-8.2 (-35.8 – 10.2)	18.1 (-20.1 – 52.9)	-	-
JUPITER	-	6.7 (2.8 – 9.4)	-	7.4 (3.2 – 10.2)	-	2.6 (-0.6 – 4.3)	-
4S	51.2 (35.7 – 63.9)	-	-	-	-	-	-
HPS*	18.6 (13.6 – 23.6)	-	-	12.6 (7.5 – 17.1)	2.4 (-3.4 – 7.3)	-	-
ALERT*	6.2 (-9.2 – 15.8)	-	-	-8.2 (-32.1 – 8.8)	-10.6 (-35.1 – 7.5)	-	-
LIPS*	-	-	-	-	-	-	-
AFCAPS*	13.4 (6.1 – 19.6)	-	-	-	-	-	-
HOPE	2.3 (0.4 – 3.8)	-	-	2.7 (0.5 – 4.4)	-	-	-

Abbreviations: CHD: Coronary heart disease; CV: Cardiovascular

*RR used instead of HR.

Figure 1. Flowchart of Included Studies

Figure 2. Forest Plots of Postponement of Cardiovascular Mortality Standardized to Five Year of Trial Duration, grouped according to Prevention Type.

Abbreviations: Sec, secondary prevention; Mix, mixed prevention.

Author Manuscript

Figure 3. Forest Plots of Postponement of Death from Non-Vascular and Non-Cardiovascular Mortality Standardized to Five Year of Trial Duration, grouped according to Prevention Type.

Abbreviations: Mix, mixed prevention.

*RR used instead of HR.

Author Manuscript

Figure 4. Forest Plots of Postponement of Any Myocardial Infarction, Standardized to Five Year of Trial Duration, grouped according to Prevention Type.

Abbreviations: Pri, Primary prevention; Sec, secondary prevention; Mix, mixed prevention.

*RR used instead of HR.

Figure 5. Forest Plots of Postponement of Any Stroke, Standardized to Five Year of Trial Duration, grouped according to Prevention Type.

Abbreviations: Pri, primary prevention; Sec, secondary prevention; Mix, mixed prevention.

*RR used instead of HR.

Author Manuscript

References.

1. Taylor FC, Huffman M, Ebrahim S. Statin therapy for primary prevention of cardiovascular disease. *JAMA*. 2013 Dec 11;310(22):2451–2.
2. Christensen PM, Brosen K, Brixen K, Andersen M, Kristiansen IS. A randomized trial of laypersons' perception of the benefit of osteoporosis therapy: Number needed to treat versus postponement of hip fracture. *Clin Ther*. 2003;25(10):2575–2585.
3. Hux JE, Naylor CD. Communicating the Benefits of Chronic Preventive Therapy Does the Format of Efficacy Data Determine Patients' Acceptance of Treatment? *Med Decis Making*. 1995;15(2):152–157.
4. Malenka DJ, Baron JA, Johansen S, Wahrenberger JW, Ross JM. The framing effect of relative and absolute risk. *J Gen Intern Med*. 1993;8(10):543–548.
5. Laupacis A, Sackett DL, Roberts RS. An assessment of clinically useful measures of the consequences of treatment. *N Engl J Med*. 1988 Jun 30;318(26):1728–33.
6. Wright JC, Weinstein MC. Gains in life expectancy from medical interventions—standardizing data on outcomes. *N Engl J Med*. 1998;339(6):380–386.
7. Hansen MR, Hróbjartsson A, Pottegård A, Damkier P, Larsen KS, Madsen KG, et al. Postponement of Death by Statin Use: a Systematic Review and Meta-analysis of Randomized Clinical Trials. *J Gen Intern Med [Internet]*. 2019 May 9 [cited 2019 May 24]; Available from: <http://link.springer.com/10.1007/s11606-019-05024-4>
8. Trialists CT, others. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170 000 participants in 26 randomised trials. *The Lancet*. 2010;376(9753):1670–1681.
9. Kenneth J. Rothman. *Epidemiology: An Introduction*. 2nd ed.
10. Julian PT Higgins and Sally Green. *Cochrane Handbook for Systematic Reviews of Interventions*. Vol. 2008.
11. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JPA, et al. The PRISMA Statement for Reporting Systematic Reviews and Meta-Analyses of Studies That Evaluate Health Care Interventions: Explanation and Elaboration. *PLoS Med*. 2009 Jul 21;6(7):e1000100.
12. Harmsen CG, Kristiansen IS, Larsen PV, Nexoe J, Stovring H, Gyrd-Hansen D, et al. Communicating risk using absolute risk reduction or prolongation of life formats: cluster-randomised trial in general practice. *Br J Gen Pract*. 2014 Apr 1;64(621):e199–207.
13. Trewby PN, Reddy AV, Trewby CS, Ashton VJ, Brennan G, Inglis J. Are preventive

drugs preventive enough? A study of patients' expectation of benefit from preventive drugs. *Clin Med.* 2002;2(6):527–533.

14. Shah ND, Dunlay SM, Ting HH, Montori VM, Thomas RJ, Wagie AE, et al. Long-term Medication Adherence after Myocardial Infarction: Experience of a Community. *Am J Med.* 2009 Oct;122(10):961.e7-961.e13.
15. Chan DC, Shrank WH, Cutler D, Jan S, Fischer MA, Liu J, et al. Patient, physician, and payment predictors of statin adherence. *Med Care.* 2010;48(3):196–202.
16. Ford I, Murray H, Packard CJ, Shepherd J, Macfarlane PW, Cobbe SM. Long-term follow-up of the West of Scotland Coronary Prevention Study. *N Engl J Med.* 2007 Oct 11;357(15):1477–86.
17. Stovring H, Harmsen CG, Wisloff T, Jarbol DE, Nexoe J, Nielsen JB, et al. A competing risk approach for the European Heart SCORE model based on cause-specific and all-cause mortality. *Eur J Prev Cardiol.* 2013 Oct 1;20(5):827–36.
18. Knopp RH. Efficacy and Safety of Atorvastatin in the Prevention of Cardiovascular End Points in Subjects With Type 2 Diabetes: The Atorvastatin Study for Prevention of Coronary Heart Disease Endpoints in Non-Insulin-Dependent Diabetes Mellitus (ASPEN). *Diabetes Care.* 2006 Jul 1;29(7):1478–85.
19. Sever PS, Dahlof B, Poulter NR, Wedel H, Beevers G, Caulfield M, et al. Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial--Lipid Lowering Arm (ASCOT-LLA): a multicentre randomised controlled trial. *Lancet.* 2003 Apr 5;361(9364):1149–58.
20. Colhoun HM, Betteridge DJ, Durrington PN, Hitman GA, W Neil HA, Livingstone SJ, et al. Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial. *The Lancet.* 2004;364(9435):685–696.
21. Wanner C, Krane V, März W, Olschewski M, Mann JF, Ruf G, et al. Atorvastatin in patients with type 2 diabetes mellitus undergoing hemodialysis. *N Engl J Med.* 2005;353(3):238–248.
22. Amarenco P, Bogousslavsky J, Callahan A 3rd, Goldstein LB, Hennerici M, Rudolph AE, et al. High-dose atorvastatin after stroke or transient ischemic attack. *N Engl J Med.* 2006 Aug 10;355(6):549–59.
23. Shepherd J, Cobbe SM, Ford I, Isles CG, Lorimer AR, Macfarlane PW, et al. Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia. *N Engl J Med.*


1995;333(20):1301–1308.

24. Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. *N Engl J Med.* 1996;335(14):1001–1009.
25. Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. The Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group. *N Engl J Med.* 1998 Nov 5;339(19):1349–57.
26. Shepherd J, Blauw GJ, Murphy MB, Bollen ELEM, Buckley BM, Cobbe SM, et al. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): a randomised controlled trial. *Lancet.* 2002 Nov 23;360(9346):1623–30.
27. Kjekshus J, Apetrei E, Barrios V, Böhm M, Cleland JG, Cornel JH, et al. Rosuvastatin in older patients with systolic heart failure. *N Engl J Med.* 2007;357(22):2248–2261.
28. GISSI-HF Investigators, Tavazzi L, Maggioni AP, Marchioli R, Barlera S, Franzosi MG, et al. Effect of rosuvastatin in patients with chronic heart failure (the GISSI-HF trial): a randomised, double-blind, placebo-controlled trial. *Lancet.* 2008 Oct 4;372(9645):1231–9.
29. Fellstrom BC, Jardine AG, Schmieder RE, Holdaas H, Bannister K, Beutler J, et al. Rosuvastatin and cardiovascular events in patients undergoing hemodialysis. *N Engl J Med.* 2009 Apr 2;360(14):1395–407.
30. Ridker PM, Danielson E, Fonseca FA, Genest J, Gotto Jr AM, Kastelein JJ, et al. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med.* 2008;359(21):2195.
31. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S). *Lancet.* 1994 Nov 19;344(8934):1383–9.
32. Heart Protection Study Collaborative Group. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet.* 2002 Jul 6;360(9326):7–22.
33. Holdaas H, Fellström B, Jardine AG, Holme I, Nyberg G, Fauchald P, et al. Effect of fluvastatin on cardiac outcomes in renal transplant recipients: a multicentre, randomised, placebo-controlled trial. *The Lancet.* 2003;361(9374):2024–2031.
34. Serruys PWJC, de Feyter P, Macaya C, Kokott N, Puel J, Vrolix M, et al. Fluvastatin for prevention of cardiac events following successful first percutaneous coronary intervention: a randomized controlled trial. *JAMA J Am Med Assoc.* 2002 Jun 26;287(24):3215–22.
35. Downs JR, Clearfield M, Weis S, Whitney E, Shapiro DR, Beere PA, et al. Primary

prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: results of AFCAPS/TexCAPS. *Jama*. 1998;279(20):1615–1622.


36. Yusuf S, Lonn E, Pais P, Bosch J, López-Jaramillo P, Zhu J, et al. Blood-Pressure and Cholesterol Lowering in Persons without Cardiovascular Disease. *N Engl J Med*. 2016 May 26;374(21):2032–43.

Author Manuscript



Author Manuscript

This article is protected by copyright



Trial_name

ES (95% CI)

Weight

