

## The association between cannabis use and testicular function in men

### A systematic review and meta-analysis

Belladelli, Federico; Del Giudice, Francesco; Kasman, Alex; Kold Jensen, Tina; Jørgensen, Niels; Salonia, Andrea; Eisenberg, Michael L.

*Published in:*  
Andrology

*DOI:*  
10.1111/andr.12953

*Publication date:*  
2021

*Document version:*  
Accepted manuscript

#### *Citation for published version (APA):*

Belladelli, F., Del Giudice, F., Kasman, A., Kold Jensen, T., Jørgensen, N., Salonia, A., & Eisenberg, M. L. (2021). The association between cannabis use and testicular function in men: A systematic review and meta-analysis. *Andrology*, 9(2), 503-510. <https://doi.org/10.1111/andr.12953>

Go to publication entry in University of Southern Denmark's Research Portal

#### **Terms of use**

This work is brought to you by the University of Southern Denmark.  
Unless otherwise specified it has been shared according to the terms for self-archiving.  
If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim.  
Please direct all enquiries to [puresupport@bib.sdu.dk](mailto:puresupport@bib.sdu.dk)

DR FEDERICO BELLADELLI (Orcid ID : 0000-0002-2165-659X)

ALEX KASMAN (Orcid ID : 0000-0003-0523-8176)

DR NIELS JØRGENSEN (Orcid ID : 0000-0003-4827-0838)

PROFESSOR ANDREA SALONIA (Orcid ID : 0000-0002-0595-7165)

Article type : Review Article

## **The association between cannabis use and testicular function in men: a systematic review and meta-analysis**

Federico Belladelli<sup>a,b,c</sup>, Francesco Del Giudice<sup>c,d</sup>, Alex Kasman<sup>c</sup>, Tina Kold Jensen<sup>e</sup>, Niels Jørgensen<sup>f</sup>, Andrea Salonia<sup>a</sup>, Michael L Eisenberg<sup>c</sup>

<sup>a</sup> Division of Experimental Oncology/Unit of Urology; URI; IRCCS Ospedale San Raffaele, Milan, Italy.

<sup>b</sup> University Vita-Salute San Raffaele, Milan, Italy.

<sup>c</sup> Department of Urology, School of Medicine, Stanford University, Stanford, California.

<sup>d</sup> Department of Urology, University Sapienza, Rome, Italy.

<sup>e</sup> Department of Environmental Health, University of Southern Denmark, Odense, Denmark.

<sup>f</sup> Department of Growth and Reproduction, International Center for Research and Research Training in Endocrine Disruption of Male Reproduction and Child Health, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark.

Word Count: 3006

### CORRESPONDING AUTHOR:

Michael L. Eisenberg

**This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as [doi: 10.1111/ANDR.12953](https://doi.org/10.1111/ANDR.12953)**

This article is protected by copyright. All rights reserved

Director, Male Reproductive Medicine and Surgery  
Associate Professor, Department of Urology and Obstetrics & Gynecology  
Stanford University School of Medicine  
Tel. +1 650.725.5746; Fax +1 650.498.5346  
Email: eisenberg@stanford.edu

## ABSTRACT

**Objective:** To evaluate the association between cannabis use and testicular function (as assessed through semen quality and serum hormone levels) in different populations.

**Evidence Review:** Systematic review and meta-analysis of population-based retrospective cohort studies. PRISMA guidelines were used for abstracting data and assessing data quality and validity. Data were pooled using a fixed-effect or random-effects model depending on the heterogeneity of studies included. Pooled risk ratio (RR) of having any sperm abnormality and Testosterone, FSH, and LH standardized mean differences among male cannabis users and non-users, and meta-regression analysis according to age, and year of publication.

**Results:** Nine studies were evaluated which included 4014 men with semen data and 4787 with hormonal data. Overall among 1158 cannabis users, 44.9% had impaired semen parameters. Compared to 24.5% of the 2856 nonusers. The relative risk among cannabis users for any abnormal semen parameter was 1.159 (95%CI: 0.840; 1.599,  $p=0.369$ ). The standardized mean difference between users and non-users testosterone levels was -0.139 (95%CI: -0.413; 0.134,  $p=0.318$ ). For FSH, the standardized mean difference estimate was -0.142 (95%CI: -0.243; -0.0425,  $p=0.005$ ), while for LH the standardized mean difference estimate was -0.318 (95%CI: -0.810-0.175;  $p=0.206$ ).

**Conclusions:** The current evidence does not suggest clinically significant associations between cannabis use and testicular function. However, we cannot exclude an effect of cannabis due to the limited and heterogeneous studies. Additional, well-designed studies will be needed to define the association between cannabis use and the male reproductive system.

**Keywords:** Sperm, Male Infertility, Cannabis, Testosterone, FSH, LH

## 1. INTRODUCTION

Approximately 15% of couples are not able to achieve pregnancy after 1 year of trying and are classified as infertile (1). Within couples, male factor infertility contributes roughly 50% of cases (2). Cannabis use has been implicated in impaired fertility; however, its effect on semen parameters and reproductive hormone profiles remains uncertain (3,4).

Cannabis is one of the most commonly used drugs and is becoming increasingly legalized within the United States and worldwide. The number of US cannabis users increased by 60% between 2007 and 2017 and almost 10 million are daily or near-daily users. Roughly 3.8 per cent of the global population aged 15–64 years used cannabis at least once in 2017 (5). To date, in the United States, thirty-three states and the District of Columbia have passed laws broadly legalizing cannabis (6) and other countries are following.

The active component of cannabis is D9-tetrahydrocannabinol (THC), which binds cannabinoid receptors present in the brain, the nervous system and, among other locations, the reproductive organs (7). There has been conflicting literature regarding the effect of cannabis on the reproductive system; however, recent reviews and studies using both animal and human systems have suggested that cannabis impairs male fertility, semen quality and hormones levels (8,9,10,11,12,13). In contrast, others have reported no changes in semen parameters and testosterone levels among cannabis users (14,15,16,17). In addition, many studies have been conducted among men with very high intake and are therefore may not be applicable to all users.

Since most of cannabis consumers are males of reproductive age (18) and prior literature is heterogeneous regarding the association between cannabis use and testicular function, it is important to further investigate. The aim of this systematic review and meta-analysis is to critically evaluate the literature for the association between cannabis use and testicular function (i.e. semen parameters, testosterone, and gonadotropins).

## **2. MATERIAL AND METHODS**

This meta-analysis was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. The following research question was established based on the PICO criteria (19): what is the association in men of smoking cannabis and testicular function as measured by semen quality, testosterone levels, and gonadotropins? Furthermore, our goal was to compare current evidence within available studies.

### **2.1 Evidence acquisition**

We performed a systematic review of the literature in PubMed, Embase, and Cochrane from inception to January 2020, to identify studies that evaluated cannabis use, semen analysis, and hormonal profiles. Search terms included: “marijuana/cannabis and semen parameters”, “marijuana/cannabis and sperm concentrations”, “marijuana/cannabis and sperm count”, “marijuana/cannabis and sperm morphology”, “marijuana/cannabis and testosterone”, “marijuana/cannabis and FSH levels”, “marijuana/cannabis and LH levels”; secondary fields: male general health; male infertility; male hypogonadism; male factor infertility; infertility.

The reference lists of the included studies were also screened for relevant articles. Nine original population-based retrospective cohort studies, one original prospective population-based cohort study and two cross sectional study were included and critically evaluated (Level of Evidence: III-2)

## **2.2 Selection of the studies and criteria of inclusion**

This analysis was restricted to data collected from original articles that examined men using cannabis presenting with semen analysis and/or hormonal profiles (testosterone, FSH/LH). Studies were considered eligible if an internal or external control populations enrolled were defined by age-matched men (age>18 years old) who had never consumed cannabis also presenting with semen analysis and/or hormonal profiles (testosterone, FSH/LH).

Articles were excluded if they met one or more of the following criteria: presence of men who have undergone testosterone replacement or other gonadotoxic hormonal treatments. Case-control cohort studies, case reports, abstracts and meeting reports were excluded from the analysis.

Two authors (FB and FDG) independently screened the titles and abstracts of all articles. The full-text articles were examined independently by three authors (FB, FDG, and AK) to determine whether or not they met the inclusion criteria. Final inclusion was determined by the consensus of all investigators. Selected articles meeting the inclusion criteria were then critically analyzed.

The following data were extracted from the included studies by using a standardized form: country of origin, publication year, sample size, participants age, number of cannabis smokers, numbers of control, population description and main findings of the study.

## **2.3 Assessment of quality for studies included and Statistical Analysis**

To assess the risk of bias (RoB) each report was reviewed using the NIH Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies (20). The authors independently assessed the methodological quality based on sequence generation, allocation concealment, enrollment of control groups, incomplete outcome data, selective outcome reporting, and

additional sources of bias. Publication bias was tested by visual assessment of the Deeks' funnel plot (21). We compared effects on subfertile semen parameters using pooled relative risk and 95% Confidence Intervals (CI). Variability in the intervention effects as a consequence of clinical or methodological diversity, for example the reason for semen analysis, among the studies was evaluated in form of heterogeneity (22). According to heterogeneity assessment, the pooled relative risk estimate was calculated using a fixed or random-effects model (23). We compared effects on total testosterone, LH and FSH using the pooled standardized mean difference estimate as the studies we included presented their results in terms of average hormone levels. Our results are graphically displayed as forest plots, with pooled relative risk and standardized mean difference indicating overall risk for cannabis using men to present with impaired semen and hormonal analyses respectively. Evaluation for presence of heterogeneity was done using (24): (1) Cochran's Q-test with  $p < 0.05$  signifying heterogeneity; (2) Higgins  $I^2$  test with inconsistency index ( $I^2$ ) = 0%–40%, heterogeneity might not be important; 30%–60%, moderate heterogeneity; 50%–90%, substantial heterogeneity; and 75%–100%, considerable heterogeneity. Sensitivity analysis with and without men recruited in fertility clinic were performed for testosterone, FSH, and LH in order to investigate possible bias but no differences was observed. The number of studies investigating semen parameters in men not recruited in a fertility clinic was too small to perform a sensitivity analysis.

Metaregression was performed to explore potential bias to the overall effects, and the proportion of the studies was assessed by a random-effects regression model. Calculations were accomplished using the Comprehensive Meta-Analysis Software, version v.2 .0 (CMA, Biostat, Englewood, NJ, USA).

### **3. RESULTS**

#### **3.1 Search results**

The initial search yielded 274 articles (PubMed: 208; Cochrane: 55; and Embase: 11). 183 were excluded as they were duplicates appearing in multiple databases. Of the remaining 91, 76 were further excluded as they either did not examine cannabis effects on conventional semen parameters (34), contained animal experiments (17) or were review papers or editorials (10). Full-text articles were then reevaluated and critically analyzed for the

remaining 15 articles. After another in-depth review, a further six did not meet the inclusion criteria. The remaining nine studies were included in our review (Fig. 1); four of them were used in the semen parameters analysis, six for the testosterone analysis and four for the gonadotropin analysis. RoB assessment according to NIH Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies for each of the individual studies is illustrated in eFig. 1.

### **3.2 Study location, types, and populations**

The patient description, main findings, and study characteristics of each article are summarized in Table 1. Of the nine studies included, five were conducted in the US (9,14,15,16,17), three in Europe (Denmark, UK, and Italy) (10,11,12), and one in Jamaica (13). Four studies investigated only hormone levels in a cannabis user group versus a non-user group (9,10,14,19). Five studies recruited men evaluated at an infertility clinic (11,13,17,18,20). One study examined semen parameters and hormone profile in cannabis users who participated in the Danish military draft (12), while two studies analyzed data from male participants of the U.S. National Health and Nutrition Examination Survey (NHANES) (15,19). The final study recruited physically active men from a university campus (16).

### **3.3 Study sample sizes, participant ages, and meta regression results**

In total, 520 (44.9%) men had impaired semen parameters among 1158 cannabis users. Among 2856 nonusers, 699 (24.5%) men had an impaired semen analysis which was defined as the presence of any abnormal parameter according to the WHO 5th edition reference values. A man was then defined as having “normal” sperm if all values were within the normal reference range or “subfertile” sperm if any value was abnormal. With regard to testosterone levels, there were a total of 1717 cannabis users and 1458 nonusers while LH and FSH levels were assessed in 735 cannabis users and 823 nonusers. Detailed information regarding the frequency/duration of cannabis use was not consistently available and thus a sub-analysis could not be performed. Meta regression analyses were performed in order to investigate the association of moderator variables on study effect size using regression-based techniques. The regression results showed that the year of publication affected overall effects of cannabis use on semen parameters (slope estimate: -0.129; 95% CI: -0.190, -0.069;  $P <$



0.001) and testosterone levels (slope estimate: 0.026; 95% CI: 0.003, 0.048;  $P=0.026$ ), while age affected semen parameters (slope estimate: -0.021; 95% CI: -0.033, -0.010;  $P < 0.001$ ) and FSH levels (slope estimate: -0.019; 95% CI: -0.032, -0.005;  $P = 0.007$ ) (eFig. 2).

### **3.4 Association of cannabis use and semen parameters**

The six studies analyzing semen parameters reported the association of any abnormality (using WHO 5th edition reference values) with cannabis consumption with a range of RR estimates from 0.829 to 1.502. The pooled relative risk for any abnormal semen parameter under a random-effects model was: 1.159 (95%CI: 0.840;1.599,  $p=0.369$ ) (Fig. 2A) with evidence for heterogeneity between the studies ( $Q= 19.4813$  (d.f.=3),  $p=0.0002$ ;  $I^2= 84.60\%$ ). A sub-analysis of the three studies with morphology data (i.e. men with  $< 4\%$  of morphologically normal sperm) was performed (Fig. 3B). The pooled relative risk of abnormal morphology with cannabis use under a random-effects model was: 0.899 (95%CI: 0.557;1.451,  $p=0.663$ ) with a considerable heterogeneity between the studies:  $Q=12.9128$  (d.f.=2),  $p=0.0016$ ;  $I^2= 84.51\%$ .

### **3.5 Association of cannabis use and total serum testosterone concentration**

The eight studies analyzing testosterone levels in relation to cannabis consumption had standardized mean difference estimates from -2.259 to 0.192. The pooled standardized mean difference estimate under a random-effects model was -0.139 (95%CI: -0.413;0.134,  $p=0.318$ ) (Fig. 3) with evidence of heterogeneity between the studies ( $Q= 39.1372$  (d.f.=5),  $p < 0.0001$ ;  $I^2= 87.22\%$ ). Inspection of the funnel plot suggests that there was a small-study effect with Kolodny et al's analysis tending to have smaller standardized mean difference estimates. Therefore, we developed a second analysis without this particular study which revealed a pooled standardized mean difference estimate under a fixed effect model of 0.0999 (95%CI: 0.0288-0.171) ( $p = 0.006$ ) without significant heterogeneity between the studies ( $Q= 4.6084$  (d.f.=4),  $p=0.3299$ ;  $I^2= 13.20$ ).

### **3.6 Association of cannabis use and serum FSH and LH concentrations**

The five studies analyzing serum FSH and LH levels in relation to cannabis consumption reported standardized mean difference estimates from -3.207 to 0.136 for LH and -0.718 to -0.0000000160 for FSH. For FSH, the pooled standardized mean difference estimate under a fixed-effect model was -0.142 (95%CI: -0.243; -0.0425,  $p=0.005$ ) with an absence of considerable heterogeneity between the studies ( $Q= 10.3188$  (d.f.=3),  $p= 0.0160$ ;  $I^2= 70.93\%$ ) (Fig. 4A). For LH, the pooled standardized mean difference estimate under a random-effects model was -0.318 (95%CI: -0.810-0.175;  $p= 0.206$ ) with considerable heterogeneity between the studies ( $Q= 28.3352$  (d.f.=3),  $p< 0.0001$ ;  $I^2= 89.41\%$ ) (Fig. 4B).

#### 4. DISCUSSION

The present meta-analysis did not identify an association between cannabis use and testicular function. Specifically, the current analysis was unable to demonstrate impairment of semen quality in general, or on morphology, specifically with cannabis use. With regard to the association of cannabis on reproductive hormones, total testosterone levels did not differ significantly between cannabis users and non-users. LH levels were not different between cannabis users and non-users, while FSH levels were lower in cannabis users compared to non-users. While the effects of cannabis on FSH levels were statistically significant, the changes were quantitatively small making the clinical significance uncertain. By meta-

regression analysis, it was found that the year of publication was associated with the relative risk of cannabis use and impaired semen parameters as well as the standardized mean differences in testosterone levels in cannabis users and non-users suggesting a temporal trend in the reported associations. Importantly, the lack of any negative associations between cannabis use and semen quality does not imply evidence of safety.

The existing literature is heterogenous in describing the associations of cannabis on male fertility. In particular, current studies appear to be conflicting with several demonstrating a possible link between cannabis consumption and abnormal semen analysis while others do not. Pacey et al, compared 318 cannabis users to 1652 non-users and demonstrated that cannabis use is a risk factor for poor sperm morphology (11). Gundersen et al examined 1215 men, median 19 years old, from the general population, and thus were unselected regarding testicular function. They detected that cannabis use was associated with lower sperm concentration and total sperm count (12). Similarly, Carrol and colleagues reported that cannabis users examined in a fertility clinic were at greater risk of being diagnosed with asthenozoospermia and teratozoospermia (13). In contrast, Nassan et al showed that cannabis users among men from infertile couples had significantly higher sperm concentrations (17).

The literature is similarly discordant regarding the effect of cannabis use on total testosterone levels. Most studies in our analyses suggest that there is no difference in testosterone levels between cannabis users and non-users; however, several showed a significant effect (14,15,16). Two studies suggested that cannabis users had higher testosterone levels (12,15). In contrast, Kolodny et al, reported a decrease in mean total testosterone levels in subjects using cannabis (9). This study of forty men showed a small study effect evident in the funnel plot and different results compared to the other studies on the subject.

The association between cannabis and gonadotropins is also mixed in the literature with studies showing positive, negative, and absent associations. The populations studied also varied. While most studies analyzed young men (18-37 years old), two studies examined men up to the age of 50 (11,15). Populations were also heterogenous with regards to recruitment. Three studies recruited men from fertility clinics (11,13,17) and two studies recruited men from college campuses complicating comparisons between reports (14,16). The rate of cannabis use in our study is different from the one reported by 2019 WHO Drug report (41.3% vs 3.9%) as most of the studies we included analyzed similar numbers of users

and non-users (case control design) to better compare testicular function in the two populations.

As the clinical utility of semen values have been questioned (25,26), cannabis use has also been investigated in relation to fecundability. Kasman et al examined a representative sample of US men and women and found no significant association between cannabis use and time to pregnancy across all cannabis user groups including daily smokers (27). Wise et al analyzed data from couples attempting to conceive and found male cannabis consumption at levels of  $\geq 1$  time/week was associated with an increase in fecundability (28). Nassan et al followed 421 women who underwent 730 ART cycles and evaluated the association of baseline cannabis smoking with ART outcomes. 200 of these women (368 cycles) were part of a couple in which the male partner's cannabis use was also studied. No statistically significant differences have been showed in the probabilities of implantation, clinical pregnancy, or live birth according to women's cannabis use. In contrast, when the male partner reported cannabis use at enrollment, the couple had a significantly higher probability of live birth, independent of the women's cannabis smoking status (29). Given the increasing legalization of cannabis worldwide, particularly in the US, this subject requires further research in regard to both laboratory and clinical endpoints. This is particularly true for those undergoing IVF, as Domar et al found many infertility patients do not follow recommendations on lifestyle habits while undergoing IVF such as utilizing cannabis (30).

Several limitations warrant mention. First, cannabis use profiles were not reported in most studies which limits the interpretability of the results as the differences in frequency and quantity of cannabis use are common (31). Such variability in the profiles of cannabis use also prevented the examination of a dose-dependent effect of cannabis use. In addition, while we used a categorization of semen quality according to the WHO reference levels, we cannot exclude the possibility of a difference between users and non-users of cannabis if we analyzed the actual values of semen variables. Additionally, the absence of pregnancy outcomes limits the ability to assess for the most important reproductive clinical endpoints of cannabis use. Furthermore, recruitment methods may lead to bias, either related to age or recruitment location (e.g. universities versus fertility clinics). Finally, all studies utilized self-reported cannabis use which may be not reliable due to the social stigma or fear of repercussions. However, the current literature does suggest the validity of survey methodology (32).

## 5. CONCLUSION

Currently, the number and quality of the studies focusing on cannabis and reproductive and sexual health remain limited. However, our systematic review and meta-analysis suggests a negligible clinical effect of cannabis use on testicular function. Due to the low number of studies and the heterogeneity of the existing studies, we cannot exclude a potential effect of cannabis on testicular function and the current analysis does not prove safety. Future studies in diverse populations with detailed information on cannabis use are needed to further examine the association of cannabis and male reproductive health.

## 6. ACKNOWLEDGMENT SECTION

The authors have no conflicts of interest to disclose.

## REFERENCES

1. World Health Organization. WHO Manual for the Standardized Investigation and Diagnosis of the Infertile Couple. Cambridge: Cambridge University Press; 2000.
2. Boivin J, Bunting L, Collins JA, Nygren KG. International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care. *Hum Reprod* 2007; 22: 1506–12.
3. Rajanahally S, Raheem O, Rogers M, et al. The relationship between cannabis and male infertility, sexual health, and neoplasm: a systematic review. *Andrology*. 2019;7(2):139–147.
4. Brents LK. Marijuana, the Endocannabinoid System and the Female Reproductive System. *Yale J Biol Med*. 2016;89(2):175–191.
5. United Nations Office on Drugs and Crime, World Drug Report 2019

6. Governing: State Marijuana Laws in 2019 Map. Available at <http://www.governing.com/gov-data/safety-justice/state-marijuana-laws-map-medical-recreational.html>. Accessed Jan 08, 2020.
7. Rossato M, Pagano C, Vettor R. The cannabinoid system and male reproductive functions. *J Neuroendocrinol*. 2008;20 Suppl 1:90–93.
8. Payne KS, Mazur DJ, Hotaling JM, Pastuszak AW. Cannabis and Male Fertility: A Systematic Review. *J Urol*. 2019;202(4):674–681.
9. Kolodny RC, Masters WH, Kolodner RM, Toro G. Depression of plasma testosterone levels after chronic intensive marihuana use. *N Engl J Med*. 1974;290(16):872–874.
10. Vescovi PP, Pedrazzoni M, Michelini M, Maninetti L, Bernardelli F, Passeri M. Chronic effects of marihuana smoking on luteinizing hormone, follicle-stimulating hormone and prolactin levels in human males. *Drug Alcohol Depend*. 1992;30(1).
11. Pacey AA, Povey AC, Clyma JA, et al. Modifiable and non-modifiable risk factors for poor sperm morphology. *Hum Reprod*. 2014;29(8):1629–1636.
12. Gundersen TD, Jørgensen N, Andersson AM, et al. Association Between Use of Marijuana and Male Reproductive Hormones and Semen Quality: A Study Among 1,215 Healthy Young Men. *Am J Epidemiol*. 2015;182(6):473–481.
13. Carroll K, Pottinger AM, Wynter S, DaCosta V. Marijuana use and its influence on sperm morphology and motility: identified risk for fertility among Jamaican men. *Andrology*.
14. Cushman P Jr. Plasma testosterone levels in healthy male marijuana smokers. *Am J Drug Alcohol Abuse*. 1975;2(2):269–275.
15. Thistle JE, Graubard BI, Braunlin M, et al. Marijuana use and serum testosterone concentrations among U.S. males. *Andrology*.
16. Lisano JK, Smith JD, Mathias AB, et al. Performance and Health-Related Characteristics of Physically Active Males Using Marijuana. *J Strength Cond Res*. 2019;33(6):1658–1668.
17. Nassan FL, Arvizu M, Mínguez-Alarcón L, et al. Marijuana smoking and markers of testicular function among men from a fertility centre. *Hum Reprod*. 2019;34(4).
18. Jordan T, Ngo B, Jones CA. The use of cannabis and perceptions of its effect on fertility among infertility patients. *Hum Reprod Open*. 2020;2020(1): hoz041. Published 2020 Feb 13.
19. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol*. 2009;62(10): e1–e34.
20. National Institute of Health and Department of Health and Human Services, “Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies,” 2015l.
21. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177–188.

22. Mantel N, Haenszel W. Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst.* 1959;22(4):719–748.
23. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327(7414):557–560.
24. Melsen WG, Bootsma MC, Rovers MM, Bonten MJ. The effects of clinical and statistical heterogeneity on the predictive values of results from meta-analyses. *Clin Microbiol Infect.* 2014;20(2):123-129. doi:10.1111/1469-0691.12494
25. Guzik DS, Overstreet JW, Factor-Litvak P, et al. Sperm morphology, motility, and concentration in fertile and infertile men. *N Engl J Med.* 2001;345(19):1388–1393.
26. Buck Louis GM, Sundaram R, Schisterman EF, et al. Semen quality and time to pregnancy: the Longitudinal Investigation of Fertility and the Environment Study. *Fertil Steril.* 2014;101(2):453–462.
27. Kasman AM, Thoma ME, McLain AC, Eisenberg ML. Association between use of marijuana and time to pregnancy in men and women: findings from the National Survey of Family Growth. *Fertil Steril.* 2018;109(5):866–871.
28. Wise LA, Wesselink AK, Hatch EE, et al. Marijuana use and fecundability in a North American preconception cohort study. *J Epidemiol Community Health.* 2018;72(3):208–215.
29. Nassan FL, Arvizu M, Mínguez-Alarcón L, et al. Marijuana smoking and outcomes of infertility treatment with assisted reproductive technologies. *Hum Reprod.* 2019;34(9):1818–1829.
30. Domar AD, Rooney KL, Milstein M, Conboy L. Lifestyle habits of 12,800 IVF patients: Prevalence of negative lifestyle behaviors, and impact of region and insurance coverage. *Hum Fertil (Camb).* 2015;18(4):253–257.
31. Callaghan RC, Sanches M, Benny C, Stockwell T, Sherk A, Kish SJ. Who consumes most of the cannabis in Canada? Profiles of cannabis consumption by quantity. *Drug Alcohol Depend.* 2019;205:107587. doi:10.1016/j.drugalcdep.2019.107587
32. Smith MJ, Alden EC, Herrold AA, et al. Recent Self-Reported Cannabis Use Is Associated With the Biometrics of Delta-9-Tetrahydrocannabinol. *J Stud Alcohol Drugs.* 2018;79(3):441–446. doi:10.15288/jsad.2018.79.441

## FIGURE LEGENDS

**Figure 1.** PRISMA flow diagram

**Figure 2.** Relative Risk of **A)** impaired semen quality; **B)** sperm morphology abnormalities in cannabis users. **CI:** confidence interval; **I<sup>2</sup>:** inconsistency.

**Figure 3.** Standardized mean difference of Testosterone levels between cannabis users and non-users. **S.m.d.:** standardized mean difference; **CI:** confidence interval; **I<sup>2</sup>:** inconsistency.

**Figure 4.** Standardized mean difference of **A)** FSH levels; **B)** LH levels between cannabis users and non-users. **S.m.d.:** standardized mean difference; **CI:** confidence interval; **I<sup>2</sup>:** inconsistency.






Figure 1. PRISMA flow diagram.

andr\_12953\_f1.tiff






Figure 2. Relative Risk of A) Impaired semen quality, B) sperm morphology abnormalities in cannabis users. CI: confidence interval; I<sup>2</sup>: inconsistency.

andr\_12953\_f2.tiff

Author Manuscript



**Figure 3.** Standardized mean difference of Testosterone levels between cannabis users and non-users. **s.m.d.:** standardized mean difference; **CI:** confidence interval;  $I^2$ : inconsistency.

andr\_12953\_f3.tiff





Figure 4. Standardized mean difference of A) FBH levels; B) LH levels between cannabis users and non-users. S.M.D.: standardized mean difference; CI: confidence interval;  $I^2$ : inconsistency.

andr\_12953\_f4.tiff