

# Clinical Utility of $\beta_3$ -Adrenoreceptor Agonists for the Treatment of Overactive Bladder: A Review of the Evidence and Current Recommendations

Jan Krhut <sup>1,2</sup>, Barbora Skugarevská<sup>1,2</sup>, David Míka<sup>1,2</sup>, Lars Lund <sup>3,4</sup>, Peter Zvara<sup>3-5</sup>

<sup>1</sup>Department of Urology, University Hospital, Ostrava, Czech Republic; <sup>2</sup>Department of Surgical Studies, Medical Faculty, Ostrava University, Ostrava, Czech Republic; <sup>3</sup>Research Unit of Urology, Department of Clinical Research, University of Southern Denmark, Odense, Denmark; <sup>4</sup>Department of Urology, Odense University Hospital, Odense, Denmark; <sup>5</sup>Biomedical Laboratory, University of Southern Denmark, Odense, Denmark

Correspondence: Jan Krhut, Department of Urology, University Hospital, Tř 17 Listopadu 1790, Ostrava 70852, Czech Republic, Tel +420 59-737-5307, Fax +420 59-737-5301, Email jan.krhut@fno.cz

**Abstract:** This nonsystematic review provides a summary of current evidence on the use of  $\beta_3$ -adrenoreceptor agonists ( $\beta_3$ -ARAs) for the treatment for lower urinary tract symptoms. Soon after their discovery in 1989,  $\beta_3$ -ARs were identified as a predominant adrenoreceptor subtype in the human urinary bladder. Although it is widely believed that  $\beta_3$ -ARAs cause detrusor relaxation, the effect on bladder afferent signaling likely plays an important role in their mechanism of action as well. In 2011 and 2012, mirabegron was approved for clinical use in overactive bladder (OAB) patients. Pooled analysis of data from prospective randomized studies on >60,000 OAB patients showed that when compared to placebo, mirabegron was superior with respect to reducing the frequency, number, and severity of urgency episodes, number of incontinence episodes and increasing dry rate, but not in reduction of nocturia episodes. The only side effect showing significantly higher incidence than placebo was nasopharyngitis. Mirabegron is approved for OAB treatment in all age-groups and in pediatric patients with neurogenic bladder. Vibegron is another  $\beta_3$ -ARA approved for OAB treatment in the US and Japan. Several large, multicenter, double-blind, randomized trials have documented statistically significant superiority of vibegron over placebo on all efficacy end points. Other  $\beta_3$ -ARAs are being developed; however, to date none has been introduced to clinical use. All  $\beta_3$ -ARAs provide efficacy similar to anticholinergics. They have a favorable safety profile and are well tolerated. Due to their different mechanisms of action, combination of  $\beta_3$ -ARAs with anticholinergic compounds allows for increased efficacy.

**Keywords:** urinary bladder, beta-mimetics, efficacy, safety, mirabegron, vibegron

## Introduction

Overactive bladder (OAB) syndrome is defined as urinary urgency, usually with urinary frequency and nocturia, with or without urgency urinary incontinence in the absence of urinary tract infection or other obvious pathology.<sup>1</sup> OAB affects 12%–16% of the adult population in Europe and the US.<sup>2</sup> Although OAB is not associated with high mortality, it has a detrimental impact on quality of life.<sup>3</sup> Lifestyle interventions and behavioral therapy are considered first-line OAB treatment. If those interventions fail, pharmacological treatment using anticholinergics (AChs) or  $\beta_3$ -adrenoreceptor agonists ( $\beta_3$ -ARAs) is recommended. The general aim of this nonsystematic review is to provide an overview of current evidence on the use of  $\beta_3$ -ARAs for OAB treatment.

## Role of $\beta_3$ Adrenoreceptors in Bladder Control under Physiological Conditions

The main role of the lower urinary tract (LUT) is accumulation of urine during the storage phase and its periodical elimination during the voiding phase of the micturition cycle. The function of the LUT is under constant control of the neural system, including peripheral nerves, the spinal cord, and supraspinal centers in the brain. Any disruption of this

control pathway may lead to development of LUT symptoms (LUTSs).<sup>4</sup> At the peripheral nerve level, parasympathetic efferents are responsible for detrusor contraction during voiding, while sympathetic nerves drive detrusor relaxation during the storage phase.

Adrenoreceptors (ARs), both  $\alpha$  and  $\beta$  classes, mediate the effect of the sympathetic nervous system by binding norepinephrine and epinephrine. The role of ARs in the regulation of heart function and vascular and pulmonary tonus has been known for >70 years.<sup>5</sup> There are three subtypes of  $\beta$ -ARs ( $\beta_1$ ,  $\beta_2$ , and  $\beta_3$ ), encoded by three genes. Stimulation of  $\beta_1$ -ARs has positive chronotropic, dromotropic, and inotropic effects on the heart, while stimulation of  $\beta_2$ -ARs results in smooth-muscle relaxation (bronchodilatation, uterolysis).  $\beta_3$ -ARs were first described in 1989,<sup>6</sup> and are expressed in several tissue types and organs, such as the brain, retina, myocardium, adipose tissue, and myometrium. In the urinary system, particularly the urinary bladder,  $\beta_3$ -ARs play a key role.<sup>7</sup>  $\beta_3$ -AR are the predominant AR subtype in the human urinary bladder.  $\beta_3$ -ARs account for >97% of all ARs expressed in the urinary bladder, while  $\beta_1$ -ARs and  $\beta_2$ -ARs account for only 1.5% and 1.4%, respectively.<sup>8</sup>

Activation of  $\beta_3$ -ARs leads to an increase in intracellular cyclic adenosine monophosphate (cAMP) levels and subsequent activation of cAMP-dependent PKA. PKA phosphorylates myosin light chain kinase (MLCK), which suppresses calcium-calmodulin-dependent interaction of myosin with actin, resulting in detrusor-muscle relaxation.<sup>9</sup> In addition, the increase in cAMP activity results in decreased intracellular calcium-ion concentration by removal of  $\text{Ca}^{2+}$  ions from the cytoplasm. Some suggest the involvement of other cAMP-independent mechanisms, such as the large-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  ( $\text{BK}_{\text{Ca}}$ ) channels in the process of  $\beta_3$ -AR activation-induced detrusor relaxation.<sup>10</sup> Some authors have proposed  $\beta_3$ -AR activation-induced decrease in acetylcholine release from both neuronal and nonneuronal sources (eg, from the urothelium and detrusor) during physiological bladder filling, thus suppressing parasympathetic activity.<sup>11</sup> However, despite extensive scientific inquiry over the last 30 years, the complex role of  $\beta_3$ -ARs in the control of bladder relaxation during the filling phase has not been fully elucidated.

## Mechanism of Action of $\beta_3$ -Adrenoreceptor Agonists in OAB Treatment

Early studies using animal OAB models indicated that the effect of  $\beta_3$ -ARAs was mainly through a direct relaxant effect on the detrusor, due to activation of cAMP and  $\text{BK}_{\text{Ca}}$  channels.<sup>12,13</sup> This theory has been questioned by several studies documenting that plasma concentrations after oral administration of mirabegron do not reach the concentration necessary to achieve detrusor relaxation during in vitro experiments using both animal and human isolated bladder strips. Since abundant expression of  $\beta_3$ -ARs has been documented at afferent nerve endings and interstitial cells in the suburothelial layer, as well as in the urothelium, it was proposed that a reduction in bladder afferent signaling could be the most important mechanism leading to detrusor relaxation.<sup>14</sup> This theory was supported by an evident decrease in afferent activity of both  $\text{A}_\delta$  and C fibers after mirabegron administration, which was associated with decreased spontaneous phasic activity of the urinary bladder wall in animal studies.<sup>15</sup>

Detrusor contraction during physiological voiding is mediated mainly by  $\text{M}_2$  and  $\text{M}_3$  muscarinic receptors. While  $\text{M}_3$  receptors probably play a key role in the initiation of detrusor contraction, the overrepresented  $\text{M}_2$  receptors are responsible for its maintenance. Nonadrenergic, noncholinergic mediators, such as ATP or bradykinin, are associated with increased nonvoiding detrusor-contraction activity under pathological conditions.  $\beta_3$ -ARAs inhibit detrusor contractions elicited by other stimuli more effectively than those elicited by cholinergic agonists.<sup>16</sup> This may explain the observation that  $\beta_3$ -ARAs attenuate pathologically increased phasic detrusor activity without interfering with physiological micturition contraction. In the long term, animal studies have shown that  $\beta_3$ -ARAs can protect bladder function and morphology during chronic ischemia. Should these data be confirmed in humans,  $\beta_3$ -ARAs may represent a new modality to prevent chronic ischemia-related bladder pathology.<sup>17</sup>

## $\beta_3$ -Adrenoreceptor Agonists Currently Available for OAB Treatment

Mirabegron was originally intended as a treatment for diabetes, but was later repurposed for the treatment of OAB.<sup>18</sup> It obtained Japanese marketing approval for the treatment of OAB in July 2011. The US Food and Drug Administration (FDA) and European Medicines Agency approved mirabegron for clinical use in the US in June 2012 and Europe in December 2012. In May 2018, the FDA approved mirabegron in combination with solifenacin for the treatment of OAB.

In March 2021, the FDA expanded the indication to treatment of neurogenic detrusor overactivity (NDO) in children aged 3 years and older. Vibegron was developed in a collaboration of Kyorin Pharmaceutical, Kissei Pharmaceutical, and Urovant Sciences. Based on results of the global phase III EMPOWUR trial, vibegron was granted approval in Japan in September 2018 and the US in December 2020.<sup>19</sup>

Another drug under clinical development is solabegron. It showed OAB-symptom reduction in a randomized, placebo-controlled phase II trial. Solabegron 125 mg produced a statistically significant difference in incontinence episodes and micturition frequency over 24 hours and an increase in voided volume.<sup>20</sup> More recently (June 2021), Velicept Therapeutics, which acquired all assets related to solabegron from GlaxoSmithKline in March 2011, announced that solabegron had met the primary end point in a phase IIb study in patients with OAB. This 12-week placebo-controlled study enrolled 435 women and demonstrated a statistically significant improvement in number of micturitions, number of urgency episodes, and urge urinary incontinence episodes over 24 hours in a solabegron-treated group compared to placebo. These results have not been published yet. In contrast, ritobegron (Kissei Pharmaceutical) failed to meet the primary end point in a phase II trial (NCT00742833) in 2008, and further clinical development ceased. Several phase I studies using other  $\beta_3$ -ARAs (AJ9677, 138-335, TRK380, and ZD7114) were performed during the early 2000s; however, none of these compounds advanced to the next stages of clinical development.<sup>21</sup>

## Mirabegron

### Pharmacological Properties

Mirabegron is rapidly absorbed after oral administration, reaching maximum plasma concentration in 3–5 hours whether taken with or without food. When administered as a single daily dose of 50 mg, steady-state concentrations are usually achieved within 7 days. The compound is highly lipophilic, metabolized in the liver, and eliminated in the urine (55%) and feces (34%), mainly in unchanged form. Terminal elimination half-life is about 23–25 hours. Mirabegron shows minimal interactions with other frequently prescribed drugs, such as digoxin, warfarin, metformin, and oral contraceptives. On the other hand, caution is advised when prescribing mirabegron to patients on certain antiarrhythmics (thioridazine) or tricyclic antidepressants (imipramine, desipramine).<sup>22</sup> Due to a lack of data, mirabegron should not be prescribed to patients with severe impairment of renal and/or liver function.<sup>22,23</sup>

### Clinical Efficacy

A strong body of evidence from numerous prospective randomised studies on >60,000 participants (66%–75% female) proved the efficacy of mirabegron. When compared to placebo, mirabegron was superior with respect to reducing frequency, number, and severity of urgency episodes, number of incontinence episodes, and increasing dry rate. In contrast, reduction in nocturia episodes for mirabegron has been seen in few studies. After 12 weeks of treatment with mirabegron, one may expect a reduction in number of voids of 16%–25%, number of severe urgency episodes of 33%–40%, and number of incontinence episodes of 53–63%. Increase in volume per void of 13%–15% has also been documented.<sup>24</sup> A majority of studies have documented both reduction of symptom severity and improvement in treatment satisfaction and quality of life.<sup>25</sup>

The fact that a significant proportion of studies enrolled patients who had discontinued previous ACh treatment may imply that mirabegron is more effective than AChs. However, this is not the case. Based on randomized trials comparing mirabegron with AChs, mirabegron 50 mg once daily showed comparable overall efficacy; however, individual AChs were more efficacious than mirabegron with respect to certain variables (eg, solifenacin 10 mg was superior with respect to micturition frequency, fesoterodine 8 mg with respect to number of incontinence episodes, and trospium 60 mg, solifenacin 10 mg, and fesoterodine 8 mg with respect to dry rate).<sup>26</sup> In the FAVOR study, which included patients after withdrawal of AChs, there were no significant differences observed in mirabegron efficacy between those who discontinued AChs due to low efficacy and those who discontinued due to adverse effects.<sup>27</sup> Generally, the onset of effect in mirabegron recipients is usually observed within 1 month, reaching a maximum within 12–16 weeks after treatment initiation. The effect remains stable over 1 year if the drug is taken regularly.<sup>28</sup>

## Clinical Efficacy in Specific OAB-Patient Cohorts

### Mirabegron in Men

Our current knowledge on the clinical efficacy of mirabegron in men is based mainly on pooled analysis of previously conducted phase II and III trials enrolling both men and women.<sup>29</sup> Studies designed primarily to assess the efficacy of mirabegron in the treatment of OAB symptoms in men are rather sparse.<sup>30</sup> A majority of placebo-controlled studies have documented statistically significant superiority of mirabegron on some end points. However the difference in most cases was small and the clinical value of such improvement questionable. In addition, data showing that female patients have better adherence to mirabegron treatment than men may implicate lower efficacy in men than women.<sup>31</sup> This is consistent with previous findings demonstrating that male patients suffering from OAB symptoms generally have a lower response rate when treated using AChs than women. This probably reflects the fact that the etiology of male LUTSs is more complex, as it may include additional factors, such as bladder-outflow obstruction, resulting in urothelial dysfunction, chronic bladder ischemia, and inflammation.<sup>32</sup>

### Mirabegron in Adults with Neurogenic Detrusor Overactivity

Evidence on the efficacy of mirabegron in treatment of NDO is inconsistent and based on several small single-center studies on patients with NDO due to various underlying pathologies. Vasudeva et al reported results in a group of 30 patients with spinal cord injury treated with a single daily dose of mirabegron 50 mg for 6 weeks. They observed significant improvement in bladder diary-derived variables, such as mean catheterization frequency, mean catheterization volume, and mean number of incontinence episodes, as well as in urodynamic parameters (mean cystometric capacity and maximum amplitude of detrusor pressure).<sup>33</sup>

Two studies published by Krhut et al and Welk et al piloted the use of mirabegron in patients with NDO due to multiple sclerosis.<sup>34,35</sup> Subsequently, another two studies demonstrated similar efficacy for mirabegron and solifenacin in multiple sclerosis patients.<sup>36,37</sup> In contrast to an initially prospective study by Gubbiotti et al that reported improvement in OAB symptoms in patients with Parkinson's disease, the prospective randomised PaDoMi study did not find any statistical difference between mirabegron and placebo groups in bladder diary-derived variables or quality of life after 12 weeks of treatment.<sup>38,39</sup> Solid evidence on mirabegron efficacy in other diseases resulting in NDO is lacking. In summary, the evidence on mirabegron in NDO patients is limited, and further well-designed prospective trials in well-specified homogeneous populations are warranted.

### Mirabegron in the Elderly

Given the increasing prevalence of OAB symptoms with age, special attention should be paid to elderly OAB patients. It has been demonstrated that they suffer from more comorbidities, especially cardiovascular, neurological, and psychiatric, than age-matched individuals without OAB.<sup>40</sup> This is often associated with polypharmacy, with many of the most commonly prescribed drugs having a strong ACh effect that can be further increased by drug interactions.<sup>41</sup> Additional use of AChs for OAB may lead to a significant increase in ACh load with an increased risk of cognitive impairment and dementia.<sup>42</sup> Other common ACh side effects, such as dry mouth, constipation, and QT prolongation, to which older people are particularly susceptible, should also be taken into consideration. Therefore, the use of mirabegron or other  $\beta_3$ -ARAs may represent a valuable alternative to AChs in this vulnerable population.

The efficacy of mirabegron in patients aged >65 years was assessed in a large double-blind, randomized, placebo-controlled, multicenter study that enrolled 888 subjects with OAB wet in the US and Canada. After 12 weeks of treatment, the authors found significant superiority of mirabegron over placebo with respect to reduction in frequency, number of urgency and incontinence episodes, and mean voided volume.<sup>43</sup> These findings are consistent with those published previously, which reported a pooled analysis of patients aged >65 years enrolled in three phase III studies.<sup>44</sup> These data were also supported by work of Lozano-Ortega et al who proved that mirabegron was similarly as effective as ACh across all efficacy end points in an older population.<sup>45</sup>

### Mirabegron in Children

OAB affects 5%–12% of 5- to 10-year-old children and 0.5% of adolescents (16–18 years of age). Around a third of individuals who had had OAB during childhood are likely to suffer from OAB in adulthood.<sup>46</sup> Pharmacokinetic data on

mirabegron in a pediatric population were published recently; however, clinical data are limited.<sup>47</sup> Blais et al performed an open-label study that enrolled 58 patients (14 girls, 44 boys) with a mean age of 10.1 years. Clinical response, defined as 50% reduction in OAB symptoms, was achieved in 89% of patients.<sup>48</sup> Based on data from a retrospective study on 45 children with idiopathic OAB, Kim et al suggested that mirabegron had comparable efficacy to solifenacin.<sup>49</sup> Recently, a beneficial effect of treatment with mirabegron on both subjective variables and urodynamic parameters was documented in a cohort of 68 children suffering from NDO.<sup>50</sup> Given the low prevalence of neurogenic OAB in children, the existing data were found to be sufficient to grant approval for mirabegron as treatment of NDO in children aged  $\geq 3$  years.

## Clinical Efficacy of Mirabegron in Combination Treatment

Both AChs and  $\beta_3$ -ARAs are standard treatments for OAB. Given their different modes of action, simultaneous administration of both drug classes is a logical way to improve efficacy if monotherapy is unable to control OAB symptoms sufficiently. In addition, especially in older individuals, the combination of AChs and  $\beta_3$ -ARAs allows for limiting unwanted ACh load. Combinations of AChs and  $\beta_3$ -ARAs have been assessed in several large-scale, multicentric prospective, randomised phase II and III studies that recruited mostly women (66%–77%) suffering from OAB wet (BESIDE, SYNERGY, SYNERGY II, and SYMPHONY). These trials documented the superiority of the combination over monotherapy on a majority of end points, showing a 31%–50% higher likelihood of achieving continence using the combination than monotherapy.<sup>51</sup> A pooled analysis of the BESIDE study of patients aged  $>65$  years documented that the combination of mirabegron and solifenacin was superior to solifenacin monotherapy without increasing the rate of adverse events.<sup>52</sup>

There is a well-known association between OAB and benign prostatic hyperplasia–induced bladder-outlet obstruction (BOO) in men. Up to 50% of men with LUTSs and urodynamically confirmed BOO have detrusor overactivity.<sup>53</sup>  $\alpha_1$  blockers ( $\alpha_1$ Bs) are considered first-line pharmacological treatment for LUTSs in men with clinical evidence of BOO. They have proved to improve voiding symptoms and well-being in men; however, they often fail to improve OAB symptoms. Therefore, AChs are recommended to be administered simultaneously or as an add-on for those with persistent OAB symptoms. In order to avoid ACh burden, a combination of  $\alpha_1$ Bs with mirabegron has been proposed in place of AChs. This concept has been assessed in two large, prospective, double-blind, placebo-controlled studies (PLUS,  $n=676$ ; MATCH,  $n=568$ ) and several smaller trials.<sup>54,55</sup> Despite some controversies over outcomes in individual variables, all studies generally support the benefit of combination therapy over monotherapy.<sup>56</sup>

## Safety and Tolerability

A recently published systematic review of mirabegron safety compared to placebo analyzed 6,135 patients recruited in ten trials. Most frequently reported side effects were nasopharyngitis, dry mouth, hypertension, constipation, headache, dyspepsia, urinary tract infection, dizziness, blurred vision, nausea, cardiovascular events, influenza, prolonged QT interval, and upper respiratory tract infection. With the exception of nasopharyngitis, no significant difference was found in the incidence of side effects between mirabegron 50 mg and placebo (OR 1.54,  $p=0.03$ ).<sup>57</sup>

Several other studies have evaluated the safety of mirabegron in specific patient subgroups. Nitti et al reported the effect of mirabegron on urodynamic parameters in aged  $> 45$  years with LUTSs and BOO. They found that mirabegron 50 mg and 100 mg did not negatively affect voiding efficacy compared to placebo after 12 weeks of treatment ( $-0.4$  mL/s in maximal flow rate and  $-1.4$  cm H<sub>2</sub>O in detrusor pressure at maximum urinary flow).<sup>58</sup> These findings correlate with clinical data indicating low incidence of acute urinary retention episodes and minimal increase in postvoid residual in men treated with mirabegron.<sup>59</sup> Very good safety profiles of mirabegron in children and neurogenic patients have been documented in several studies.<sup>60,61</sup> Recently, Griebeling et al found no increased risk of cognitive impairment in elderly patients receiving mirabegron for 12 weeks (Pillar study).<sup>62</sup> Another study based on data obtained from national health registers (Denmark, Spain, Sweden, UK, US) did not reveal any increased risk of cancer among mirabegron users compared with ACh medications in men or women.<sup>63</sup>

## Adherence and Persistence

OAB is a chronic condition that requires long-term treatment, with only a small proportion of patients able to discontinue the treatment due to symptom cessation.<sup>64</sup> Mirabegron seems to have a well-balanced efficacy:side effects ratio; therefore, adherence to mirabegron and treatment persistence should be significantly better than AChs. However, a recent systematic literature review of real-world data documented that 1-year persistence rates were 12%–25% and 32%–38% for antimuscarinic agents and mirabegron, respectively. Median time to discontinuation was 5–6.5 months for AChs and 5.6–7.4 months for mirabegron.<sup>65</sup>

## Vibegron

### Pharmacological Properties

Vibegron reaches peak plasma concentration 1–3 hours after oral administration, and constant blood concentration is achieved in 7 days of once-daily administration. It is excreted in feces and urine as unchanged drug. The plasma concentration of the drug increases in people >65 years of age and those with moderate–severe renal impairment. Vibegron does not induce or inhibit the activity of CYP2D6 and CYP3A4 enzymes, and thus drug–drug interactions with most frequently prescribed agents are limited. The elimination half-life is approximately 70 hours.<sup>19</sup>

### Clinical Efficacy

A multicenter phase IIb dose-finding study randomizing 1,395 subjects showed significant superiority of vibegron 50 mg and 100 mg over placebo with respect to decreased frequency, reduced urgency-incontinence episodes, incontinence episodes, and urgency episodes per day. Data from the second part of this study documented the additive effect of the concomitant administration of vibegron plus tolterodine 4 mg once daily.<sup>66</sup> A large multicenter double-blind randomized phase III study (EMPOWUR, n=1,518) comparing vibegron 75 mg to placebo and tolterodine 4 mg extended release as active control documented statistically significant superiority of vibegron over placebo on all efficacy end points.<sup>67</sup> In the open-label extension of this study (n=505), significant superiority of vibegron 75 mg over tolterodine was observed in respect of mean change from baseline in urgency-incontinence episodes after 52 weeks of treatment. No other outcomes for efficacy end points differed.<sup>68</sup> Meaningful clinical data on the efficacy of virabegron in specific OAB-patients cohort are not yet available.

### Safety and Tolerability

Based on the results of the available phase IIb and III trials, virabegron seems generally safe and well tolerated.<sup>69</sup> The incidence and severity of drug-related side effects are comparable to placebo. A recently published pooled analysis of the EMPOWUR study confirmed the very good safety profile of virabegron in patients aged  $\geq 65$  years.<sup>70</sup>

## Recommendations on the Use of $\beta_3$ -Adrenoreceptor Agonists in OAB Treatment

Orally administered AChs are considered the standard pharmacological treatment.  $\beta_3$ -ARAs should be offered as an alternative to AChs in both men and women. If monotherapy is not effective, combination treatment with AChs and  $\beta_3$ -ARAs may lead to greater benefit.<sup>71–73</sup>

## Conclusion

$\beta_3$ -ARAs represent a valuable addition to pharmacological treatment of OAB syndrome. They provide better efficacy than placebo and similar efficacy to ACh.  $\beta_3$ -ARAs have a favorable safety profile and are well tolerated. Due to their different mechanisms of action, combinations of  $\beta_3$ -ARAs with AChs,  $\alpha_1$ Bs, or other compounds allow for higher efficacy.

## Disclosure

JK is a researcher and consultant for Astellas, Coloplast, STIMVIA, and Medtronic outside the submitted work. The authors report no other conflicts of interest in this work.

## References

1. Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. *Neurourol Urodyn*. 2002;21:167–178. doi:10.1002/nau.10052
2. Irwin DE, Milsom I, Hunskaar S, et al. Population-based survey of urinary incontinence, overactive bladder, and other lower urinary tract symptoms in five countries: results of the EPIC study. *Eur Urol*. 2006;50:1306–1314. doi:10.1016/j.eururo.2006.09.019
3. Milsom I, Abrams P, Cardozo L, et al. How widespread are the symptoms of an overactive bladder and how are they managed? A population-based prevalence study. *BJU Int*. 2001;87:760–766. doi:10.1046/j.1464-410x.2001.02228.x
4. de Groat WC, Griffiths D, Yoshimura N. Neural control of the lower urinary tract. *Compr Physiol*. 2015;5:327–396. doi:10.1002/cphy.c130056
5. Ahlquist RP. A study of the adrenergic receptors. *Am J Physiol*. 1948;153:586–600. doi:10.1152/ajplegacy.1948.153.3.586
6. Emorine LJ, Marullo S, Briand-Sutren MM, et al. Molecular characterization of the human beta 3-adrenergic receptor. *Science*. 1989;245:1118–1121. doi:10.1126/science.2570461
7. Michel MC, Vrydag W. Alpha1-, alpha2- and beta-adrenoceptors in the urinary bladder, urethra and prostate. *Br J Pharmacol*. 2006;147(Suppl 2):S88–119. doi:10.1038/sj.bjp.0706619
8. Nomiya M, Yamaguchi O. A quantitative analysis of mRNA expression of alpha 1 and beta-adrenoceptor subtypes and their functional roles in human normal and obstructed bladders. *J Urol*. 2003;170:649–653. doi:10.1097/01.ju.0000067621.62736.7c
9. Yamaguchi O, Chapple CR. Beta3-adrenoceptors in urinary bladder. *Neurourol Urodyn*. 2007;26:752–756. doi:10.1002/nau.20420
10. Cernecka H, Kersten K, Maarsingh H, et al.  $\beta_3$ -adrenoceptor-mediated relaxation of rat and human urinary bladder: roles of BKCa channels and Rho kinase. *Naunyn Schmiedebergs Arch Pharmacol*. 2015;388:749–759. doi:10.1007/s00210-015-1128-z
11. Andersson KE. Antimuscarinic mechanisms and the overactive detrusor: an update. *Eur Urol*. 2011;59:377–386. doi:10.1016/j.eururo.2010.11.040
12. Andersson KE, Arner A. Urinary bladder contraction and relaxation: physiology and pathophysiology. *Physiol Rev*. 2004;84:935–986. doi:10.1152/physrev.00038.2003
13. Petkov GV, Nelson MT. Differential regulation of Ca<sup>2+</sup>-activated K<sup>+</sup> channels by beta-adrenoceptors in Guinea pig urinary bladder smooth muscle. *Am J Physiol Cell Physiol*. 2005;288:C1255–1263. doi:10.1152/ajpcell.00381.2004
14. Aizawa N, Homma Y, Igawa Y. Effects of mirabegron, a novel  $\beta_3$ -adrenoceptor agonist, on primary bladder afferent activity and bladder microcontractions in rats compared with the effects of oxybutynin. *Eur Urol*. 2012;62(6):1165–1173. doi:10.1016/j.eururo.2012.08.056
15. Igawa Y, Aizawa N, Michel MC.  $\beta_3$ -Adrenoceptors in the normal and diseased urinary bladder—What are the open questions? *Br J Pharmacol*. 2019;176:2525–2538. doi:10.1111/bph.14658
16. Michel MC, Sand C. Effect of pre-contraction on  $\beta$ -adrenoceptor-mediated relaxation of rat urinary bladder. *World J Urol*. 2009;27:711–715. doi:10.1007/s00345-009-0416-y
17. Sawada N, Nomiya M, Hood B, et al. Protective effect of a  $\beta_3$ -adrenoceptor agonist on bladder function in a rat model of chronic bladder ischemia. *Eur Urol*. 2013;64:664–671. doi:10.1016/j.eururo.2013.06.043
18. Tyagi P, Tyagi V. Mirabegron, a  $\beta_3$ -adrenoceptor agonist for the potential treatment of urinary frequency, urinary incontinence or urgency associated with overactive bladder. *IDrugs*. 2010;13:713–722.
19. Keam SJ. Vibegron: first Global Approval. *Drugs*. 2018;78:1835–1839. doi:10.1007/s40265-018-1006-3
20. Ohlstein EH, von Keitz A, Michel MC. A multicenter, double-blind, randomized, placebo-controlled trial of the  $\beta_3$ -adrenoceptor agonist solabegron for overactive bladder. *Eur Urol*. 2012;62:834–840. doi:10.1016/j.eururo.2012.05.053
21. Michel MC.  $\beta$ -adrenergic receptor subtypes in the urinary tract. In: Andersson KE, Michel MC, editors. *Handbook of Experimental Pharmacology Vol. 202: Urinary Tract*. Berlin Heidelberg: Springer; 2011:307–318.
22. European Medicines Agency. Betmiga prolonged-release tablets: summary of product characteristics; 2017. Available from: <http://www.ema.europa.eu/>. Accessed August 19, 2021.
23. Krauwinkel W, van Dijk J, Schaddelee M, et al. Pharmacokinetic properties of mirabegron, a  $\beta_3$ -adrenoceptor agonist: results from two phase I, randomized, multiple-dose studies in healthy young and elderly men and women. *Clin Ther*. 2012;34:2144–2160. doi:10.1016/j.clinthera.2012.09.010
24. Deeks ED. Mirabegron: a review in overactive bladder syndrome. *Drugs*. 2018;78:833–844. doi:10.1007/s40265-018-0924-4
25. Freeman R, Foley S, Rosa Arias J, et al. Mirabegron improves quality-of-life, treatment satisfaction, and persistence in patients with overactive bladder: a multi-center, non-interventional, real-world, 12-month study. *Curr Med Res Opin*. 2018;34:785–793. doi:10.1080/03007995.2017.1419170
26. Kelleher C, Hakimi Z, Zur R, et al. Efficacy and tolerability of mirabegron compared with antimuscarinic monotherapy or combination therapies for overactive bladder: a systematic review and network meta-analysis. *Eur Urol*. 2018;74:324–333. doi:10.1016/j.eururo.2018.03.020
27. Ko KJ, Choo MS, Chang YS, et al. A multicenter prospective study for overactive bladder patient treatment satisfaction with mirabegron after being unsatisfied with antimuscarinic therapy (FAVOR study). *Neurourol Urodyn*. 2020;39:2417–2424. doi:10.1002/nau.24505
28. Yamaguchi O, Ikeda Y, Ohkawa S. Phase III study to assess long-term (52-week) safety and efficacy of mirabegron, a  $\beta_3$ -adrenoceptor agonist, in Japanese patients with overactive bladder. *Low Urin Tract Symptoms*. 2017;9:38–45. doi:10.1111/luts.12107
29. Tubaro A, Batista JE, Nitti VW, et al. Efficacy and safety of daily mirabegron 50 mg in male patients with overactive bladder: a critical analysis of five phase III studies. *Ther Adv Urol*. 2017;9:137–154. doi:10.1177/1756287217702797
30. Shin DG, Kim HW, Yoon SJ, et al. Mirabegron as a treatment for overactive bladder symptoms in men (MIRACLE study): efficacy and safety results from a multicenter, randomized, double-blind, placebo-controlled, parallel comparison phase IV study. *Neurourol Urodyn*. 2019;38:295–304. doi:10.1002/nau.23852
31. Carlson KV, Rovner ES, Nair KV, et al. Persistence with mirabegron or antimuscarinic treatment for overactive bladder syndrome: findings from the PERSPECTIVE registry study. *Low Urin Tract Symptoms*. 2021;13:425–434. doi:10.1111/luts.12382

32. De Nunzio C, Brucker B, Bschiepfer T, et al. Beyond antimuscarinics: a review of pharmacological and interventional options for overactive bladder management in men. *Eur Urol.* 2021;79:492–504. doi:10.1016/j.eururo.2020.12.032
33. Vasudeva P, Prasad V, Yadav S, et al. Efficacy and safety of mirabegron for the treatment of neurogenic detrusor overactivity resulting from traumatic spinal cord injury: a prospective study. *Neurol Urodyn.* 2021;40:666–671. doi:10.1002/nau.24604
34. Krhut J, Borovička V, Bilková K, et al. Efficacy and safety of mirabegron for the treatment of neurogenic detrusor overactivity - Prospective, randomized, double-blind, placebo-controlled study. *Neurol Urodyn.* 2018;37:2226–2233. doi:10.1002/nau.23566
35. Welk B, Hickling D, McKibbin M, et al. A pilot randomized-controlled trial of the urodynamic efficacy of mirabegron for patients with neurogenic lower urinary tract dysfunction. *Neurol Urodyn.* 2018;37:2810–2817. doi:10.1002/nau.23774
36. Glykas I, Fragkoulis C, Mitsikostas DD, et al. B3 agonists or anticholinergics in the treatment of the lower urinary tract dysfunction in patients with multiple sclerosis? - A randomized study. *World J Urol.* 2021;39:3049–3056. doi:10.1007/s00345-020-03555-8
37. Brucker BM, Jericevic D, Rude T, et al. Mirabegron versus solifenacin in multiple sclerosis patients with overactive bladder symptoms: a prospective comparative nonrandomized study. *Urology.* 2020;145:94–99. doi:10.1016/j.urology.2020.08.008
38. Gubbiotti M, Conte A, Di Stasi SM, et al. Feasibility of mirabegron in the treatment of overactive bladder in patients affected by Parkinson's disease: a pilot study. *Ther Adv Neurol Disord.* 2019;12:1756286419843458. doi:10.1177/1756286419843458
39. Cho SY, Jeong SJ, Lee S, et al. Mirabegron for treatment of overactive bladder symptoms in patients with Parkinson's disease: a double-blind, randomized placebo-controlled trial (Parkinson's Disease Overactive bladder Mirabegron, PaDoMi Study). *Neurol Urodyn.* 2021;40:286–294. doi:10.1002/nau.24552
40. Asche CV, Kim J, Kulkarni AS, et al. Presence of central nervous system, cardiovascular and overall co-morbidity burden in patients with overactive bladder disorder in a real-world setting. *BJU Int.* 2012;109(4):572–580. doi:10.1111/j.1464-410X.2011.10436.x
41. Hines LE, Murphy JE. Potentially harmful drug-drug interactions in the elderly: a review. *Am J Geriatr Pharmacother.* 2011;9(6):364–377. doi:10.1016/j.amjopharm.2011.10.004
42. Rutman MP, Horn JR, Newman DK, et al. Overactive bladder prescribing considerations: the role of polypharmacy, anticholinergic burden, and CYP2D6 drug-drug interactions. *Clin Drug Investig.* 2021;41(4):293–302. doi:10.1007/s40261-021-01020-x
43. Herschorn S, Staskin D, Schermer CR, et al. Safety and tolerability results from the PILLAR study: a phase IV, double-blind, randomized, placebo-controlled study of mirabegron in patients  $\geq 65$  years with overactive bladder-wet. *Drugs Aging.* 2020;37:665–676. doi:10.1007/s40266-020-00783-w
44. Wagg A, Cardozo L, Nitti VW, et al. The efficacy and tolerability of the  $\beta$ -adrenoceptor agonist mirabegron for the treatment of symptoms of overactive bladder in older patients. *Age Ageing.* 2014;43:666–675. doi:10.1093/ageing/afu017
45. Lozano-Ortega G, Walker DR, Johnston K, et al. Comparative safety and efficacy of treatments for overactive bladder among older adults: a network meta-analysis. *Drugs Aging.* 2020;37:801–816. doi:10.1007/s40266-020-00792-9
46. Franco I. Overactive bladder in children. *Nat Rev Urol.* 2016;13:520–532. doi:10.1038/nrurol.2016.152
47. Rittig S, Baka-Ostrowska M, Tøndel C, et al. The pharmacokinetics, safety, and tolerability of mirabegron in children and adolescents with neurogenic detrusor overactivity or idiopathic overactive bladder and development of a population pharmacokinetic model-based pediatric dose estimation. *J Pediatr Urol.* 2020;16:31.e1–31.e10. doi:10.1016/j.jpuro.2019.10.009
48. Blais AS, Nadeau G, Moore K, et al. Prospective pilot study of mirabegron in pediatric patients with overactive bladder. *Eur Urol.* 2016;70:9–13. doi:10.1016/j.eururo.2016.02.007
49. Kim SC, Park M, Chae C, et al. Efficacy and tolerability of mirabegron compared with solifenacin for children with idiopathic overactive bladder: a preliminary study. *Investig Clin Urol.* 2021;62:317–323. doi:10.4111/icu.20200380
50. Baka-Ostrowska M, Bolong DT, Persu C, et al. Efficacy and safety of mirabegron in children and adolescents with neurogenic detrusor overactivity: an open-label, Phase 3, dose-titration study. *Neurol Urodyn.* 2021;40:1490–1499. doi:10.1002/nau.24657
51. Rutledge EC, Hernandez N, Gonzalez RR. Contemporary landmark trials update in the management of idiopathic overactive bladder. *Curr Bladder Dysfunct Rep.* 2020;15:377–385. doi:10.1007/s11884-020-00617-w
52. Gibson W, MacDiarmid S, Huang M, et al. Treating overactive bladder in older patients with a combination of mirabegron and solifenacin: a prespecified analysis from the BESIDE study. *Eur Urol Focus.* 2017;3(6):629–638. doi:10.1016/j.euf.2017.08.008
53. Eapen R, Radomski S. Review of the epidemiology of overactive bladder. *Res Rep Urol.* 2016;8:71–76. doi:10.2147/RRU.S102441
54. Kaplan SA, Herschorn S, McVary KT, et al. Efficacy and safety of mirabegron versus placebo add-on therapy in men with overactive bladder symptoms receiving tamsulosin for underlying benign prostatic hyperplasia: a randomized, phase 4 study (PLUS). *J Urol.* 2020;203:1163–1171. doi:10.1097/JU.0000000000000738
55. Kakizaki H, Lee KS, Yamamoto O, et al. Mirabegron add-on therapy to tamsulosin for the treatment of overactive bladder in men with lower urinary tract symptoms: a randomized, placebo-controlled study (MATCH). *Eur Urol Focus.* 2020;6:729–737. doi:10.1016/j.euf.2019.10.019
56. Ichihara K, Masumori N, Fukuta F, et al. A randomized controlled study of the efficacy of tamsulosin monotherapy and its combination with mirabegron for overactive bladder induced by benign prostatic obstruction. *J Urol.* 2015;193:921–926. doi:10.1016/j.juro.2014.09.091
57. Hou J, Xu F, Du H, et al. Adverse events associated with mirabegron 50mg versus placebo: a systematic review and meta-analysis. *Prog Urol.* 2021;31:SS1166–2. doi:10.1016/j.puro.2021.05.005
58. Nitti VW, Rosenberg S, Mitcheson DH, et al. Urodynamics and safety of the  $\beta$ 3 -adrenoceptor agonist mirabegron in males with lower urinary tract symptoms and bladder outlet obstruction. *J Urol.* 2013;190:1320–1327. doi:10.1016/j.juro.2013.05.062
59. Moss MC, Rezan T, Karaman UR, et al. Treatment of concomitant OAB and BPH. *Curr Urol Rep.* 2017;18:1. doi:10.1007/s11934-017-0649-z
60. Sager C, Sanmartino M, Burek C, et al. Efficacy and safety of Mirabegron as adjuvant treatment in children with refractory neurogenic bladder dysfunction. *J Pediatr Urol.* 2020;16:655.e1–655.e6. doi:10.1016/j.jpuro.2020.07.020
61. Krhut J, Wohlfahrt P, Pudich J, et al. Cardiovascular safety of mirabegron in individuals treated for spinal cord injury- or multiple sclerosis-induced neurogenic detrusor overactivity. *Int Urol Nephrol.* 2021;53:1089–1095. doi:10.1007/s11255-020-02774-7
62. Griebing TL, Campbell NL, Mangel J, et al. Effect of mirabegron on cognitive function in elderly patients with overactive bladder: MoCA results from a phase 4 randomized, placebo-controlled study (PILLAR). *BMC Geriatr.* 2020;20:109. doi:10.1186/s12877-020-1474-7
63. Phiri K, Hallas J, Linder M, et al. A study of cancer occurrence in users of mirabegron and antimuscarinic treatments for overactive bladder. *Curr Med Res Opin.* 2021;37:867–877. doi:10.1080/03007995.2021.1891035
64. Krhut J, Martan A, Zachoval R, et al. Is it possible to cure the symptoms of the overactive bladder in women? *Int Urol Nephrol.* 2018;50:433–439. doi:10.1007/s11255-017-1777-6

65. Ali M, Grogan S, Powell S, et al. Qualitative analysis of factors influencing patient persistence and adherence to prescribed overactive bladder medication in UK primary care. *Adv Ther.* 2019;36:3110–3122. doi:10.1007/s12325-019-01098-y
66. Mitcheson HD, Samanta S, Muldowney K, et al. Vibegron (RVT-901/MK-4618/KRP-114V) administered once daily as monotherapy or concomitantly with tolterodine in patients with an overactive bladder: a multicenter, phase iib, randomized, double-blind, controlled trial. *Eur Urol.* 2019;75:274–282. doi:10.1016/j.eururo.2018.10.006
67. Staskin D, Frankel J, Varano S, et al. International phase III, randomized, double-blind, placebo and active controlled study to evaluate the safety and efficacy of vibegron in patients with symptoms of overactive bladder: EMPOWUR. *J Urol.* 2020;204:316–324. doi:10.1097/JU.0000000000000807
68. Staskin D, Frankel J, Varano S, et al. Once-daily vibegron 75 mg for overactive bladder: long-term safety and efficacy from a double-blind extension study of the international phase 3 trial (EMPOWUR). *J Urol.* 2021;205:1421–1429. doi:10.1097/JU.0000000000001574
69. Su S, Liang L, Lin J, et al. Systematic review and meta-analysis of the efficacy and safety of vibegron vs antimuscarinic monotherapy for overactive bladder. *Medicine.* 2021;100(5):e23171. doi:10.1097/MD.00000000000023171
70. Varano S, Staskin D, Frankel J, et al. Efficacy and safety of once-daily vibegron for treatment of overactive bladder in patients aged  $\geq 65$  and  $\geq 75$  years: subpopulation analysis from the EMPOWUR randomized, international, phase III study. *Drugs Aging.* 2021;38:137–146. doi:10.1007/s40266-020-00829-z
71. Lightner DJ, Gomelsky A, Souter L, et al. Diagnosis and treatment of overactive bladder (non-neurogenic) in adults: AUA/SUFU guideline amendment 2019. *J Urol.* 2019;202:558–563. doi:10.1097/JU.0000000000000309
72. Harding CK, Lapitan MC, Arlandis S, et al. *EAU Guidelines on Management of Non-Neurogenic Female Lower Urinary Tract Symptoms (LUTS)*. Arnhem, The Netherlands: EAU Guidelines Office; 2021.
73. Gravas S, Cornu JN, Gacci M, et al. *EAU Guidelines on Management of Non-Neurogenic Male Lower Urinary Tract Symptoms (LUTS), Incl. Benign Prostatic Obstruction (BPO)*. Arnhem, The Netherlands: EAU Guidelines Office; 2021.

## Research and Reports in Urology

Dovepress

### Publish your work in this journal

Research and Reports in Urology is an international, peer-reviewed, open access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of adult and pediatric urology in the clinic and laboratory including the following topics: Pathology, pathophysiology of urological disease; Investigation and treatment of urological disease; Pharmacology of drugs used for the treatment of urological disease. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/research-and-reports-in-urology-journal>