



University of Southern Denmark

Adverse effects with semaglutide

a protocol for a systematic review with meta-analysis and trial sequential analysis

Sillassen, Christina Dam Bjerregaard; Kamp, Caroline Barkholt; Petersen, Johanne Juul; Faltermeier, Pascal; Siddiqui, Faiza; Grand, Johannes; Dominguez, Helena; Frølich, Anne; Gæde, Peter Haulund; Glud, Christian; Mathiesen, Ole; Jakobsen, Janus

Published in:

BMJ Open

DOI:

10.1136/bmjopen-2024-084190

Publication date:

2024

Document version:

Final published version

Document license:

CC BY-NC

Citation for pulished version (APA):

Sillassen, C. D. B., Kamp, C. B., Petersen, J. J., Faltermeier, P., Siddiqui, F., Grand, J., Dominguez, H., Frølich, A., Gæde, P. H., Glud, C., Mathiesen, O., & Jakobsen, J. (2024). Adverse effects with semaglutide: a protocol for a systematic review with meta-analysis and trial sequential analysis. *BMJ Open*, 14(6), Article e084190. <https://doi.org/10.1136/bmjopen-2024-084190>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.






Unless otherwise specified it has been shared according to the terms for self-archiving.

If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim. Please direct all enquiries to puresupport@bib.sdu.dk

BMJ Open Adverse effects with semaglutide: a protocol for a systematic review with meta-analysis and trial sequential analysis

Christina Dam Bjerregaard Sillassen ^{1,2,3} Caroline Barkholt Kamp ^{1,3}
 Johanne Juul Petersen ^{1,4} Pascal Faltermeier,^{1,5} Faiza Siddiqui ¹
 Johannes Grand ⁶ Helena Dominguez,^{7,8} Anne Frølich,^{9,10} Peter Haulund Gæde,^{2,3}
 Christian Gluud,^{1,3} Ole Mathiesen,^{11,12} Janus Jakobsen^{1,3}

To cite: Sillassen CDB, Kamp CB, Petersen JJ, *et al.* Adverse effects with semaglutide: a protocol for a systematic review with meta-analysis and trial sequential analysis. *BMJ Open* 2024;**14**:e084190. doi:10.1136/bmjopen-2024-084190

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-084190>).

Received 12 January 2024
Accepted 24 May 2024



© Author(s) (or their employer(s)) 2024. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Christina Dam Bjerregaard Sillassen;
christina.sillassen@ctu.dk

ABSTRACT

Introduction Semaglutide is increasingly used for the treatment of type 2 diabetes mellitus, overweight and other conditions. It is well known that semaglutide lowers blood glucose levels and leads to significant weight loss. Still, a systematic review has yet to investigate the adverse effects with semaglutide for all patient groups.

Methods and analysis We will conduct a systematic review and search major medical databases (Cochrane Central Register of Controlled Trials, Medline, Embase, Latin American and Caribbean Health Sciences Literature, Science Citation Index Expanded, Conference Proceedings Citation Index—Science) and clinical trial registries from their inception and onwards to identify relevant randomised clinical trials. We expect to conduct the literature search in July 2024. Two review authors will independently extract data and perform risk-of-bias assessments. We will include randomised clinical trials comparing oral or subcutaneous semaglutide versus placebo. Primary outcomes will be all-cause mortality and serious adverse events. Secondary outcomes will be myocardial infarction, stroke, all-cause hospitalisation and non-serious adverse events. Data will be synthesised by meta-analyses and trial sequential analysis; risk of bias will be assessed with Cochrane Risk of Bias tool—version 2, an eight-step procedure will be used to assess if the thresholds for statistical and clinical significance are crossed, and the certainty of the evidence will be assessed by Grading of Recommendations, Assessment, Development and Evaluations.

Ethics and dissemination This protocol does not present any results. Findings of this systematic review will be published in international peer-reviewed scientific journals. **PROSPERO registration number** CRD42024499511.

INTRODUCTION

Glucagon-like peptide 1 receptor agonists (GLP-1 RAs) are relatively new drugs that mimic the effects of incretin.¹ Glucagon-like peptide 1 is produced by enteroendocrine L cells in the gastrointestinal tract.¹ Some metabolic effects of glucagon-like peptide

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Both risks of random and systematic errors will be taken into account by using a predefined detailed Preferred Reporting Items for Systematic Review and Meta-Analysis Protocols compliant methodology, risk-of-bias assessments using the Cochrane Risk of Bias tool—version 2, an eight-step assessment of the statistical and clinical significance, and trial sequential analysis.
- ⇒ The involvement of two independent investigators for data extraction and risk-of-bias assessments enhances the review's credibility.
- ⇒ The adverse effects of semaglutide might not be disease or dose specific, and pooling all patient groups and types of semaglutide increase the statistical power.
- ⇒ Statistical heterogeneity may be expected as the review includes trials with all types of study participants and all types of semaglutide.
- ⇒ We expect a lack of relevant data, as we expect that a large proportion of the available trials have not focused on assessing adverse effects.

1 are enhancement of glucose-dependent stimulation of insulin secretion, inhibition of glucagon secretion, reduction of gastrointestinal motility and reduction of gastric emptying, thus enhancing satiety and reducing food intake.² Treatment with GLP-1 RAs entails supraphysiological stimulation of the glucagon-like peptide receptor, and therefore, treatment with GLP-1 RAs results in supraphysiological effects of glucagon-like peptide 1.

One type of GLP-1 RAs is semaglutide. Since 2017, semaglutide has been approved by the U.S. Food & Drug Administration for treating patients with type 2 diabetes mellitus.³ In 2021, semaglutide was approved by the U.S. Food & Drug Administration for

treating overweight and obesity.³ Semaglutide is widely and increasingly used for the treatment of overweight or type 2 diabetes mellitus.

Semaglutide is available in both peroral and subcutaneous formulas:

- ▶ The peroral semaglutide is Rybelsus used for the treatment of type 2 diabetes mellitus in the following doses: 3 mg, 7 mg and 14 mg one time a day.⁴
- ▶ Subcutaneous Ozempic is used for treating type 2 diabetes mellitus and off-label usage for treating overweight and obesity in the following doses: 0.25 mg, 0.5 mg, 1 mg or 2 mg one time a week.⁵
- ▶ Subcutaneous Wegovy is used for treating overweight and obesity in the following doses: 0.25 mg, 0.5 mg, 1 mg, 1.7 mg and 2.4 mg one time a week.⁴

Previous evidence

Several previous systematic reviews have shown that semaglutide lowers blood glucose levels and leads to significant weight loss compared with placebo. Most of these systematic reviews have also assessed some of the adverse effects of semaglutide.^{4 6-14} Generally, the most frequent serious and non-serious adverse events have been related to the gastrointestinal tract (see online supplemental material). Animal studies have indicated an association between treatment with GLP-1 RA and pancreatitis, pancreatic cancer and thyroid cancer.¹⁵ Recently, concerns have been raised regarding suicidal behaviour due to semaglutide.¹⁶ The results of previous reviews are conflicting, and none of the systematic reviews included all relevant patient groups. Moreover, the latest published trials investigating semaglutide versus placebo (the ‘Semaglutide Effects on Cardiovascular Outcomes in People With Overweight or Obesity’ (SELECT) trial¹⁷ and the ‘Semaglutide in Patients with Heart Failure with Preserved Ejection Fraction and Obesity’ (STEP-HFpEF) trial¹⁸) have not been included in the previous reviews.^{4 6-14} A detailed description of the previous reviews may be found in online supplemental material.

Why is this review important?

The results of previous reviews assessing the effects of semaglutide are conflicting, and no previous review has included all patient groups treated with semaglutide. The effects of semaglutide on blood glucose levels and weight loss might differ between the different patient groups being treated, but the adverse effects with semaglutide might be comparable. Including all patient groups in the analyses will increase the statistical power if meta-analyses are deemed warranted. There is a lack of a systematic review including all relevant trials assessing the adverse effects with semaglutide.

METHODS AND ANALYSES

The protocol is reported following the reporting guideline provided in the Preferred Reporting Items for Systematic Reviews and Meta-Analysis Protocols (PRISMA-P)

statement^{19 20} and is registered in the PROSPERO database (CRD42024499511).

Criteria for considering studies for this review

Types of trials

We will include randomised clinical trials, irrespective of publication status, publication year and language. We will not include quasi-randomised studies (ie, trials with inappropriate allocation strategies for interventions²¹), non-randomised studies (eg, case-control studies) or observational studies.

Types of participants

All patients treated with oral or subcutaneous semaglutide irrespectively of the indication for the treatment (eg, type 2 diabetes mellitus, obesity, chronic kidney disease, patients with a high risk of a cardiovascular event, women with polycystic ovary syndrome, etc).

Types of interventions

We will accept all dosages of oral and subcutaneous semaglutide.

We will accept placebo or ‘no intervention’ as control interventions.

Cointerventions

We will accept any cointervention, if the cointervention is intended to be delivered similarly in the intervention and control groups.

Outcome measures

Primary outcomes

1. All-cause mortality.
2. Proportion of participants with one or more serious adverse events. We will use the International Conference on Harmonization of technical requirements for registration of pharmaceuticals for human use—Good Clinical Practice (ICH-GCP) definition of a serious adverse event, which is any untoward medical occurrence that resulted in death, was life-threatening, required hospitalisation or prolonging of existing hospitalisation, and resulted in persistent or significant disability or jeopardised the participant.²² If the trialists do not use the ICH-GCP definition, we will include the data if the trialists use the term “serious adverse event”. If the trialists do not use the ICH-GCP definition or the term serious adverse event, we will also include the data if the event clearly fulfils the ICH-GCP definition of a serious adverse event. We will exploratorily assess each type of serious adverse event separately.

Secondary outcomes

1. Myocardial infarction (as defined by trialists).
2. Stroke (as defined by trialists).
3. All-cause hospitalisation (as defined by trialists).
4. Non-serious adverse events. Each type of adverse event will be analysed and presented separately.

Exploratory outcomes

1. Pancreatitis (as defined by trialists).
2. Any type of cancer (as defined by trialists).
3. Suicides or suicide attempts (as defined by trialists).
4. Composite outcome of death from cardiovascular causes, non-fatal myocardial infarction or non-fatal stroke (as defined by trialists).
5. Vision changes (blurred vision, retinopathy or macular complications) (as defined by trialists).

Assessment time points

We will assess all outcomes at maximum follow-up.

Search methods for identification of studies

Electronic searches

We will search Cochrane Central Register of Controlled Trials, Medical Literature Analysis and Retrieval System Online (Medline) (Medline Ovid), Excerpta Medica database (Embase) (Embase Ovid), Latin American and Caribbean Health Sciences Literature (VHL Regional Portal), Science Citation Index Expanded (Web of Science) and Conference Proceedings Citation Index—Science (Web of Science) to identify relevant trials. We will search all databases from their inception to the present. Trials will be included irrespective of language, publication status, publication year and publication type. For a detailed search strategy for all electronic searches, see online supplemental material.

We expect to conduct the literature search in July 2024.

Searching other resources

The reference lists of relevant trial publications will be checked for any unidentified randomised clinical trials. To identify unpublished trials, we will search clinical trial registries (eg, clinicaltrials.gov, clinicaltrialregister.eu, who.int/ictrp, chictr.org.cn) of the World.

We will also include unpublished trials if we identify these and assess relevant retraction statements and errata for included trials. We will also try to obtain clinical study reports from regulatory authorities as these are known to be more transparent in reporting adverse events.^{23–28} We will also search preprint servers (bioRxiv, medRxiv) for unpublished trials.

Data collection

We will perform and report the review as recommended by the PRISMA statement.²⁹ Analyses will be performed using Stata V.16 (StataCorp LLC)³⁰ and trial sequential analysis.^{31 32}

Selection of randomised clinical trials

Two review authors will independently screen titles and abstracts. We will retrieve all relevant full-text study reports/publications, and two review authors will independently screen the full-texts and will record reasons for the exclusion of the ineligible studies. The same two review authors will resolve any disagreements through discussion, or if required, they will consult a third author (JJ).

Data extraction and management

Two review authors will independently extract data from included trials in a predefined form. Disagreements will be resolved by discussion with a third author (JJ). The two review authors will assess duplicate publications and companion papers of a trial together to evaluate all available data simultaneously (maximise data extraction, correct bias assessment). Each trial will be named after the first author and year of the primary publication, and all secondary publications will be classified under that name. We will contact the trial authors by email to specify any missing data, which may not be reported sufficiently or at all in the publication.

Trial characteristics

We will extract the following data: bias risk components (as defined below), trial design (parallel, factorial or crossover), number of intervention groups, length of follow-up, estimation of sample size, and inclusion and exclusion criteria.

Participant characteristics

We will extract the following data: number of randomised participants, number of analysed participants, number of participants lost to follow-up/withdrawals/crossover, age range (mean or median), sex ratio, primary diagnosis (eg, type 2 diabetes mellitus, obesity, hypertension, chronic kidney disease, etc), haemoglobin A1c (mean and SD) and body mass index (mean and SD).

Experimental intervention characteristics

We will extract the following data: type of semaglutide (oral or subcutaneous), dosage and length of treatment period.

Control intervention characteristics

We will extract the following data: type of control (placebo, matching placebo, 'active' placebo, or no intervention), dosage and length of treatment period.

Outcomes

All outcomes listed above will be extracted from each randomised clinical trial. For each outcome, we will identify if there are deviations from intended interventions, if outcomes are missing, inappropriately measured, or selectively reported according to the criteria described later in the 'missing outcome data' bias domain, the 'risk of bias in measurement of the outcome' bias domain and the 'risk of bias in selection of the reported result' bias domain.

Notes

We will search for information regarding industry funding of either personal or academic activities for each trial author. We will judge a publication at high risk of for-profit bias if a trial is sponsored by the industry or if just one author has affiliation to the industry.³³ We will note in the 'Characteristics of included studies' table if outcome data were not reported in a usable way. Two review

authors will independently transfer data into the Stata file.³⁰ Disagreements will be resolved through discussion, or if required, we will consult with a third author (JJ).

Assessment of risk of bias in the included trials

Our bias risk assessment will be based on the Cochrane Risk of Bias tool—version 2 (RoB 2) as recommended in The Cochrane Handbook of Systematic Reviews of Interventions.²¹ We will evaluate the methodology in respect of the following bias domains:

- ▶ Bias arising from the randomisation process.
- ▶ Bias due to deviation from intended interventions (effect of assignment to intervention).
- ▶ Bias due to missing outcome data.
- ▶ Bias in measurement of outcomes.
- ▶ Bias arising from selective reporting of results.
- ▶ Overall assessment of risk of bias.

We will assess the domains ‘deviations from intended interventions’, ‘missing outcome data’, ‘risk of bias in measurement of the outcome’ and ‘risk of bias in selection of the reported result’ for each outcome result. Thus, we can assess the bias risk for each outcome assessed in addition to each trial. The overall risk of bias of a result or trial will be judged to be low if all domains are assessed at low risk of bias. If one or more domains are assessed at either some concerns or high risk of bias, the overall risk of bias will be assessed at high. Our primary conclusions will be based on the results of our primary outcome results with overall low risk of bias. Both our primary and secondary conclusions will be presented in the summary of findings tables.

Differences between the protocol and the review

We will conduct the review according to this published protocol and report any deviations from it in the ‘Differences between the protocol and the review’ section of the systematic review.

Measurement of treatment effect

Dichotomous outcomes

We will calculate risk ratios with 95% CI for dichotomous outcomes, as well as the trial sequential analysis-adjusted CIs (see below).

Dealing with missing data

We will use intention-to-treat data if provided by the trialists.³⁴ We will, as the first option, contact all trial authors to obtain any relevant missing data (ie, for data extraction and for assessment of risk of bias, as specified above).

Dichotomous outcomes

We will not impute missing values for any outcomes in our primary analysis. In our sensitivity analyses (see paragraph below), we will impute data.

Assessment of heterogeneity

We will primarily visually investigate forest plots for signs of heterogeneity. We will second assess the presence of statistical heterogeneity using I^2 statistic (ie,

the percentage of total variation across trials apart from random variation, with 0% indicating no observed heterogeneity and increasing percentages indicating increasing heterogeneity)^{21 35 36} and restricted maximum likelihood method to estimate the heterogeneity variance.^{37 38} We will only conclude that the heterogeneity is high if the between trial variance translates to differences important to patients (based on the effect sizes defined by in the trial sequential analyses, see below). We will investigate heterogeneity through subgroup analyses (see ‘Subgroup analyses and integration of heterogeneity’ section below). We may ultimately decide that a meta-analysis should be avoided if heterogeneity is high.²¹

Assessment of reporting biases

We will use a funnel plot to assess reporting bias if 10 or more trials are included.²¹ We will visually inspect funnel plots to assess the risk of bias. We are aware of the limitations of a funnel plot (ie, a funnel plot assesses small study bias).²¹ From this information, we will assess possible reporting bias. For dichotomous outcomes, we will test asymmetry with the Harbord test³⁹ if τ^2 is less than 0.1 and with the R ucker test if τ^2 is more than 0.1.²¹ In addition, we will assess possible reporting bias for dichotomous outcomes with Egger and Begg tests.^{40 41}

Unit of analysis issues

We will only include randomised clinical trials. For trials using crossover design, only data from the first period will be included,^{21 42} so there will be no unit of analysis issues. We will not include cluster randomised trials.

Meta-analysis

We will undertake the meta-analyses in accordance with the *Cochrane Handbook of Systematic Reviews of Interventions*,²¹ Keus *et al*⁴³ and the eight-step assessment suggested by Jakobsen *et al*.⁴⁴ We will use the statistical software Stata to analyse data.³⁰ We will assess our intervention effects with both a random-effect meta-analysis⁴⁵ and a fixed-effect meta-analysis for each treatment comparison separately.⁴⁶ We will primarily report the most conservative point estimate of the two (highest p-value) and consider the less conservative result a sensitivity analysis.⁴⁴

We will assess a total of two primary outcomes and consider a p value of 0.03 (adjustment per recommendations by Jakobsen *et al*) or less as the threshold for statistical significance.⁴⁴ We will investigate possible heterogeneity through subgroup analyses. We will use the eight-step procedure to assess if the thresholds for significance are crossed.⁴⁴ We will include only the relevant intervention groups (arms) where multiple trial arms are reported in a single trial. If two comparisons are combined in the same meta-analysis, we will split the control group to avoid double counting.

Trial sequential analysis

We wish to control the risks of type I and type II errors. We will, therefore, perform trial sequential analysis on all outcomes, to calculate the required information size (that

is, the number of participants needed in a meta-analysis to detect or reject a certain intervention effect) and the cumulative Z-curve's breach of relevant trial sequential monitoring boundaries.^{31 32 47–53} A more detailed description of trial sequential analysis can be found in the trial sequential analysis manual³¹ and at <http://www.ctu.dk/tsa/>. For dichotomous outcomes, we will estimate the required information size based on the observed proportion of patients with an outcome in the control group (the cumulative proportion of patients with an event in the control groups relative to all patients in the control groups), a relative risk reduction or a relative risk increase of 25%, an alpha of 3% for all outcomes (including exploratory outcomes), a beta of 10% and the observed diversity as suggested by the trials in the meta-analysis.

Subgroup analyses and integration of heterogeneity

Subgroup analyses

We will perform the following subgroup analyses when analysing the primary outcomes (all-cause mortality and serious adverse events).

1. Subgroup analyses of each patient group (eg, type 2 diabetes mellitus, obesity, hypertension, chronic kidney disease, etc).
2. Subgroup analyses of subcutaneous semaglutide compared with oral semaglutide.
3. Subgroup analyses of trials at high risk of bias compared with trials at low risk of bias.
4. Subgroup analyses of different doses of semaglutide (at or above the median compared with below the median).
5. Subgroup analyses of different lengths of intervention (at or above the median compared with below the median).
6. Subgroup analyses of different levels of treatment compliance (high degree of compliance compared with low degree of compliance (as defined by trialists)).

We will use the formal test for subgroup interactions in Stata.³⁰ We will perform relevant unanticipated subgroup analyses, if we identify these.

Sensitivity analysis

To assess the potential impact of the missing data for dichotomous outcomes, we will perform the two following sensitivity analyses on all primary and secondary dichotomous outcomes.

- ▶ 'Best-worst-case' scenario: we will assume that all participants lost to follow-up in the experimental group survived, had no serious adverse events, had no non-serious adverse events and was not hospitalised, and that all those participants lost to follow-up in the control group did not survive, had a serious adverse event, had a non-serious adverse event and was hospitalised.
- ▶ 'Worst-best-case' scenario: we will assume that all participants lost to follow-up in the experimental group did not survive, had a serious adverse event, had a non-serious adverse event, and was hospitalised,

and that all those participants lost to follow-up in the control group survived, had no serious adverse events, had no non-serious adverse events and was not hospitalised.

We will present the results of both scenarios in our review. Other post hoc sensitivity analyses might be warranted if unexpected clinical or statistical heterogeneity is identified during the analysis of the review results.⁴⁴

Summary of findings table

We will create a summary of findings table, including each of the prespecified outcomes. We will use the five Grading of Recommendations, Assessment, Development and Evaluations (GRADE) considerations (bias risk of the trials, consistency of effect, imprecision, indirectness, and publication bias) to assess the certainty of the body of evidence.^{44 54–56} We will assess imprecision using trial sequential analysis by downgrading three levels if the accrued sample is less than 33% of diversity-adjusted required information size (DARIS); two levels if between 33% and 66% of DARIS; one level if more than 66% of DARIS; and no downgrade if the cumulative Z-curve crosses the boundary for benefit, futility or harm. We will justify all decisions to downgrade the certainty of the evidence using footnotes, and we will make comments to aid the reader's understanding of the review where necessary. First, we will present our results in the summary of findings table based on the results from the trials at overall low risk of bias, and second, we will present the results based on all trials.⁵⁷

Patient and public involvement

Patients and the public were not involved in the design, or conduct, or reporting, or dissemination plans of our research.

ETHICS AND DISSEMINATION

This systematic review does not require ethical approval and informed consent, as it does not use identifiable patient data. Findings of this systematic review will be published in international peer-reviewed scientific journals.

DISCUSSION

This systematic review with aggregate data meta-analyses and trial sequential analyses aims to assess the adverse effects of semaglutide versus placebo (both beneficial and harmful effects) on hard binary outcomes, including all patient groups. Primary outcomes will be all-cause mortality and serious adverse events. Secondary outcomes will be myocardial infarction, stroke, all-cause hospitalisation and non-serious adverse events.

This protocol has several strengths. The predefined detailed methodology follows the PRISMA-P statement, the eight-step assessment suggested by Jakobsen *et al*,⁴⁴



trial sequential analysis,^{31 32} and GRADE assessments.⁵⁴ Thus, this protocol considers both risks of design errors, random errors and systematic errors. The adverse effects of semaglutide might not be disease or dose specific, and pooling all patient groups and types of semaglutide increase the statistical power.

Our protocol also has limitations. The primary limitation is the risk of potential statistical heterogeneity as we include all types of participants and all types and doses of semaglutide. Furthermore, we anticipate that a large proportion of the available trials will investigate the effects of semaglutide on either glycaemic control or weight loss as the primary outcome and have less focus on the adverse effects with semaglutide. Insufficient data may complicate the systematic assessments of the adverse effects with semaglutide so we cannot draw definitive conclusions. Some randomised trials will presumably only focus on short-term effects of semaglutide (down to 12 weeks of treatment and then follow-up), which may reduce the likelihood of clinical events occurring within the treatment period.

Undetected low adherence to semaglutide and, thus, undetected lower exposure to semaglutide may be an issue, especially in trials with long treatment period. This may be a potential limitation.

Last, it is a limitation that we will only be able to include aggregate data and not individual patient data.

Author affiliations

¹Centre for Clinical Intervention Research, Rigshospitalet Copenhagen Trial Unit, Copenhagen, Denmark

²Department of Cardiology and Endocrinology, Slagelse Hospital, Slagelse, Denmark

³Department of Regional Health Research, Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark

⁴Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark

⁵MSH Medical School Hamburg University of Applied Sciences and Medical University, Hamburg, Germany

⁶Amager-Hvidovre Hospital, Department of Cardiology, Copenhagen University Hospital, Copenhagen, Denmark

⁷Bispebjerg and Frederiksberg Hospital, Department of Cardiology, Copenhagen University Hospital, Copenhagen, Denmark

⁸Department of Biomedicine, Health Faculty, University of Copenhagen, Copenhagen, Denmark

⁹Innovation and Research Centre for Multimorbidity, Slagelse Hospital, Slagelse, Denmark

¹⁰Section of General Practice, University of Copenhagen Faculty of Health and Medical Sciences, Copenhagen, Denmark

¹¹Department of Anaesthesiology, Zealand University Hospital Koge Centre for Anaesthesiological Research, Koge, Denmark

¹²Department of Clinical Medicine, University of Copenhagen Faculty of Health and Medical Sciences, Copenhagen, Denmark

X Johannes Grand @JohannesGrand

Acknowledgements CBK, JJP, PF, FS, CG, and JJ thank CTU, Centre for Clinical Intervention Research, for employment.

Contributors The study idea, design and conduct were conceived by CDBS and JJ. CDBS authored the initial draft and worked closely with JJ on crafting the first version of the manuscript. Subsequently, CBK, JJP, PF, FS, JG, HD, AF, PHG, CG and OM assisted in refining the study design, critically revised the manuscript and approved the submitted version. CDBS is the guarantor. CDBS, FS, JJP and PF will be responsible for data collection. Analyses will be conducted by CDBS, JJP, FS, PF and JJ.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests PG has received lecture fees for Novo Nordisk, AstraZeneca, Eli Lilly, Bayer, MSD and has served in Advisory Boards for Novo Nordisk, AstraZeneca, Bayer.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

Author note We are anticipating to start the data analysis in 6 months (ie, July 2024).

ORCID iDs

Christina Dam Bjerregaard Sillassen <http://orcid.org/0009-0004-9247-5245>

Caroline Barkholt Kamp <http://orcid.org/0000-0002-7756-4694>

Johanne Juul Petersen <http://orcid.org/0000-0001-9837-1958>

Faiza Siddiqui <http://orcid.org/0000-0002-8358-6259>

Johannes Grand <http://orcid.org/0000-0002-5511-4668>

REFERENCES

- Lebrun LJ, Lenaerts K, Kiers D, *et al*. Enteroendocrine L cells sense LPS after gut barrier injury to enhance GLP-1 secretion. *Cell Rep* 2017;21:1160–8.
- Aaboe K, Krarup T, Madsbad S, *et al*. GLP-1: physiological effects and potential therapeutic applications. *Diabetes Obes Metab* 2008;10:994–1003.
- U.S. Food & Drug Administration. FDA Approves New Drug Treatment for Chronic Weight Management, First Since 2014 [press release], 2021. Available: <https://www.fda.gov/news-events/press-announcements/fda-approves-new-drug-treatment-chronic-weight-management-first-2014>
- Palanisamy S, Yien ELH, Shi LW, *et al*. Systematic review of efficacy and safety of newer antidiabetic drugs approved from 2013 to 2017 in controlling HbA1C in diabetes patients. *Pharmacy (Basel)* 2018;6:57.
- Administration USFD. OZEMPIC (semaglutide) injection, for subcutaneous use, Available: https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/209637s012lbl.pdf
- Zaazouee MS, Hamdallah A, Helmy SK, *et al*. Semaglutide for the treatment of type 2 diabetes mellitus: A systematic review and network meta-analysis of safety and efficacy outcomes. *Diabet Metab Syndr : Clin Res Rev* 2022;16:102511.
- Iannone A, Natale P, Palmer SC, *et al*. Clinical outcomes associated with drugs for obesity and overweight: A systematic review and network meta-analysis of randomized controlled trials. *Diabetes Obes Metab* 2023;25:2535–44.
- Gao X, Hua X, Wang X, *et al*. Efficacy and safety of Semaglutide on weight loss in obese or overweight patients without diabetes: A systematic review and meta-analysis of randomized controlled trials. *Front Pharmacol* 2022;13.
- Shi F-H, Li H, Cui M, *et al*. Efficacy and safety of once-weekly Semaglutide for the treatment of type 2 diabetes: A systematic review and meta-analysis of randomized controlled trials. *Front Pharmacol* 2018;9:576.
- Li X, Qie S, Wang X, *et al*. The safety and efficacy of once-weekly glucagon-like Peptide-1 receptor agonist Semaglutide in patients

- with type 2 diabetes mellitus: a systematic review and meta-analysis. *Endocrine* 2018;62:535–45.
- 11 Vosoughi K, Atieh J, Khanna L, et al. Association of glucagon-like peptide 1 analogs and agonists administered for obesity with weight loss and adverse events: A systematic review and network meta-analysis. *EClinicalMedicine* 2021;42:101213.
 - 12 Zhang R, Hou Q-C, Li B-H, et al. Efficacy and safety of subcutaneous Semaglutide in adults with overweight or obese: a subgroup meta-analysis of randomized controlled trials. *Front Endocrinol (Lausanne)* 2023;14:1132004.
 - 13 Iqbal J, Wu H-X, Hu N, et al. Effect of glucagon-like Peptide-1 receptor agonists on body weight in adults with obesity without diabetes mellitus—a systematic review and meta-analysis of randomized control trials. *Obes Rev* 2022;23:e13435.
 - 14 Xie Z, Hu J, Gu H, et al. Comparison of the efficacy and safety of 10 glucagon-like Peptide-1 receptor agonists as add-on to metformin in patients with type 2 diabetes: a systematic review. *Front Endocrinol (Lausanne)* 2023;14:1244432.
 - 15 Filippatos TD, Panagiotopoulou TV, Elisaf MS. Adverse effects of GLP-1 receptor agonists. *Rev Diabet Stud* 2014;11:202–30.
 - 16 European Medicines Agency. EMA statement on ongoing review of GLP-1 receptor agonists [press release]. 2023. Available: <https://www.ema.europa.eu/en/news/ema-statement-ongoing-review-glp-1-receptor-agonists>
 - 17 Lincoff AM, Brown-Frandsen K, Colhoun HM, et al. Semaglutide and cardiovascular outcomes in obesity without diabetes. *N Engl J Med* 2023;389:2221–32.
 - 18 Kosiborod MN, Abildström SZ, Borlaug BA, et al. Semaglutide in patients with heart failure with preserved ejection fraction and obesity. *N Engl J Med* 2023;389:1069–84.
 - 19 Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015;350:g7647.
 - 20 Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4.
 - 21 Higgins J, Thomas J, Chandler J, et al. Cochrane Handbook for systematic reviews of interventions: Cochrane. 2023. Available: www.training.cochrane.org/handbook
 - 22 Committee for Human Medicinal Products. Guideline for good clinical practice E6(R2), 2016. Available: https://www.ema.europa.eu/en/documents/scientific-guideline/ich-guideline-good-clinical-practice-e6r2-step-5_en.pdf
 - 23 Sharma T, Guski LS, Freund N, et al. Suicidality and aggression during antidepressant treatment: systematic review and meta-analyses based on clinical study reports. *BMJ* 2016;352:i65.
 - 24 Schroll JB, Penninga EI, Gøtzsche PC. Assessment of adverse events in protocols, clinical study reports, and published papers of trials of Orlistat: A document analysis. *PLOS Med* 2016;13:e1002101.
 - 25 Jørgensen L, Gøtzsche PC, Jefferson T. Benefits and harms of the human Papillomavirus (HPV) vaccines: systematic review with meta-analyses of trial data from clinical study reports. *Syst Rev* 2020;9:43.
 - 26 Gøtzsche PC, Jørgensen KJ. EMA's mishandling of an investigation into suspected serious neurological harms of HPV vaccines. *BMJ Evid Based Med* 2022;27:7–10.
 - 27 Hodkinson A, Dietz KC, Lefebvre C, et al. The use of clinical study reports to enhance the quality of systematic reviews: a survey of systematic review authors. *Syst Rev* 2018;7:117.
 - 28 Howick J. Exploring the Asymmetrical relationship between the power of finance bias and evidence. *Perspect Biol Med* 2019;62:159–87.
 - 29 Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 2010;8:336–41.
 - 30 StataCorp. Stata Statistical Software. College Station, TX: StataCorp LLC, 2019. Available: <http://www.stata.com>
 - 31 Thorlund K, Engström J, Wetterslev J, et al. User Manual for Trial Sequential Analysis (TSA) [1-119], 2017. Available: http://ctu.dk/wp-content/uploads/2021/03/2017-10-10-TSA-Manual-ENG_ER.pdf
 - 32 Copenhagen Trial Unit. TSA - trial sequential analysis, Available: <http://www.ctu.dk/tsa>
 - 33 Lundh A, Sismondo S, Lexchin J, et al. Industry sponsorship and research outcome. *Cochrane Database Syst Rev* 2012;12:MR000033.
 - 34 Jakobsen JC, Gluud C, Wetterslev J, et al. When and how should multiple imputation be used for handling missing data in randomised clinical trials – a practical guide with Flowcharts. *BMC Med Res Methodol* 2017;17:162.
 - 35 Higgins JPT, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ* 2003;327:557–60.
 - 36 Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
 - 37 Veroniki AA, Jackson D, Viechtbauer W, et al. Methods to estimate the between-study variance and its uncertainty in meta-analysis. *Res Synth Methods* 2016;7:55–79.
 - 38 Langan D, Higgins JPT, Jackson D, et al. A comparison of heterogeneity variance Estimators in simulated random-effects meta-analyses. *Res Synth Methods* 2019;10:83–98.
 - 39 Harbord RM, Egger M, Sterne JAC. A modified test for small-study effects in meta-analyses of controlled trials with binary endpoints. *Stat Med* 2006;25:3443–57.
 - 40 Egger M, Smith GD, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629–34.
 - 41 Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088–101.
 - 42 Elbourne DR, Altman DG, Higgins JPT, et al. Meta-analyses involving cross-over trials: methodological issues. *Int J Epidemiol* 2002;31:140–9.
 - 43 Keus F, Wetterslev J, Gluud C, et al. Evidence at a glance: error matrix approach for Overviewing available evidence. *BMC Med Res Methodol* 2010;10:90.
 - 44 Jakobsen JC, Wetterslev J, Winkel P, et al. Thresholds for statistical and clinical significance in systematic reviews with meta-analytic methods. *BMC Med Res Methodol* 2014;14:120.
 - 45 DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;7:177–88.
 - 46 Demets DL. Methods for combining randomized clinical trials: strengths and limitations. *Stat Med* 1987;6:341–50.
 - 47 Wetterslev J, Thorlund K, Brok J, et al. Trial sequential analysis may establish when firm evidence is reached in cumulative meta-analysis. *J Clin Epidemiol* 2008;61:64–75.
 - 48 Brok J, Thorlund K, Gluud C, et al. Trial sequential analysis reveals insufficient information size and potentially false positive results in many meta-analyses. *J Clin Epidemiol* 2008;61:763–9.
 - 49 Brok J, Thorlund K, Wetterslev J, et al. Apparently conclusive meta-analyses may be inconclusive—trial sequential analysis adjustment of random error risk due to repetitive testing of accumulating data in apparently conclusive neonatal meta-analyses. *Int J Epidemiol* 2009;38:287–98.
 - 50 Thorlund K, Devereaux PJ, Wetterslev J, et al. Can trial sequential monitoring boundaries reduce spurious inferences from meta-analyses *Int J Epidemiol* 2009;38:276–86.
 - 51 Wetterslev J, Thorlund K, Brok J, et al. Estimating required information size by Quantifying diversity in random-effects model meta-analyses. *BMC Med Res Methodol* 2009;9:86.
 - 52 Thorlund K, Anema A, Mills E. Interpreting meta-analysis according to the adequacy of sample size. an example using isoniazid Chemoprophylaxis for tuberculosis in purified protein derivative negative HIV-infected individuals. *Clin Epidemiol* 2010;2:57–66.
 - 53 Imberger G, Thorlund K, Gluud C, et al. False-positive findings in Cochrane meta-analyses with and without application of trial sequential analysis: an empirical review. *BMJ Open* 2016;6:e011890.
 - 54 Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924–6.
 - 55 Guyatt GH, Oxman AD, Schünemann HJ, et al. GRADE guidelines: a new series of articles in the Journal of clinical epidemiology. *J Clin Epidemiol* 2011;64:380–2.
 - 56 Schünemann HJ, Best D, Vist G, et al. Letters, numbers, symbols and words: how to communicate grades of evidence and recommendations. *CMAJ* 2003;169:677–80.
 - 57 Schünemann H, Brozek J, Guyatt G, eds. GRADE Handbook for grading quality of evidence and strength of recommendations. updated October 2013. 2013. Available: guidelinedevelopment.org/handbook