

Syddansk Universitet

Large N Scalars From Glueballs to Dynamical Higgs Models

Sannino, Francesco

Published in:
Physical Review D

DOI:
[10.1103/PhysRevD.93.105011](https://doi.org/10.1103/PhysRevD.93.105011)

Publication date:
2016

Document version
Early version, also known as pre-print

Citation for published version (APA):
Sannino, F. (2016). Large N Scalars: From Glueballs to Dynamical Higgs Models. Physical Review D, 93(10), 1-8. [105011]. <https://doi.org/10.1103/PhysRevD.93.105011>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Large N Scalars: From Glueballs to Dynamical Higgs Models

Francesco Sannino*

*CP³-Origins & the Danish IAS, University of Southern Denmark,
Campusvej 55, DK-5230 Odense M, Denmark.*

We construct effective Lagrangians, and corresponding counting schemes, valid to describe the dynamics of the lowest lying large N stable massive composite state emerging in strongly coupled theories. The large N counting rules can now be employed when computing quantum corrections via an effective Lagrangian description. The framework allows for systematic investigations of composite dynamics of non-Goldstone nature. Relevant examples are the lightest glueball states emerging in any Yang-Mills theory. We further apply the effective approach and associated counting scheme to composite models at the electroweak scale. To illustrate the formalism we consider the possibility that the Higgs emerges as: the lightest glueball of a new composite theory; the large N scalar meson in models of dynamical electroweak symmetry breaking; the large N pseudodilaton useful also for models of near-conformal dynamics. For each of these realisations we determine the leading N corrections to the electroweak precision parameters. The results nicely elucidate the underlying large N dynamics and can be used to confront first principle lattice results featuring composite scalars with a systematic effective approach.

Preprint: CP³-Origins-2015-035 DNRF90& DIAS-2015-35

I. INTRODUCTION

Strong dynamics continues to pose a formidable challenge. Several analytical and numerical ingenious techniques have been invented, exploited and are routinely used to elucidate some of its physical properties. The large number of underlying colours limit is a time-honoured example [1–3]. It has been extensively used in Quantum Chromodynamics (QCD) and string theory, and it constitutes the backbone of the gauge-gravity duality program. We will further highlight here the power of the large N expansion by introducing a four-dimensional calculable framework permitting to investigate the dynamics of the lightest stable non-Goldstone large N composite state.

t' Hooft and Witten demonstrated that Yang-Mills theories at large number of colours admit an

*Electronic address: sannino@cp3.dias.sdu.dk

effective description in terms of an infinite number of non-interacting absolutely stable hadronic states of arbitrary spin [1–3]. By capitalising on this central result we focus on the physics of the lightest massive scalar state that is known to play an important role in QCD [5–14] and in various models of dynamical electroweak symmetry breaking as summarised in [15]. The framework can also be used to consistently determine quantum corrections to compare with first principle lattice simulations of composite dynamics featuring scalars [16–19].

The paper is organised as follows. In Section II we introduce the effective theory for the lightest massive glueball scalar state emerging within a pure Yang-Mills theory, and provide the associated counting scheme. Here we discuss the intriguing interplay between momentum and large N expansions. The framework goes beyond the glueball example and lays the foundation of the subsequent analyses. We then extend the framework to several models of composite electroweak dynamics in Section III where the Higgs is identified with the lightest composite state. In Section IV we determine the large N dependence, for each model, of the electroweak precision parameters [20] stemming from different dynamical Higgs realisations. We summarise our results in Section V.

II. LARGE N EFFECTIVE THEORY FOR THE LIGHTEST GLUEBALL STATE

Consider the lightest scalar state stemming from an $SU(N)$ pure Yang-Mills theory, which is also expected to be the lowest lying state of the full theory. At infinite number of colours the effective Lagrangian for this state is simply the one of a free scalar field [1–3]

$$L_{GB} = \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 + \mathcal{O}(N^{-1}). \quad (1)$$

It is possible to go beyond the free-field limit by first defining with Λ_H the intrinsic composite scale of the theory that permits to expand the effective Lagrangian both in $1/N$ and ∂^2/Λ_H^2 as follows

$$L_{GB} = \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \sum_{q=0, p=0}^{\infty} V_{q,p} \left(\frac{\partial^2}{\Lambda_H^2} \right)^q \left(\frac{1}{N} \frac{h}{\Lambda_H} \right)^p. \quad (2)$$

Here $V_{q,p}$ are dimensionless coefficients of order unity with $V_{0,0} = 1$. The expansion in $1/N$ takes care of the large N suppression of higher point correlators, while the higher derivative terms take into account integrating out heavier states. Since the heavier states couple via $1/N$ suppressed interactions we must also have $V_{q,0} = 0$ with $q \geq 1$.

To leading order in the double expansion we have:

$$L_{GB} = \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \left[1 + \frac{V_{0,1}}{N} \frac{h}{\Lambda_H} \right]. \quad (3)$$

This shows that the trilinear coupling of the scalar is naturally suppressed in this limit. Expanding a little further we have:

$$L_{GB} = \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \left[1 + \frac{V_{0,1}}{N} \frac{h}{\Lambda_H} + \frac{V_{1,1}}{N} \frac{\partial^2}{\Lambda_H^2} \frac{h}{\Lambda_H} + \frac{V_{0,2}}{N^2} \frac{h^2}{\Lambda_H^2} \right]. \quad (4)$$

We have ordered the terms in such a way that the $1/N$ order counts as ∂^2/Λ_H^2 , however, we can imagine several different situations. For example we can work in the very low momentum region. In this limit we can order $\partial^2/\Lambda_H^2 \sim 1/N^2$ and therefore to the next leading order we drop the derivative terms and obtain

$$L_{GB} = \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \left[1 + \frac{V_{0,1}}{N} \frac{h}{\Lambda_H} + \frac{V_{0,2}}{N^2} \frac{h^2}{\Lambda_H^2} \right]. \quad (5)$$

The effective theory features small self couplings, even though it stems from a highly nonperturbative underlying gauge theory. The glueball mass receives $1/N^2$ corrections at the fundamental theory level. By computing the one loop corrections to the h two-point function one can check that the effective theory correctly reproduces the expected large N corrections. Of course, the effective Lagrangian is not renormalizable in the usual sense but because the $1/N$ and ∂^2/Λ_H^2 ordering it is possible to organise and subtract the divergences order by order in this double expansion. Furthermore the coefficients of this effective theory can be determined, for a given underlying Yang-Mills gauge theory, via lattice simulations [4].

III. LARGE N SCALARS FOR DYNAMICAL HIGGS MODELS

We now extend the framework presented above in order to introduce consistent effective descriptions of dynamical Higgs models. We are not concerned with fitting the latest experimental data but focus instead on elucidating the associated large N dynamics and effective theory structure.

We shall first introduce different examples and then, for each of these example we compute the electroweak precision observables in Section IV, more specifically the S and T parameters [20].

A. The Dynamical Higgs as the lightest Large N glueball

We start by considering the logical possibility that the dynamical Higgs state is the lightest glueball state of a new fundamental composite theory. Besides pure Yang-Mills one can also

consider theories with matter displaying large distance conformality and then add an explicit source of conformal breaking, such as fermion masses. This has been shown to occur via lattice simulations [16] and via controllable perturbative examples [21, 22].

Within this scenario one can envision the newly discovered particle at the Large Hadron Collider (LHC) to be the lightest glueball state of a new Yang-Mills theory with a new N -independent string tension proportional to Λ_H . The scale is not automatically related, here, with the electroweak symmetry breaking scale $v \simeq 246$ GeV or $4\pi v$. Therefore dynamical spontaneous breaking of the electroweak symmetry is triggered by either another strongly coupled sector or, if within the same theory, via a distinct dynamically induced chiral symmetry scale¹. Since the state h is a singlet with respect to the Standard Model (SM) symmetries we can write

$$\begin{aligned}
\mathcal{L}_{\text{Glueball-Higgs}} = & \mathcal{L}_{\text{SM}} + \left(1 + \frac{2r_\pi}{N\Lambda_H} h + \frac{s_\pi}{N^2\Lambda_H^2} h^2 \right) \frac{v^2}{4} \text{Tr} D_\mu U^\dagger D^\mu U \\
& + \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \left[1 + \frac{V_{0,1}}{N} \frac{h}{\Lambda_H} + \frac{V_{0,2}}{N^2} \frac{h^2}{\Lambda_H^2} \right] \\
& - m_t \left(1 + \frac{r_t}{N\Lambda_H} h \right) \left[\bar{q}_L U \left(\frac{1}{2} + T^3 \right) q_R + \text{h.c.} \right] \\
& - m_b \left(1 + \frac{r_b}{N\Lambda_H} h \right) \left[\bar{q}_L U \left(\frac{1}{2} - T^3 \right) q_R + \text{h.c.} \right] + \dots \\
& + \mathcal{O} \left(\frac{1}{4\pi v}, \frac{\partial^2}{\Lambda_H^2} \right)
\end{aligned} \tag{6}$$

where \mathcal{L}_{SM} is the SM Lagrangian without Higgs and Yukawa sectors, the ellipses denote Yukawa interactions for SM fermions other than the top-bottom doublet $q \equiv (t, b)$, and $\mathcal{O}(1/\Lambda_H)$ includes higher-dimensional operators, which are suppressed by powers of $1/\Lambda_H$. In this Lagrangian U is the usual exponential map of the Goldstone bosons produced by the breaking of the electroweak symmetry, $U = \exp(i2\pi^a T^a/v)$, with covariant derivative $D_\mu U \equiv \partial_\mu U - igW_\mu^a T^a U + ig'UB_\mu T^3$, $2T^a$ are the Pauli matrices, with $a = 1, 2, 3$. The kinetic term and potential of the SM Higgs have been replaced by the effective theory for the lightest glueball state. The tree-level SM is recovered for $r_\pi = r_t = r_b = N\frac{\Lambda_H}{v}$ and $s_\pi = N^2\frac{\Lambda_H^2}{v^2}$. Here we will keep these couplings of order unity. Also we have not speculated on the hidden sector providing the link between the new glueball theory and the SM sector, but required it to respect the large N counting for the insertion of an extra Glueball-Higgs. We have also ordered the higher derivatives on h such that they are subleading when compared to the $1/N$ operators retained here.

¹ We remind the reader that in theories with an intact centre group the confining and the chiral scale are well separated [23].

If we consider the infinite number of colours limit of the new Yang-Mills theory first we arrive at a perturbative self-interacting glueball state coupled to the SM also via perturbative couplings. We have, therefore, at our disposal an organisation structure that allows to go beyond the tree-level [24]. We shall investigate the dependence on the number of new colors N in the section dedicated to the electroweak parameters.

B. The Large N physics of the Dynamical Higgs

In time-honoured models of dynamical electroweak symmetry breaking [25, 26] the Higgs can be identified with a fermion-antifermion meson². Depending on the new fermion representation with respect to the underlying gauge group one can have different large N countings [15] such as the Corrigan and Ramond one [27]. The counting is incorporated directly in the pion decay constant $F_{\Pi}^2 = d(R)\Lambda_{TC}^2$ with $d(R)$ the dimension of the technicolor theory and Λ_{TC} an intrinsic scale independent on the number of colours [29, 38].

Let's assume for definitiveness that we have an $SU(\bar{N})$ underlying theory featuring a doublet of techniquarks transforming according to the representation R of the composite theory and therefore we can set $v = F_{\Pi}(\bar{N})$. For a generic N it is sufficient to replace v with $v\sqrt{d/\bar{d}}$. For example for the fundamental representation $d = N$ and $\bar{d} = \bar{N}$. Differently from the previous Glueball-Higgs example here also the pion sector is affected by the large N scaling since the self-interactions among the composite pions are also controlled by N . Choosing for definitiveness the underlying fermions to belong to the fundamental representation we have

$$\begin{aligned}
\mathcal{L}_{TC}(N) = & \mathcal{L}_{SM} + \frac{N}{\bar{N}} \left(1 + \frac{2r_{\pi}}{v} \sqrt{\frac{\bar{N}}{N}} h + \frac{\bar{N} s_{\pi}}{N v^2} h^2 \right) \frac{v^2}{4} \text{Tr} D_{\mu} U^{\dagger} D^{\mu} U \\
& + \frac{1}{2} \partial_{\mu} h \partial^{\mu} h - \frac{m_h^2}{2} h^2 \left[1 + \frac{\sqrt{\bar{N}} V_{0,1}}{\sqrt{N}} \frac{h}{v} + \frac{\bar{N} V_{0,2}}{N} \frac{h^2}{v^2} \right] \\
& - \frac{y_t v}{\sqrt{2\bar{N}}} \sqrt{\bar{N}} \left(1 + \frac{\sqrt{\bar{N}} r_t}{\sqrt{N} v} h \right) \left[\bar{q}_L U \left(\frac{1}{2} + T^3 \right) q_R + \text{h.c.} \right] \\
& - \frac{y_b v}{\sqrt{2\bar{N}}} \sqrt{\bar{N}} \left(1 + \frac{\sqrt{\bar{N}} r_b}{\sqrt{N} v} h \right) \left[\bar{q}_L U \left(\frac{1}{2} - T^3 \right) q_R + \text{h.c.} \right] + \dots \\
& + \mathcal{O} \left(\frac{1}{4\pi v'} \frac{\partial^2}{v^2} \right). \tag{7}
\end{aligned}$$

² This does not always have to be the case, meaning that the lightest state can be, in principle, made by multi-fermion states.

We have therefore:

$$m_W^2 = g^2 v^2 \frac{N}{4N}, \quad m_q = y_q v \sqrt{\frac{N}{2N}}, \quad (8)$$

with q a given quark, g the weak coupling and y_q the Yukawa coupling. Substituting v in terms of m_W and the weak coupling via

$$\sqrt{\frac{N}{N}} v = 2 \frac{m_W}{g}, \quad (9)$$

in the effective Lagrangian we arrive at

$$\begin{aligned} \mathcal{L}_{TC}(N) = & \mathcal{L}_{SM} + \left(1 + g \frac{2r_\pi}{2m_W} h + g^2 \frac{s_\pi}{4m_W^2} h^2 \right) \frac{m_W^2}{g^2} \text{Tr} D_\mu U^\dagger D^\mu U \\ & + \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \left[1 + gV_{0,1} \frac{h}{2m_W} + g^2 V_{0,2} \frac{h^2}{4m_W^2} \right] \\ & - m_t \left(1 + g \frac{r_t}{2m_W} h \right) \left[\bar{q}_L U \left(\frac{1}{2} + T^3 \right) q_R + \text{h.c.} \right] \\ & - m_b \left(1 + g \frac{r_b}{2m_W} h \right) \left[\bar{q}_L U \left(\frac{1}{2} - T^3 \right) q_R + \text{h.c.} \right] + \dots \end{aligned} \quad (10)$$

Nicely the $1/\sqrt{N}$ cost of introducing an extra power of the composite field h is monitored by the corresponding power in g .

C. The large N dynamical pseudo-dilaton

It is possible to imagine that the Higgs state of the SM is associated to the spontaneous breaking of a conformal symmetry. There are several possible realisations according to which the breaking of the conformal dynamics can be either associated to a nonperturbative sector [30, 31] or a perturbative one [21, 22].

The large N counting together with the request to satisfy the conformal relations can be both ensured by imposing

$$N\Lambda_H = f, \quad r_q = r_\pi = s_\pi = 1, \quad (11)$$

with f , in general, a new scale. The Lagrangian in (12) then becomes:

$$\begin{aligned}
\mathcal{L}_{\text{Dilaton}}(N) = & \mathcal{L}_{\text{SM}} + \left(1 + \frac{2h}{N\Lambda_H} + \frac{h^2}{N^2\Lambda_H^2}\right) \frac{v^2}{4} \text{Tr} D_\mu U^\dagger D^\mu U \\
& + \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{m_h^2}{2} h^2 \left[1 + \frac{V_{0,1}}{N} \frac{h}{\Lambda_H} + \frac{V_{0,2}}{N^2} \frac{h^2}{\Lambda_H^2}\right] \\
& - m_t \left(1 + \frac{h}{N\Lambda_H}\right) \left[\bar{q}_L U \left(\frac{1}{2} + T^3\right) q_R + \text{h.c.}\right] \\
& - m_b \left(1 + \frac{h}{N\Lambda_H}\right) \left[\bar{q}_L U \left(\frac{1}{2} - T^3\right) q_R + \text{h.c.}\right] + \dots
\end{aligned} \tag{12}$$

This framework can be immediately extended to any dilatonic-like interpretation of the state h , such as the one coming from a near conformal-like technicolor dynamics where f (Λ_H) is identified with v . Because we would like to investigate the explicit N dependence we hold fix the N independent scale Λ_H . Clearly this limit corresponds to the Glueball-Higgs case with extra constraints for the h couplings.

We are now ready to investigate the first consequences of the large N counting.

IV. S AND T PARAMETERS

As a relevant application of the formalism introduced above we study two important correlators, i.e. the S and T parameters [33] for the three different types of dynamical Higgs models discussed above.

Defining by S the difference between S in the full theory S_{theory} , and the value of S in the SM, i.e. S_{SM} we arrive at [34]

$$\begin{aligned}
S = & \left(1 - \frac{\kappa_1^2}{4}\right) \left[\frac{f(m_Z/m_h)}{6\pi} + \frac{1}{12\pi} \log \frac{(4\pi\Lambda_H)^2}{m_h^2} + \frac{5}{72\pi} \right] + 16\pi c(4\pi\Lambda_H) \\
& + \frac{1}{12\pi} \log \frac{m_h^2}{m_{h,\text{ref}}^2} + \frac{f(m_Z/m_{h,\text{ref}}) - f(m_Z/m_h)}{6\pi}.
\end{aligned} \tag{13}$$

where κ_1 is the coefficient of the linear term in h/v multiplying the operator $\text{Tr} [D_\mu U D^\mu U^\dagger]$ which in the SM is equal to two, and

$$f(x) \equiv \frac{2x^2 + x^4 - 3x^6 + (9x^4 + x^6) \log x}{(1 - x^2)^3}. \tag{14}$$

The c term in (13) is a needed counterterm. For the T parameter one obtains [34]

$$T = -\frac{3}{16\pi \cos^2 \theta_w} \left(1 - \frac{\kappa_1^2}{4}\right) \log \frac{(4\pi\Lambda_H)^2}{m_h^2} + T_{\text{SM}}^h(m_h) - T_{\text{SM}}^h(m_{h,\text{ref}}), \tag{15}$$

where we have absorbed the finite part of the counterterm in the actual value of Λ_H . $m_{h,\text{ref}}$ is the reference value of the Higgs mass, θ_w is the Weinberg angle, and $T_{\text{SM}}^h(m_h)$ is given in equation (22) of [34].

We can now determine the large N behaviour of these relevant parameters coming from the various large N dynamical Higgs models introduced earlier.

A. S and T for the Large N Glueball and Dilaton Higgs Models

In the glueball case we have

$$\kappa_1^{\text{GB}} = 2 \frac{vr_\pi}{N\Lambda_H}, \quad (16)$$

and the SM limit is recovered when $N\Lambda_H = vr_\pi$. The precision parameters are then

$$\begin{aligned} S_{\text{GB}} = & \left(1 - \frac{v^2 r_\pi^2}{N^2 \Lambda_H^2}\right) \left[\frac{f(m_Z/m_h)}{6\pi} + \frac{1}{12\pi} \log \frac{(4\pi\Lambda_H)^2}{m_h^2} + \frac{5}{72\pi} \right] + 16\pi c_{\text{GB}}(4\pi\Lambda_H) \\ & + \frac{1}{12\pi} \log \frac{m_h^2}{m_{h,\text{ref}}^2} + \frac{f(m_Z/m_{h,\text{ref}}) - f(m_Z/m_h)}{6\pi}, \end{aligned} \quad (17)$$

and

$$T_{\text{GB}} = -\frac{3}{16\pi \cos^2 \theta_w} \left(1 - \frac{v^2 r_\pi^2}{N^2 \Lambda_H^2}\right) \log \frac{(4\pi\Lambda_H)^2}{m_h^2} + T_{\text{SM}}^h(m_h) - T_{\text{SM}}^h(m_{h,\text{ref}}). \quad (18)$$

We have chosen $m_h = 125$ GeV and $m_{h,\text{ref}} = 117$ GeV and $c_{\text{GB}}(4\pi\Lambda_H)$ is a counterterm that depends on the specific underlying theory. For the Glueball-Higgs theory we absorb the unknown counterterm in the definition of Λ_H and therefore set it to zero in the numerical evaluation. We expect that higher glueball states will not modify the N counting.

The first observation is that if the glueball-Higgs scale Λ_H is larger than the electroweak scale v there is a positive contribution to the S parameter and an associated negative one for the T parameter. Vice-versa we observe a reduction (increase) of the S (T) parameter if Λ_H is smaller than v . This is an intriguing general result given the fact that the scale of compositeness is $4\pi\Lambda_H$ can be kept above the electroweak scale.

Increasing N while keeping fixed Λ_H and r_π one arrives at the following N -independent results

$$\begin{aligned} \lim_{N \rightarrow \infty} S_{\text{GB}} = & \left[\frac{f(m_Z/m_h)}{6\pi} + \frac{1}{12\pi} \log \frac{(4\pi\Lambda_H)^2}{m_h^2} + \frac{5}{72\pi} \right] + 16\pi c_{\text{GB}}(4\pi\Lambda_H) \\ & + \frac{1}{12\pi} \log \frac{m_h^2}{m_{h,\text{ref}}^2} + \frac{f(m_Z/m_{h,\text{ref}}) - f(m_Z/m_h)}{6\pi}, \end{aligned} \quad (19)$$




FIG. 1: We show the dependence on the number of underlying glueball-Higgs colors for the S (left-panel) and T (right-panel) for $\Lambda_H = 500$ GeV (blue curve), 200 GeV (magenta), 100 GeV (red) and 50 GeV (green). The composite scale $4\pi\Lambda_H$ is always higher than the electroweak scale of 246 GeV, and further assumed $r_\pi \approx 1$.

and

$$\lim_{N \rightarrow \infty} T_{GB} = -\frac{3}{16\pi \cos^2 \theta_w} \log \frac{(4\pi\Lambda_H)^2}{m_h^2} + T_{SM}^h(m_h) - T_{SM}^h(m_{h,ref}). \quad (20)$$

The corrections appear at $\mathcal{O}(N^2)$.

In Figure 1 we plot S and T as function of the number of colors for different values of Λ_H . Because the corrections are in $1/N^2$ the large N limit is approached quickly. We have also assumed $r_\pi \approx 1$ that is its natural order of magnitude and, in any event, can be partially reabsorbed in Λ_H . We compare the result with the experimental value of precision data in Fig. 2 for $\Lambda_H = 200$ GeV (magenta) and $N = 1, 2$. In red we have $\Lambda_H = 100$ GeV and $N = 2, 3, 4$. Finally we plot the 50 GeV (green) case for $N = 4, 5, 6$. The composite scale $4\pi\Lambda_H$ is always higher than the electroweak scale of 246 GeV. The left most point on each curve corresponds to the smallest N . The experiments prefer smaller values of Λ_H with N in the range two to four. Larger values of Λ_H require N to be away from the large N limit and therefore we cannot conclude on the viability of the $\Lambda_H = 200$ GeV case. Increasing further Λ_H it is clearly not preferred by precision observables. If, therefore, a Glueball-Higgs model does describe the Higgs we expect soon new states to be discovered with masses in the range 600 – 1200 GeV.

We stress that by requiring to be in agreement with precision measurements the couplings of the Higgs to the standard model gauge bosons are also close to the experimental values. This occurs because the product $N\Lambda_H$ is constrained to be near the electroweak scale.




FIG. 2: Comparison with the precision electroweak constraints for the glueball-Higgs for $\Lambda_H = 200$ GeV (magenta) and $N = 1, 2, 100$ GeV (red) and $N = 2, 3, 4,$ and 50 GeV (green) for $N = 4, 5, 6$. The composite scale $4\pi\Lambda_H$ is always higher than the electroweak scale of 246 GeV. The left most point on each curve corresponds to the smallest N , and further assumed $r_\pi = 1$.

For the Dilaton-Higgs example we have

$$\kappa_1^{\text{Dilaton}} = 2 \frac{v}{N\Lambda_H}, \quad (21)$$

that corresponds to the results above but now with r_π exactly equal to one.

B. S and T for the large N Dynamical Higgs

It is interesting to explore what happens for the large N dynamical Higgs. The main difference with respect to the previous case is that the electroweak scale and the dynamical Higgs scale are now identified. Among the possible underlying models that can lead to this kind of effective dynamics there are time-honoured examples such as minimal models of (near-conformal) technicolor [35–37].

$$\kappa_1^{TC} = 2r_\pi \sqrt{\frac{N}{N}}, \quad (22)$$

yielding

$$S_{TC} = \left(1 - \frac{\bar{N}}{N} r_\pi^2\right) \left[\frac{f(m_Z/m_h)}{6\pi} + \frac{1}{12\pi} \log \frac{(4\pi v)^2}{m_h^2} + \frac{5}{72\pi} \right] + 16\pi c(4\pi v) \quad (23)$$

$$+ \frac{1}{12\pi} \log \frac{m_h^2}{m_{h,\text{ref}}^2} + \frac{f(m_Z/m_{h,\text{ref}}) - f(m_Z/m_h)}{6\pi}, \quad (24)$$

and

$$T_{TC} = -\frac{3}{16\pi \cos^2 \theta_w} \left(1 - \frac{\bar{N}}{N} r_\pi^2\right) \log \frac{(4\pi v)^2}{m_h^2} + T_{\text{SM}}^h(m_h) - T_{\text{SM}}^h(m_{h,\text{ref}}). \quad (25)$$

Differently from the Glueball-Higgs case we have at our disposal only the N dependence of the effective coupling which goes as $1/N$ for the fundamental representation (chosen here) or as $1/N(N \pm 1)$ for two-index representations [38]. We show in Fig. 3 the comparison with the precision electroweak constraints for the dynamical-Higgs for $v = 246$ GeV and $N = 3, 4, 5, 6$. The left most point on each curve corresponds to the smallest N , and further assumed $r_\pi = 0.9$ (left panel), $r_\pi = 1$ (central panel), $r_\pi \approx 1.1$ (right panel). It is clear from the results that it is possible to abide the electroweak precision constraints for larger number of colours provided that r_π is larger than in the SM.

The computation elucidate another important point, i.e. that at large N the leading corrections, expected to be proportional to N , do not come from the dynamical-Higgs sector but rather from the tree-level exchange of spin one resonances [34]. A simple way to understand this point is to observe that the dynamical-Higgs corrections appear, at the effective Lagrangian level, from loops of h and technipions, i.e. the longitudinal components of the SM gauge bosons. At the more fundamental level these corrections are subleading in N and therefore respect the counting provided by our effective approach. Furthermore these dynamical-Higgs corrections were not taken into account in [33], have been amended in [34] and here we provide the intrinsic N dependence.

V. CONCLUSIONS AND TOP CORRECTIONS

We introduced effective field theories and associated counting schemes to consistently describe the lightest massive large N stable composite scalar state emerging in any theory of composite dynamics. The framework allows for systematic investigations of composite dynamics featuring non-Goldstone (and Goldstone) scalars. As time-honoured examples we discussed the lightest glueball state stemming from Yang-Mills theories. We further applied our effective approach to models of (near-conformal) dynamical electroweak symmetry breaking. In particular we considered the following three possibilities: the Higgs is the lightest glueball of a new composite theory;




FIG. 3: Comparison with the precision electroweak constraints for the dynamical-Higgs for $v = 246$ GeV and $N = 3, 4, 5, 6$. The left most point on each curve corresponds to the smallest N , and further assumed $r_\pi = 0.9$ (left panel), $r_\pi = 1$ (central panel), $r_\pi \approx 1.1$ (right panel).

it is a large N scalar meson in models of dynamical Higgs such as technicolor, and finally we considered it to be a large N pseudodilaton in the form of a conformal compensator. For each of these models, we provided the leading N corrections to the precision parameters.

We observe that it is straightforward to show that in this framework the top corrections to the Glueball and dynamical Higgs mass can be reliably estimated in the large N limit by rescaling r_t in equation (4) of [39, 40] by the opportune power of N , for each model, and simultaneously replacing the cutoff scale by either $4\pi\Lambda_H$ or $4\pi v$.

The results provide useful insight stemming from the large N dynamics of these models and can be viewed as the stepping stone for consistent determination of quantum corrections at the effective Lagrangian level containing massive scalar states. The effective approach is directly applicable also to models of composite Goldstone Higgs dynamics [41, 42] when including the first massive scalar state [40, 43, 44], as well as to investigate interesting flavour properties [45, 46]. Finally holographic studies of the spectrum and large N properties of strongly coupled theories [47–49] can benefit from a model independent large N computation that can be performed with the effective theories constructed here.

Acknowledgments

CP³-Origins is partially supported by the Danish National Research Foundation grant DNRF:90.

-
- [1] G. 't Hooft, "A Planar Diagram Theory for Strong Interactions," Nucl. Phys. B **72**, 461 (1974).
 - [2] E. Witten, "Baryons in the 1/n Expansion," Nucl. Phys. B **160**, 57 (1979).
 - [3] E. Witten, "Large N Chiral Dynamics," Annals Phys. **128**, 363 (1980).
 - [4] B. Lucini and M. Panero, "Introductory lectures to large- N QCD phenomenology and lattice results," Prog. Part. Nucl. Phys. **75**, 1 (2014) [arXiv:1309.3638 [hep-th]].
 - [5] F. Sannino and J. Schechter, "Exploring pi pi scattering in the 1/N(c) picture," Phys. Rev. D **52**, 96 (1995) [hep-ph/9501417].
 - [6] M. Harada, F. Sannino and J. Schechter, "Simple description of pi pi scattering to 1-GeV," Phys. Rev. D **54**, 1991 (1996) [hep-ph/9511335].
 - [7] M. Harada, F. Sannino and J. Schechter, "Comment on 'Confirmation of the sigma meson'," Phys. Rev. Lett. **78**, 1603 (1997) [hep-ph/9609428].
 - [8] M. Harada, F. Sannino and J. Schechter, "Large N(c) and chiral dynamics," Phys. Rev. D **69**, 034005 (2004) [hep-ph/0309206].
 - [9] D. Black, A. H. Fariborz, S. Moussa, S. Nasri and J. Schechter, "Unitarized pseudoscalar meson scattering amplitudes in three flavor linear sigma models," Phys. Rev. D **64**, 014031 (2001) [hep-ph/0012278].
 - [10] I. Caprini, G. Colangelo and H. Leutwyler, "Mass and width of the lowest resonance in QCD," Phys. Rev. Lett. **96**, 132001 (2006) [hep-ph/0512364].
 - [11] R. Garcia-Martin, R. Kaminski, J. R. Pelaez and J. Ruiz de Elvira, "Precise determination of the f₀(600) and f₀(980) pole parameters from a dispersive data analysis," Phys. Rev. Lett. **107**, 072001 (2011) [arXiv:1107.1635 [hep-ph]].
 - [12] D. Parganlija, P. Kovacs, G. Wolf, F. Giacosa and D. H. Rischke, "Meson vacuum phenomenology in a three-flavor linear sigma model with (axial-)vector mesons," Phys. Rev. D **87**, no. 1, 014011 (2013) [arXiv:1208.0585 [hep-ph]].
 - [13] J. R. Pelez, T. Cohen, F. J. Llanes-Estrada and J. Ruiz de Elvira, "Different Kinds of Light Mesons at Large N_c ," Acta Phys. Polon. Supp. **8**, no. 2, 465 (2015).
 - [14] Z. Ghalemovi, F. Giacosa and D. H. Rischke, "Masses of Heavy and Light Scalar Tetraquarks in a Non-Relativistic Quark Model," arXiv:1507.03345 [hep-ph].
 - [15] F. Sannino, "Conformal Dynamics for TeV Physics and Cosmology," Acta Phys. Polon. B **40**, 3533 (2009) [arXiv:0911.0931 [hep-ph]].

- [16] F. Bursa *et al.*, “Improved Lattice Spectroscopy of Minimal Walking Technicolor,” *Phys. Rev. D* **84**, 034506 (2011) [arXiv:1104.4301 [hep-lat]].
- [17] Z. Fodor, K. Holland, J. Kuti, S. Mondal, D. Negradi and C. H. Wong, “Toward the minimal realization of a light composite Higgs,” *PoS LATTICE* **2014**, 244 (2015) [arXiv:1502.00028 [hep-lat]].
- [18] J. Kuti, “The Higgs particle and the lattice,” *PoS LATTICE* **2013**, 004 (2014).
- [19] Z. Fodor, K. Holland, J. Kuti, D. Negradi, C. Schroeder and C. H. Wong, “Can the nearly conformal sextet gauge model hide the Higgs impostor?,” *Phys. Lett. B* **718**, 657 (2012) [arXiv:1209.0391 [hep-lat]].
- [20] M. E. Peskin and T. Takeuchi, “A New constraint on a strongly interacting Higgs sector,” *Phys. Rev. Lett.* **65**, 964 (1990).
- [21] B. Grinstein and P. Uttayarat, “A Very Light Dilaton,” *JHEP* **1107**, 038 (2011) [arXiv:1105.2370 [hep-ph]].
- [22] O. Antipin, M. Mojaza and F. Sannino, “Light Dilaton at Fixed Points and Ultra Light Scale Super Yang Mills,” *Phys. Lett. B* **712**, 119 (2012) [arXiv:1107.2932 [hep-ph]].
- [23] A. Mocsy, F. Sannino and K. Tuominen, “Confinement versus chiral symmetry,” *Phys. Rev. Lett.* **92**, 182302 (2004) [hep-ph/0308135].
- [24] M. Hansen, K. Langaebler, C. Pica and F. Sannino, in preparation.
- [25] S. Weinberg, “Implications of Dynamical Symmetry Breaking,” *Phys. Rev. D* **13**, 974 (1976).
- [26] L. Susskind, “Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory,” *Phys. Rev. D* **20**, 2619 (1979).
- [27] E. Corrigan and P. Ramond, “A Note on the Quark Content of Large Color Groups,” *Phys. Lett. B* **87**, 73 (1979).
- [28] F. Sannino and J. Schechter, “Alternative large $N(c)$ schemes and chiral dynamics,” *Phys. Rev. D* **76**, 014014 (2007) [arXiv:0704.0602 [hep-ph]].
- [29] E. B. Kiritsis and J. Papavassiliou, “An Alternative Large N Limit for QCD and Its Implications for Low-energy Nuclear Phenomena,” *Phys. Rev. D* **42**, 4238 (1990).
- [30] F. Sannino and J. Schechter, “Chiral phase transition for $SU(N)$ gauge theories via an effective Lagrangian approach,” *Phys. Rev. D* **60**, 056004 (1999) [hep-ph/9903359].
- [31] W. D. Goldberger, B. Grinstein and W. Skiba, “Distinguishing the Higgs boson from the dilaton at the Large Hadron Collider,” *Phys. Rev. Lett.* **100**, 111802 (2008) [arXiv:0708.1463 [hep-ph]].
- [32] A. Belyaev, M. S. Brown, R. Foadi and M. T. Frandsen, “The Technicolor Higgs in the Light of LHC Data,” *Phys. Rev. D* **90**, 035012 (2014) [arXiv:1309.2097 [hep-ph]].
- [33] M. E. Peskin and T. Takeuchi, “Estimation of oblique electroweak corrections,” *Phys. Rev. D* **46**, 381 (1992).
- [34] R. Foadi and F. Sannino, “S and T parameters from a light nonstandard Higgs particle,” *Phys. Rev. D* **87**, no. 1, 015008 (2013) [arXiv:1207.1541 [hep-ph]].
- [35] T. Appelquist, P. S. Rodrigues da Silva and F. Sannino, “Enhanced global symmetries and the chiral phase transition,” *Phys. Rev. D* **60**, 116007 (1999) [hep-ph/9906555].
- [36] F. Sannino and K. Tuominen, “Orientifold theory dynamics and symmetry breaking,” *Phys. Rev. D* **71**,

- 051901 (2005) [hep-ph/0405209].
- [37] D. D. Dietrich, F. Sannino and K. Tuominen, "Light composite Higgs from higher representations versus electroweak precision measurements: Predictions for CERN LHC," *Phys. Rev. D* **72**, 055001 (2005) [hep-ph/0505059].
- [38] F. Sannino and J. Schechter, "Alternative large $N(c)$ schemes and chiral dynamics," *Phys. Rev. D* **76**, 014014 (2007) [arXiv:0704.0602 [hep-ph]].
- [39] R. Foadi, M. T. Frandsen and F. Sannino, "125 GeV Higgs boson from a not so light technicolor scalar," *Phys. Rev. D* **87**, no. 9, 095001 (2013) [arXiv:1211.1083 [hep-ph]].
- [40] G. Cacciapaglia and F. Sannino, "Fundamental Composite (Goldstone) Higgs Dynamics," *JHEP* **1404**, 111 (2014) [arXiv:1402.0233 [hep-ph]].
- [41] D. B. Kaplan and H. Georgi, *Phys. Lett. B* **136**, 183 (1984).
- [42] D. B. Kaplan, H. Georgi and S. Dimopoulos, *Phys. Lett. B* **136**, 187 (1984).
- [43] A. Arbey, G. Cacciapaglia, H. Cai, A. Deandrea, S. Le Corre and F. Sannino, "Fundamental Composite Electroweak Dynamics: Status at the LHC," arXiv:1502.04718 [hep-ph].
- [44] G. Cacciapaglia, H. Cai, A. Deandrea, T. Flacke, S. J. Lee and A. Parolini, "Composite scalars at the LHC: the Higgs, the Sextet and the Octet," arXiv:1507.02283 [hep-ph].
- [45] D. Ghosh, R. S. Gupta and G. Perez, "Is the Higgs Mechanism of Fermion Mass Generation a Fact? A Yukawa-less First-Two-Generation Model," arXiv:1508.01501 [hep-ph].
- [46] W. Altmannshofer, S. Gori, A. L. Kagan, L. Silvestrini and J. Zupan, "Uncovering Mass Generation Through Higgs Flavor Violation," arXiv:1507.07927 [hep-ph].
- [47] D. Aren, I. Iatrakis, M. Jrvinen and E. Kiritsis, *JHEP* **1311**, 068 (2013) [arXiv:1309.2286 [hep-ph]].
- [48] T. Alho, M. Jrvinen, K. Kajantie, E. Kiritsis, C. Rosen and K. Tuominen, "A holographic model for QCD in the Veneziano limit at finite temperature and density," *JHEP* **1404**, 124 (2014) [*JHEP* **1502**, 033 (2015)] [arXiv:1312.5199 [hep-ph]].
- [49] T. Alho, M. Jarvinen, K. Kajantie, E. Kiritsis and K. Tuominen, "Quantum and stringy corrections to the equation of state of holographic QCD matter and the nature of the chiral transition," *Phys. Rev. D* **91**, no. 5, 055017 (2015) [arXiv:1501.06379 [hep-ph]].