

## Choroidal thickness and myopia in relation to physical activity during childhood

Lundberg, Kristian ; Jacobsen, Nina; Vestergaard, Anders Højslet; Goldschmidt, Ernst; Peto, Tunde ; Larsen, Mikael; Wedderkopp, Niels; Grauslund, Jakob

*Publication date:*  
2017

*Citation for published version (APA):*

Lundberg, K., Jacobsen, N., Vestergaard, A. H., Goldschmidt, E., Peto, T., Larsen, M., ... Grauslund, J. (2017). *Choroidal thickness and myopia in relation to physical activity during childhood*. Poster session presented at Association for Research in Vision and Ophthalmology Annual Meeting 2017, Baltimore, United States.

### Terms of use

This work is brought to you by the University of Southern Denmark through the SDU Research Portal. Unless otherwise specified it has been shared according to the terms for self-archiving. If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim. Please direct all enquiries to [puresupport@bib.sdu.dk](mailto:puresupport@bib.sdu.dk)

### **Choroidal thickness and myopia in relation to physical activity during childhood**

Kristian Lundberg<sup>1,2</sup>, Nina Jacobsen<sup>3</sup>, Anders Vestergaard<sup>1,2</sup>, Ernst Goldschmidt<sup>4</sup>, Tunde Peto<sup>2,5</sup>, Michael Larsen Niels Wedderkopp<sup>6,7</sup>, Jakob Grauslund<sup>1,2</sup>

1 - Department of Ophthalmology, Odense University Hospital, Odense, Denmark. 2 - Department of Clinical Research, Faculty of Health Sciences, University of Southern Denmark, Odense, Denmark. 3 - Department of Ophthalmology, Rigshospitalet-Glostrup University Hospital, Copenhagen, Denmark. 4 - Danish Institute for Myopia Research, Vedbæk, Denmark. 5 - Centre for Public Health, Queen's University Belfast, United Kingdom. 6 - Research in Childhood Health, Institute for Regional Health Research, University of Southern Denmark. 7 - Sport medicine Clinic, the Orthopedic Department, Hospital of Middelfart, Institute of Regional Health Services Research, University of Southern Denmark, Middelfart, Denmark.

#### **Purpose:**

Decreasing physical activity (PA) has been suggested to be a driving force behind the rapid increase of myopia worldwide. The possible protective effects of PA might be through increased blood flow and subsequent change in thickness of the choroid. The purpose of this study was to correlate PA, myopia, and choroidal thickness (CT).

#### **Methods:**

A prospective study of 307 children from the CHAMPS-study Denmark. Objective data from a GT3X accelerometer (ActiGraph, USA) worn at 4 periods between 2009 and 2015 were used to determine the amount and intensity of PA. Intensity was estimated as counts/minutes, and cut off-points were defined at four intensity levels; sedentary (SED), light (L), moderate (M) and vigorous (V). Eye examinations were performed in 2015 and included autorefractometry in cycloplegia (Tonoref II, Nidek, Japan), axial length (AL) by biometri (Lenstar 900, Haag Streit, Switzerland) and fovea-centered radial scans of 4 sections by enhanced depth imaging optical coherence tomography (EDI-OCT) (Heidelberg Spectralis, Germany). By a validated semi-automated method we measured the CT at 17 targets per eye representing anatomically different choroidal locations (subfoveal, 1 and 3 millimeter in each direction of fovea). Linear regression and slope coefficients of repeated PA were performed to evaluate the association between PA, myopia and CT.

#### **Results:**

Mean age at follow-up was 15.4±0.7 years (range 14.3 to 17.5) and 52.4% were boys. The mean spherical equivalent (SE) was 0.3±1.5 diopters (cycloplegia) and 17.9% were myopic (SE≤-0.5 diopters). The mean AL was 23.5±0.9 mm. The mean subfoveal CT was 369±87 µm. The mean CT for myopic vs. non-myopic eyes was 259±65 µm vs. 354±71 µm (macula), 313±77 µm vs. 382±84 µm (fovea), 306±71 µm vs. 368±80 µm (1 mm zone) and 267±52 µm vs. 312±59 µm (3 mm zone), respectively. All CT measurements were thinner in myopic eyes (p<0.0001) and in boys (p<0.05). By age- and sex-adjusted linear regression there were no associations between PA and SE, AL or any CT measurements. There was no association between accumulated PA and the overall CT for SED, L, M and V PA (2.31 µm/% (p=0.22), -3.99 µm/% (p=0.15), -5.43 µm/% (p=0.57) and -0.53 µm/% (p=0.95), respectively).

#### **Conclusions:**

We found no association between physical activity and the choroidal thickness, axial length or refractive error. However, the choroid was thinner in myopic eyes and in boys. The role of choroidal thickness on the development of refraction should be investigated in prospective studies.