

Syddansk Universitet

Human Achilles tendon glycation and function in diabetes

Coupe, Christian; Svensson, Rene Brüggebusch; Kongsgaard, Mads; Kovanen, Vuokko; Grosset, Jean-Francois; Snorgaard, Ole; Bencke, Jesper; Larsen, Jytte Overgaard; Bandholm, Thomas; Christensen, Tomas Møller; Boesen, Anders Ploug; Helmark, Ida C; Aagaard, Per; Kjaer, Michael; Magnusson, Stig Peter

Published in:
Journal of Applied Physiology

DOI:
[10.1152/jappphysiol.00547.2015](https://doi.org/10.1152/jappphysiol.00547.2015)

Publication date:
2016

Document version
Peer reviewed version

Citation for published version (APA):

Coupe, C., Svensson, R. B., Kongsgaard, M., Kovanen, V., Grosset, J-F., Snorgaard, O., ... Magnusson, S. P. (2016). Human Achilles tendon glycation and function in diabetes. *Journal of Applied Physiology*, 120(2), 130-137. <https://doi.org/10.1152/jappphysiol.00547.2015>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Human Achilles tendon glycation and function in diabetes**

2 Christian Couppé^{1,2*}, Rene Brüggerbusch Svensson^{1*}, Mads Kongsgaard¹, Vuokko Kovanen³, Jean-
3 Francois Grosset^{4,5}, Ole Snorgaard⁶, Jesper Bencke⁷, Jytte Overgaard Larsen⁸, Thomas Bandholm⁹,
4 Tomas Møller Christensen¹⁰, Anders Boesen¹, Ida Carøe Helmark¹, Per Aagaard¹¹, Michael Kjær¹,
5 Stig Peter Magnusson^{1,2}
6

7 ¹IOC Sports Medicine, Department of Orthopedic Surgery M, Bispebjerg Hospital and Center for
8 Healthy Aging, Faculty of Health Sciences, University of Copenhagen, Denmark

9 ²Department of Physical Therapy, Musculoskeletal Rehabilitation Research Unit, Bispebjerg
10 Hospital, Denmark

11 ³Department of Health Sciences, University of Jyväskylä, Jyväskylä, Finland

12 ⁴ CNRS UMR 7338, Biomécanique et Bioingénierie, Université de Technologie de Compiègne,
13 60205 Compiègne cedex, France

14 ⁵ Université Paris 13, Sorbonne Paris Cité, UFR Santé Médecine et Biologie Humaine, 93017
15 Bobigny, France

16 ⁶Department of Endocrinology, Hvidovre University Hospital, Hvidovre, Denmark

17 ⁷Gait Analysis Laboratory, Department of Orthopaedics, Copenhagen University Hospital
18 Hvidovre, Hvidovre, Denmark

19 ⁸Department of Neuroscience and Pharmacology, Faculty of Health Sciences, University of
20 Copenhagen, Denmark

21 ⁹Physical Medicine & Rehabilitation Research-Copenhagen (PMR-C), Dept of Physical Therapy,
22 Denmark; Department of Orthopedic Surgery, Hvidovre Hospital, University of Copenhagen,
23 Denmark; Clinical Research Centre, Hvidovre Hospital, University of Copenhagen, Denmark

24 ¹⁰Department of Endocrinology, Bispebjerg Hospital, Denmark

25 ¹¹Institute of Sports Science and Clinical Biomechanics, SDU Muscle Research Cluster, University
26 of Southern Denmark, Odense, Denmark

27 *These authors contributed equally to this work

28
29 Running Title: The Human Diabetic Tendon
30

31 Corresponding author:

32 Christian Couppé, Institute of Sports Medicine Copenhagen,
33 Bispebjerg Hospital, bldg 8, Bispebjerg Bakke 23, DK-2400
34 Copenhagen NV, Denmark.

35 Tel: + 45 3531 5059, Fax: + 45 3531 2733, E-mail: ccouppe@gmail.com

36
37

38 **Abstract**

39 Diabetic patients have an increased risk of foot ulcers, and glycation of collagen may increase tissue
40 stiffness. We hypothesized that the level of glycemic control (glycation) may affect Achilles tendon
41 stiffness, which can influence gait pattern. We therefore investigated the relationship between
42 collagen glycation, Achilles tendon stiffness parameters and plantar pressure in poorly ($n = 22$) and
43 well ($n = 22$) controlled diabetic patients, including healthy age matched (45-70 yrs) controls ($n =$
44 11). There were no differences in any of outcome parameters (collagen cross-linking or tendon
45 stiffness) between patients with well-controlled and poorly controlled diabetes. The overall effect of
46 diabetes was explored by collapsing the diabetes groups (DB) compared to the controls. Skin
47 collagen cross-linking lysylpyridinoline (LP), hydroxyllysylpyridinoline (HP), (136%, 80%, $P <$
48 0.01) and pentosidine concentrations (55%, $P < 0.05$) were markedly greater in DB. Furthermore,
49 Achilles tendon material stiffness was higher in DB (54%, $P < 0.01$). Notably, DB also
50 demonstrated higher forefoot/ rearfoot peak plantar pressure (PPP)-ratio (33%, $P < 0.01$). Overall,
51 Achilles tendon material stiffness and skin connective tissue cross-linking were greater in diabetic
52 patients compared to controls. The higher foot pressure indicates that material stiffness of tendon
53 and other tissue (e.g skin and joint capsule) may influence on foot gait. The difference in foot
54 pressure distribution may contribute to the development of foot ulcers in diabetic patients.

55

56 Key words: Diabetes, Enzymatic and non-enzymatic collagen cross-linking, Achilles tendon
57 mechanics, Foot ulcer
58

59
60

61 **Introduction**

62 Pathological conditions of the feet remain an extensive clinical problem in persons with diabetes
63 (6), and advanced diabetes ulcerations of the forefoot are the main reason for lower extremity
64 amputations (20). In fact, approximately 25% of all hospital admissions of diabetic patients
65 encompass pathological conditions of the feet, and about 15% of all diabetes patients will develop a
66 foot ulcer (20). In addition to this, Achilles tendon problems are more pronounced in patients with
67 diabetes (1), but it is unknown to what extent this is due to altered tendon tissue properties in
68 diabetes, or rather is secondary to altered gait pattern or skin ulcers.

69 Patients with poorly controlled diabetes have elevated plasma glucose concentrations, and
70 this is associated with the accumulation of AGE (Advanced Glycation Endproducts) derived cross-
71 links in various collagenous tissues such as skin, via the Maillard reaction (37). There is evidence
72 that compromised tissue function is a consequence of such increases in AGE cross-linking (4, 37-
73 39, 46). In vitro experiments have shown that glycation increases tendon stiffness and strength (3,
74 26, 27, 45). Increased collagen and tendon stiffness, due to the accumulation of intermolecular AGE
75 cross-links, has been proposed as a concomitant factor in the development of pathological foot
76 conditions in diabetes (23, 41), but reports on AGE accumulation in the human diabetic tendon is
77 sparse (24, 52).

78 Evidence of mechanical changes in diabetic tendons is currently inconclusive, since the
79 effect of diabetes on animal tendon has been reported to result in increased (2, 35, 42) or decreased
80 (7, 12, 18) stiffness properties. In addition, it has not been investigated if the quality of glycemic
81 control in diabetic patients affects AGE cross-linking and tendon stiffness. At the micro-structural
82 level, the extent to which tendon collagen fibrils are affected by diabetes is also sparsely
83 investigated (23). A few animal studies (3, 43) and a single human study (23) have demonstrated
84 significant changes in tendon fibril morphology (increased fibril density and decreased mean fibril

85 area). The biomechanical consequences of these changes in terms of potential alterations in tendon
86 tissue stiffness currently remain unknown.

87 The influence of Achilles tendon stiffness on gait patterns in diabetic patients is unknown,
88 but elevated Achilles tendon stiffness may well decrease dorsiflexion capacity of the ankle joint,
89 and reduced dorsiflexion has been reported to increase forefoot loading (17). Moreover, excessive
90 plantar pressure has been shown to result in elevated tissue breakdown and delayed wound healing
91 in the foot (41) and could be a risk factor for diabetes related pathological foot conditions (51).
92 Therefore, the purpose of the present study was to investigate the hypothesis that poorly controlled
93 diabetes is associated with greater accumulation of AGE cross-links, greater tendon stiffness and
94 altered gait pattern compared to well-controlled diabetes, that may lead to development of foot
95 ulcers. This hypothesis was tested by examining the concentration of enzymatic and non-enzymatic
96 collagen cross-links in skin and tendon, Achilles tendon stiffness, and the modulation in plantar
97 pressure during gait in poorly and well-controlled diabetic patients compared to healthy age-
98 matched controls.

99

100 **Methods**

101 The present cross-sectional study was designed to compare the effect of glycemic control (based on
102 2 year average HbA1c) in two groups of male diabetic patients (type I and type II) with either well
103 ($n = 22$, HbA1c $< 7.5\%$; WCD) or poorly- ($n = 22$, HbA1c $> 9\%$; PCD) controlled diabetes. The
104 number of type 1 diabetic patients was: 1 in WCD and 3 in PCD. Subject characteristics are shown
105 in Table 1. A smaller healthy control group was also included to provide baseline healthy
106 characteristics ($n = 11$, HbA1c $< 6\%$; CON). Subjects were matched for age (45-70 years) and
107 physical activity. Exclusion criteria in both WCD, PCD and CON included neuropathy of non-
108 diabetic origin, severe neuropathy, foot ulcers, severe arterial insufficiency, arthritis of the ankle or

109 foot, previous foot surgery, previous Achilles tendon rupture, amputations, previous Charcot foot,
110 body mass > 110 kg and use of anti-thrombotic medication. The presence of clinical neuropathy
111 was assessed by use of Semmes-Weinstein 5.07 monofilament exam and biothesiometry. The Ethics
112 Committee of the Capital Region of Denmark approved study (journal number 25543), and all
113 procedures conformed to the Declaration of Helsinki. Written, informed consent was obtained from
114 all subjects prior to study onset.

115

116 *Physical Activity*

117 Physical activity was assessed using the International Physical Activity Questionnaire - (IPAQ,
118 Swedish version translated into Danish) quantified as weekly metabolic equivalent of task-(MET)
119 minutes.

120

121 *Blood Sampling*

122 Blood samples of 10 mL were collected before the test day and sent for standard clinical blood tests
123 for triglycerides, high and low-density lipoprotein cholesterol (HDL and LDL), total cholesterol and
124 HbA1c as a measure of mean glucose load over the previous 2-3 months (14, 40). For the diabetic
125 patients (WCD, PCD), the two-year average HbA1c was also determined based on data from their
126 medical records (3-4 measurements).

127

128 *Biopsy Sampling*

129 After biomechanical testing was performed (details given below) biopsy specimens of the Achilles
130 tendon were obtained in the non-dominant leg at the distal end of the tendon 4 cm proximal to the
131 calcaneus. Using ultrasound imaging, the biopsy site was marked on the skin and under local
132 anesthetic (1% lidocaine) the biopsy was obtained with a 16 gauge Bard Monopty triggered biopsy

133 instrument (C. R. Bard Inc, Covington GA). Skin biopsies were performed using a 4 mm biopsy
134 punch (Miltex, York PA) in the gluteal region under local anesthetic (1% lidocain). Both tendon
135 and skin biopsies were immediately frozen in liquid nitrogen for cross-link analysis, and a small
136 segment from the tendon was also placed in 0.05 M phosphate buffered 2% glutaraldehyde for
137 electron microscopy.

138

139 *Collagen Cross-links*

140 The concentrations of enzymatic cross-links lysylpyridinoline (LP) and hydroxylysylpyridinoline
141 (HP), and non-enzymatic AGE cross-link pentosidine in the biopsy samples were quantified as
142 previously described (8, 32). In brief, the tendon biopsy was hydrolysed in 6 M HCl and run on a
143 reversed-phase high performance liquid chromatography column with detection by
144 autofluorescence. The cross-link content was normalized to total collagen content based on
145 hydroxyproline measurement by 4-dimethylaminobenzaldehyde color reaction after oxidation, as
146 previously described (8, 32). Three tendon and 5 skin biopsies were lost during processing for
147 cross-link analysis.

148

149 *Electron Microscopy*

150 Transmission electron microscopy was performed as previously reported (30, 32). In brief,
151 glutaraldehyde fixated samples were stained en-bloc with OsO₄ and embedded in epon. Ultrathin
152 ($\approx 100\text{nm}$) cross-sections were cut and stained with uranyl acetate and lead citrate. Ten $10 \times 10 \mu\text{m}^2$
153 images were obtained in a random pattern across each section to avoid selection bias. In each image
154 36 unbiased counting frames and an unbiased point grid were used to determine collagen fibril
155 density, volume fraction and size. Five biopsies were lost during processing for electron
156 microscopy.

157

158

159 *Achilles Tendon Morphology*

160 Details of the tendon morphology measurements have previously been published (29). In brief, the
161 subject was sitting with the hip, knee and ankle at 90° and using a 100 mm long ultrasound probe
162 the full length of the free tendon from its insertion on the calcaneus to its fusion with the soleus
163 muscle was imaged in B-mode. Using the ultrasound “shadow” of a long needle, the calcaneus and
164 soleus insertions were marked on the skin with a permanent marker. Three evenly spaced marks
165 were placed between the two ends (proximal, mid and distal), and axial ultrasound images were
166 recorded at each point for determining tendon cross-sectional area (CSA) as previously described
167 (29). The average tendon CSA was calculated and used for analysis. The paired student's t-test
168 (systematic error), Pearson correlation coefficient (strength of relationship) and typical error percent
169 for duplicate measures within day were 0.64, 0.93 and 3% for proximal, 0.70, 0.90 and 4% for mid
170 tendon, and 0.57, 0.90 and 4% for distal tendon. The Achilles tendon moment arm was determined
171 as the distance from the foot axis of rotation (mean of medial and lateral malleoli) to the tendon line
172 of action (mid line between calcaneus and soleus insertion) as previously described (29).

173

174 *Achilles Tendon Mechanical Properties*

175 Mechanical properties of the Achilles tendon were assessed using a method that has previously been
176 described and validated in detail (29). In brief, subjects were seated in a rigid chair with the hip,
177 knee and ankle at 90°. The foot was resting on a footplate with the foot axis of rotation vertically
178 above the plate axis of rotation (see Figure 1). The knee was immobilized by a steel cross-bar to
179 prevent lower limb motion (29). A load cell fixed to the footplate was used to measure the plantar
180 flexor moment. Electromyography (EMG) electrodes were attached to the tibialis anterior and

181 soleus muscles to monitor muscle activation and correct for antagonist co-activation as previously
182 described (29). Achilles tendon deformation was monitored using B-mode ultrasound imaging
183 (Hitachi EUB-6500) with a 100 mm long 10 MHz probe positioned along the tendon to visualize
184 the insertion at the calcaneus and soleus.

185 Achilles tendon mechanics were assessed during slow (10s) isometric plantar flexion ramps
186 to maximum voluntary contraction. Force and EMG were recorded synchronously with ultrasound
187 video (29). To correct the Achilles tendon force for antagonist muscle co-activation, the relationship
188 between tibialis anterior EMG amplitude and its resulting dorsiflexor moment was determined
189 during a maximal isometric dorsiflexion lasting 5 seconds (29).

190 Tendon deformation was obtained from the ultrasound videos by feature tracking of the
191 calcaneus and soleus insertions (29). The force-deformation data were fitted to a 3rd order
192 polynomial and this fit was used for further analysis. Stiffness was measured as the slope over the
193 last 20% of tendon deformation. Material properties - stress, strain and modulus - of the Achilles
194 tendon were obtained by dividing force with the mean tendon CSA and dividing deformation with
195 the initial free tendon length. In order to compare tendon properties at identical load, all parameters
196 were also determined at the largest common tendon force observed across participants. To avoid the
197 highly nonlinear toe region commonly observed in tendon at low load, 7 participants (all from the
198 diabetic groups) with particularly low force production were omitted from this comparison. The
199 decision to omit the data points in these 7 participants were made prior to running any between-
200 group analyses. The selected common tendon force level was 1815 N. Five participants did not
201 complete all morphology and mechanical tests due to logistical reasons.

202

203 *Gait Analysis*

204 Load distribution on the foot during walking was determined using a pressure plate (4 sensors/cm²,
205 Emed, Novel, Germany) integrated into a wooden walking path. Subjects were instructed to walk
206 normally along the path and the pressure plate was hit at the third step after start. The mean pressure
207 distribution during 5 steps from each foot was calculated and pressure distribution was assessed by
208 the forefoot/rearfoot peak plantar pressure ratio (PPP-Ratio). Two participants did not complete gait
209 analyses due to logistical reasons.

210

211 *Data Reduction and Statistics*

212 The study was initially powered for the comparison of the WDC and PDC groups, with the healthy
213 controls (CON) included only as a baseline. Tendon stiffness was considered the primary outcome
214 and sample size was determined to be 21 for an effect size of 0.2 with 80% power and a
215 significance level of 5%. Differences between WCD and PCD were determined by an unpaired two-
216 tailed Student's t-test corrected for unequal variances. No differences were observed between the
217 two diabetic groups for any of the outcome variables related to the hypothesis. For this reason it was
218 decided to also report findings relative to the healthy group as a more exploratory approach, in spite
219 of this group being underpowered. Acknowledging that the study is underpowered, we also report
220 some near-significant trends as a basis for future investigation. Diabetic patients were combined
221 into a merged diabetes (DB) group and subsequently compared to CON using unpaired two-tailed
222 Students t-tests corrected for unequal variances. Pearson product-moment correlation analysis was
223 used to analyze the strength of relationships between variables within the merged diabetes group
224 (DB). $P < 0.05$ was considered significant. Results are reported as mean \pm standard error (SE)
225 unless otherwise reported. Student's t-tests were performed using Excel for Mac 2011 (Microsoft
226 corporation) while all correlation analysis was performed using Prism 6 (Graphpad Software Inc.).
227

228 **Results**

229 *Subject characteristics*

230 Diabetes duration was not different between the WCD and PCD groups. HbA1c concentration was
231 higher in PCD compared to WCD, both at present ($8.9 \pm 1.7\%$ vs. $7.2 \pm 0.9\%$, $P < 0.01$) and as 2-
232 year average ($9.4 \pm 1.4\%$ vs. $6.9 \pm 0.5\%$, $P < 0.01$). Subject characteristics are shown in Table 1.
233 Body mass was greater in DB compared to CON ($P < 0.01$). The difference in IPAQ score was not
234 significant between the groups.

235

236 *Collagen cross-linking*

237 Tendon collagen cross-link data are shown in Table 2. None of the parameters collagen,
238 pentosidine, HP and LP concentration, differed significantly between DB and CON. Tendon
239 pentosidine was positively related to age ($r = 0.42$, $P < 0.01$). Skin collagen cross-link data are
240 shown in Table 2. In contrast to tendon, skin pentosidine ($P < 0.05$), LP ($P < 0.01$) and HP ($P <$
241 0.01) concentrations were higher in DB than CON. Two year HbA1c correlated with skin HP ($r =$
242 0.34 , $p < 0.05$) and pentosidine ($r = 0.31$, $p < 0.05$).

243

244 *Collagen fibril characteristics*

245 Collagen fibril data are shown in Table 2. Tendon fibril density was greater in DB compared to
246 CON ($P < 0.05$).

247

248 *Achilles Tendon Morphology*

249 The Achilles tendon moment arm was greater in DB compared to CON (4.26 ± 0.07 vs. 3.94 ± 0.10
250 cm, $P < 0.05$). However, no other differences were observed between DB and controls with respect

251 to average Achilles tendon CSA (0.73 ± 0.02 vs. 0.79 ± 0.03 cm², $P = 0.23$) or free Achilles tendon
252 length (6.5 ± 0.2 vs. 6.1 ± 0.4 cm, $P = 0.47$).

253

254 *Mechanical Tendon Properties*

255 Mechanical properties of the Achilles tendon at maximum force are shown in Table 3. DB did not
256 differ from CON although there was a trend toward reduced Achilles tendon strain in DB compared
257 to controls (effect size 0.9%, $P = 0.075$). Mechanical properties of the Achilles tendon at largest
258 common force are shown in Table 3. DB had higher Achilles tendon modulus at common force than
259 CON ($P < 0.001$).

260

261 *Gait Analysis*

262 Gait data are shown in Table 3. DB demonstrated greater forefoot/rearfoot PPP-Ratio than CON (P
263 < 0.05).

264

265 **Discussion**

266 To the best of our knowledge the present study is the first to investigate if diabetes in humans is
267 associated with greater Achilles tendon glycation and stiffness, and altered gait. In contrast to our
268 initial hypothesis, we could not demonstrate any differences in collagen cross-linking or
269 biomechanical Achilles tendon stiffness between patients with well-controlled and poorly-
270 controlled diabetes. However, in skin collagen cross-linking (HP, LP and pentosidine
271 concentrations) was markedly greater in diabetic patients compared to healthy age-matched
272 controls. Furthermore, Achilles tendon modulus, which represents the material stiffness after
273 accounting for tendon dimensions, was higher in diabetic patients compared to controls. Notably,
274 diabetic patients also demonstrated higher forefoot/rearfoot peak plantar pressure ratio (PPP-ratio)

275 indicating a more forward distributed loading pattern on the foot. This difference in foot pressure
276 distribution may contribute to the development of foot ulcers in diabetic patients. These findings
277 lend some support to the hypothesis that diabetes leads to increased stiffness in the Achilles tendon
278 and an elevated forefoot pressure.

279

280 *Collagen cross-linking*

281 In diabetes there is an increased rate of non-enzymatic formation of AGE cross-links, which may
282 also affect the protein structure and function of connective tissue such as tendon and skin. In
283 collagen one such cross-link is pentosidine, and in the present study the concentration of
284 pentosidine was greater in skin of diabetic patients, although somewhat surprisingly not elevated in
285 the Achilles tendon. In agreement with the present skin data, previous work on experimental animal
286 and human skin composition also show increased pentosidine concentration (16, 37) and other
287 glycation products with diabetes (5, 13). In contrast, data on cross-links in the diabetic tendon are
288 scarce. A greater glycation in the tendon of diabetic human digastric muscle and diaphragm has
289 been shown, although pentosidine was not measured specifically (24, 52). In diabetic animals,
290 increased glycation of tendon has also been reported (35, 42). The difference between tendon and
291 skin data in the present study may relate to differences in tissue turnover. Tendons have very slow
292 turnover, and may even be maintained throughout adult life (25), while skin has a much more rapid
293 turnover rate (50), as also indicated by the lower pentosidine concentrations presently observed in
294 skin biopsies compared to tendon biopsies. Consequently, pentosidine in tendon most likely
295 represent an average over a longer time period than that of skin, and therefore the relative effect of
296 the period with diabetes may be smaller in tendon tissue.

297 Another factor potentially affecting the pentosidine concentration in Achilles tendons is the
298 level of physical activity of the subjects. It has recently been shown that the pentosidine

299 concentration of the patellar tendon is reduced in elderly life-long regular endurance runners
300 (master athletes) compared to sedentary controls (9), and that resistance training can reduce
301 pentosidine concentration in patellar tendons (28). If loading of tendons can ameliorate AGE
302 accumulation, it may also explain why greater AGE accumulation was observed in the diabetic
303 digastric tendon as previously mentioned, since this tendon is not weight bearing.

304 The present study also revealed markedly greater HP and LP concentrations in the skin of
305 diabetic patients compared to healthy controls. The concomitant greater in glycation and enzymatic
306 cross-links is in agreement with previous reports on skin collagen in diabetic conditions (5).
307 Conversely, in the Achilles tendon we did not observe a similar greater cross-linking (HP and LP (P
308 = 0.10)) with diabetes, which to our knowledge has not previously been examined in human
309 diabetic tendons. A simultaneous greater HP, LP and pentosidine with aging have been
310 demonstrated in the human patellar tendon (8). Based on the 'synchronized' changes in non-
311 enzymatic and enzymatic cross-links reported in both diabetes and aging, it is reasonable to
312 speculate that some mechanistic link(s) may exist between the two cross-linking processes. The
313 finding that serum two-year average HbA1c and skin pentosidine in the present study demonstrated
314 a weak relationship ($r = 0.31$, $P < 0.05$) while this was not the case in the tendon ($r = 0.03$, $P =$
315 0.84). This may indicate that the skin tissue is subjected to a systemic effect of AGEs with less
316 protection by physical activity and mechanical loading, which thereby could lead to greater
317 accumulation of non-enzymatic cross-links in skin compared to tendon. Despite superior glycemic
318 control (Hb1Ac) in WCD compared to PCD there were no differences in any of the collagen cross-
319 linking parameters examined, which is in agreement with observations by Lyons et al. who reported
320 similar skin pentosidine content in type 1 diabetic patients with better glycemic control (34).
321 Monnier et al (37) reported an approximately 20% lower skin pentosidine in diabetic patients with
322 improved glycemic control and considering the absolute difference observed in the present study,

323 there is in fact a similar difference, so the lacking effect may reflect a sample size issue.

324

325 *Collagen fibril morphology*

326 Some studies have reported on tendon microstructural changes in diabetes. Both animal and human
327 studies have reported greater collagen fibril density and decreased mean fibril area (3, 23, 43). The
328 present study revealed a 25% higher fibril density in diabetic patients compared to controls.

329 Furthermore, mean fibril diameter and mean fibril area tended ($P = 0.096$) to be reduced (11%) in
330 diabetic patients compared controls, confirming previous findings (3, 23, 43). Why diabetic tendon
331 collagen fibrils display higher fibril density is unknown. It has been speculated that closer packing
332 density could be a result of AGEs binding together collagen fibrils (3, 33). Another mechanism
333 could be that the higher density is a compensating mechanism for a lower mean fibril diameter
334 thereby maintaining total collagen content and volume fraction in agreement with our findings.
335 However these mechanisms need to be explored further.

336

337 *Achilles Tendon Mechanical Properties*

338 In the present study we observed no difference in Achilles tendon mechanics expressed in absolute
339 terms between WCD and PCD, however a 54% greater Young modulus was observed in diabetic
340 patients compared to healthy controls, indicating that qualitative differences exist between diabetic
341 and healthy Achilles tendon tissue. Diabetes has previously been associated with mechanical
342 changes in different tissues including tendon. In experimental diabetic animals greater stiffness has
343 been extensively reported in non-weight bearing rat-tail tendon (2, 19, 21, 22, 35, 42, 53) and knee
344 ligaments (15). Likewise, in various human non-weight bearing connective tissue such as blood
345 vessels (49) and the lens of the eye (44), it has been reported that diabetes induces greater tissue
346 stiffness. A modest increased stiffness has also been demonstrated in weight bearing diabetic canine

347 patellar tendon under long-term insulin therapy (31). In contrast, lower stiffness of the Achilles
348 tendon has been reported in several experimental diabetic animal studies (7, 12, 18), and this may
349 be attenuated by weight bearing physical activity (11). It was recently shown that Achilles tendon,
350 strains are less during walking in human diabetic patients than in controls, which may indicate that
351 greater tendon stiffness could be related to observed differences in the gait pattern of these patients
352 (10). To our best knowledge the present study is the first to directly measure the mechanical
353 properties of human diabetic Achilles tendons in vivo. Our data show a markedly (54%) higher
354 Achilles tendon material stiffness (modulus) compared to controls, however, absolute tendon
355 stiffness was not significantly different despite it was numerically 27% greater in diabetic patients.
356 The difference between the modulus and stiffness lies in the tendon dimensions, with the diabetic
357 tendon towards a greater tendon length and reduced cross-sectional area (neither significant), which
358 counteracts the greater material stiffness. It is possible that the Achilles tendon dimensions of
359 diabetic patients may have adapted to counteract increased material stiffness in order to maintain
360 functional stiffness, but this hypothesis cannot be addressed by the data obtained in the present
361 study.

362 Cross-linking by AGEs is the likely mechanism underpinning tissue stiffening with diabetes
363 (38), and AGE cross-links have been shown to increase tendon stiffness in vitro, where tendon is
364 incubated with a reducing sugar (26, 27, 45). In the present study the material stiffness of the
365 Achilles tendon was greater with diabetic patients, however no differences were observed in
366 pentosidine or HP, LP cross-link concentrations. In addition, collagen content also did not differ
367 between diabetic patients and healthy controls. The diabetic patients had a higher fibril density, but
368 due to their tendency ($P = 0.096$) to toward a lower fibril size the total volume fraction, and thus the
369 load bearing cross-sectional area was unaltered.

370

371 *Gait*

372 In the present study, diabetic patients demonstrated higher forefoot/rearfoot PPP-ratio indicating
373 increased forefoot loading during walking. This finding is in agreement with our initial hypothesis.
374 A forward shift in pressure could be caused by an increased ankle joint stiffness; however, the
375 hypothesized relation to absolute Achilles tendon stiffness was not observed. As previously
376 discussed, the weight bearing nature of the Achilles tendon may render it less susceptible to diabetic
377 changes than other tissues crossing the joint. Since diabetes is a systemic disease these other tissues
378 are likely also affected and may contribute to overall joint stiffness. One concern could be that the
379 difference in tendon moment arm observed between the two subject groups would influence these
380 findings, however, the moment arm was not correlated to either forefoot/rearfoot PPP-ratio or
381 tendon modulus, respectively. However, the potential influence of the Achilles tendon should not
382 be completely disregarded, since the modulus was greater and there were tendencies for both
383 greater absolute stiffness and reduced strain, and as such, a lack of sensitivity may have prevented.
384 Stiffening of the Achilles tendon material properties combined with the observed tendency for
385 decreased tendon strain (potentially causing reduced dorsiflexor ROM during the late stance phase)
386 could *per se* cause an increased magnitude of forefoot loading, and any systemic glycation effect
387 would likely also stiffen other connective tissues surrounding the joint. Notably, reduced
388 dorsiflexion ability has been shown to increase peak plantar pressure during walking (17) while
389 excessive plantar pressure has been shown to result in accelerated tissue breakdown and delayed
390 wound healing (41).

391

392 *Study Limitations*

393 The present investigation is a cross-sectional case-control study and, therefore, has inherent
394 limitations. Furthermore, while a fairly large number of diabetic patients were recruited, a larger

395 number of control subjects would have improved the statistical strength. In the present study the
396 only measured AGE marker was pentosidine, which constitutes a small fraction of AGE cross-links
397 (47). Even though pentosidine is reported to correlate well with diabetic tissue complications (48),
398 total AGE fluorescence (36, 48) and with more abundant AGEs such as carboxymethyllysine
399 (CML)(4), it is possible that investigating other AGE targets (47) could have provided additional
400 information to help explain the greater Achilles tendon mechanics in our diabetic patients.

401 In vivo mechanical measurements are also affected by several limitations. The tendon load
402 is estimated from external moments, and while muscle activation was partly accounted for by EMG
403 measurements, there are still uncertainty in such measures. In addition, the CSA used for
404 determining tendon stress was measured by ultrasound, which is less precise than for example MRI.
405 Finally, tendon deformation is also determined with ultrasound in 2D and some uncertainty may be
406 present due to out of plane motion. These factors combine to increase the variance of the
407 measurements, but should affect the groups equally.

408 There were differences in the baseline characteristics of the two groups, which could affect
409 the outcome. The diabetic group had a higher body mass, and as would be expected peak plantar
410 pressure did correlate with body mass ($r = 0.23$, $P = 0.1$), the forefoot/rearfoot PPP-ratio was not
411 correlated to body mass ($r = 0.06$, $P = 0.66$). Furthermore, tendon stiffness correlated with body
412 mass ($r = 0.34$, $P = 0.03$), but body mass was not linked to modulus ($r = 0.22$, $P = 0.14$).
413 Moreover, the moment arm in diabetic group was higher than in controls. In the present study, the
414 method used to determine moment arm may have some limitations that could have influenced our
415 results. Using e.g. x-ray would have been more precise. However, we were not able demonstrate
416 that the difference in moment arm correlated with the outcome parameters (Forefoot/rearfoot PPP-
417 ratio: $r = -0.10$, $P = 0.48$, Modulus: $r = -0.16$, $P = 0.30$). In addition, the higher moment arm in
418 the diabetic group would have underestimated modulus and thereby cannot be the reason for the

419 observed increase in the diabetic group. To our knowledge there is no evidence that diabetes results
420 in altered moment arm and so we would believe that the difference observed in the present study is
421 spurious. Altogether, if we include mass and moment arm as confounding factors in an ANOVA,
422 the main findings of increased forefoot/rearfoot PPP-ratio and modulus in the diabetic group remain
423 significant.

424

425 *Conclusions*

426 For the first time it was demonstrated that irrespective of hyperglycemia severity Achilles tendon
427 material stiffness was greater in diabetic patients compared to age-matched healthy controls. The
428 finding that well and poor glycemic controlled diabetic patients did not differ in terms of
429 biomechanical Achilles tendon properties was in contrast to our initial hypothesis. Surprisingly,
430 collagen cross-linking also did not differ in the Achilles tendon of the diabetic patients compared to
431 that of controls. In contrast, when assessed in the skin HP, LP and pentosidine cross-link
432 concentrations were markedly greater in diabetic patients compared to controls. Furthermore,
433 diabetic patients showed higher forefoot/rearfoot PPP-ratio during walking, however, a direct
434 relation to increased Achilles whole tendon stiffness was not found, indicating that altered Achilles
435 tendon material stiffness and possibly also in other tissues (e.g skin and joint capsule) may
436 influence plantar pressure distribution during gait habitual walking. Collectively, our data suggest
437 that both the material stiffness of the Achilles tendon and foot pressure distribution are altered in
438 diabetic patients. Such changes in tendon material properties and loading may have implications for
439 the development of diabetic foot ulcers.

440

441 *Acknowledgements*

442 The present study was supported by the Center for Healthy Aging, the University of Copenhagen,

443 the Danish Physiotherapy Association, and the Danish Medical Research Council.

444

445

446

447

448

449
450

451 Referenceliste

- 452 1. **Abate M, Schiavone C, Salini V, and Andia I.** Occurrence of tendon pathologies in
453 metabolic disorders. *Rheumatology (Oxford)* 52: 599-608, 2013.
- 454 2. **Andreassen TT, Seyer-Hansen K, and Bailey AJ.** Thermal stability, mechanical
455 properties and reducible cross-links of rat tail tendon in experimental diabetes. *Biochim Biophys*
456 *Acta* 677: 313-317, 1981.
- 457 3. **Bai P, Phua K, Hardt T, Cernadas M, and Brodsky B.** Glycation alters collagen fibril
458 organization. *Connect Tissue Res* 28: 1-12, 1992.
- 459 4. **Baynes JW.** From life to death--the struggle between chemistry and biology during aging:
460 the Maillard reaction as an amplifier of genomic damage. *Biogerontology* 1: 235-246, 2000.
- 461 5. **Buckingham B, and Reiser KM.** Relationship between the Content of Lysyl Oxidase
462 Dependent Cross-Links in Skin Collagen, Nonenzymatic Glycosylation, and Long-Term
463 Complications in Type-I Diabetes-Mellitus. *J Clin Invest* 86: 1046-1054, 1990.
- 464 6. **Cavanagh PR, Lipsky BA, Bradbury AW, and Botek G.** Treatment for diabetic foot
465 ulcers. *Lancet* 366: 1725-1735, 2005.
- 466 7. **Connizzo BK, Bhatt PR, Liechty KW, and Soslowsky LJ.** Diabetes Alters Mechanical
467 Properties and Collagen Fiber Re-Alignment in Multiple Mouse Tendons. *Ann Biomed Eng* 2014.
- 468 8. **Coupe C, Hansen P, Kongsgaard M, Kovanen V, Suetta C, Aagaard P, Kjaer M, and**
469 **Magnusson SP.** Mechanical properties and collagen cross-linking of the patellar tendon in old and
470 young men. *J Appl Physiol* 107: 880-886, 2009.
- 471 9. **Coupe C, Svensson RB, Grosset JF, Kovanen V, Nielsen RH, Olsen MR, Larsen JO,**
472 **Praet SF, Skovgaard D, Hansen M, Aagaard P, Kjaer M, and Magnusson SP.** Life-long
473 endurance running is associated with reduced glycation and mechanical stress in connective tissue.
474 *Age (Dordr)* 36: 9665, 2014.
- 475 10. **Cronin NJ, Peltonen J, Ishikawa M, Komi PV, Avela J, Sinkjaer T, and Voigt M.**
476 Achilles tendon length changes during walking in long-term diabetes patients. *Clin Biomech*
477 *(Bristol, Avon)* 25: 476-482, 2010.
- 478 11. **de Oliveira RR, Bezerra MA, de Lira KD, Novaes KA, Teixeira MF, Chaves Cde C,**
479 **and Moraes SR.** Aerobic physical training restores biomechanical properties of Achilles tendon in
480 rats chemically induced to diabetes mellitus. *J Diabetes Complications* 26: 163-168, 2012.
- 481 12. **de Oliveira RR, de Lira KD, Silveira PV, Coutinho MP, Medeiros MN, Teixeira MF,**
482 **and de Moraes SR.** Mechanical properties of achilles tendon in rats induced to experimental
483 diabetes. *Ann Biomed Eng* 39: 1528-1534, 2011.
- 484 13. **Dunn JA, McCance DR, Thorpe SR, Lyons TJ, and Baynes JW.** Age-dependent
485 accumulation of N epsilon-(carboxymethyl)lysine and N epsilon-(carboxymethyl)hydroxylysine in
486 human skin collagen. *Biochemistry* 30: 1205-1210, 1991.
- 487 14. **Dunn PJ, Cole RA, and Soeldner JS.** Further development and automation of a high
488 pressure liquid chromatography method for the determination of glycosylated hemoglobins.
489 *Metabolism* 28: 777-779, 1979.
- 490 15. **Duquette JJ, Grigg P, and Hoffman AH.** The effect of diabetes on the viscoelastic
491 properties of rat knee ligaments. *J Biomech Eng* 118: 557-564, 1996.
- 492 16. **Dyer DG, Dunn JA, Thorpe SR, Bailie KE, Lyons TJ, McCance DR, and Baynes JW.**
493 Accumulation of Maillard reaction products in skin collagen in diabetes and aging. *J ClinInvest* 91:
494 2463-2469, 1993.

495 17. **Fernando DJ, Masson EA, Veves A, and Boulton AJ.** Relationship of limited joint
496 mobility to abnormal foot pressures and diabetic foot ulceration. *Diabetes Care* 14: 8-11, 1991.

497 18. **Fox AJ, Bedi A, Deng XH, Ying L, Harris PE, Warren RF, and Rodeo SA.** Diabetes
498 mellitus alters the mechanical properties of the native tendon in an experimental rat model. *J*
499 *Orthop Res* 29: 880-885, 2011.

500 19. **Galeski A, Kastelic J, Baer E, and Kohn RR.** Mechanical and structural changes in rat tail
501 tendon induced by alloxan diabetes and aging. *J Biomech* 10: 775-782, 1977.

502 20. **Giurini JM.** Diabetic ulcers. *J Am Podiatr Med Assoc* 82: 594, 1992.

503 21. **Golub LM, Greenwald RA, Zebrowski EJ, and Ramamurthy NS.** The effect of
504 experimental diabetes on the molecular characteristics of soluble rat-tail tendon collagen. *Biochim*
505 *Biophys Acta* 534: 73-81, 1978.

506 22. **Gonzalez AD, Gallant MA, Burr DB, and Wallace JM.** Multiscale analysis of
507 morphology and mechanics in tail tendon from the ZDSD rat model of type 2 diabetes. *J Biomech*
508 47: 681-686, 2014.

509 23. **Grant WP, Sullivan R, Sonenshine DE, Adam M, Slusser JH, Carson KA, and Vinik**
510 **AI.** Electron microscopic investigation of the effects of diabetes mellitus on the Achilles tendon. *J*
511 *Foot Ankle Surg* 36: 272-278; discussion 330, 1997.

512 24. **Hamlin CR, Kohn RR, and Luschin JH.** Apparent accelerated aging of human collagen in
513 diabetes mellitus. *Diabetes* 24: 902-904, 1975.

514 25. **Heinemeier KM, Schjerling P, Heinemeier J, Magnusson SP, and Kjaer M.** Lack of
515 tissue renewal in human adult Achilles tendon is revealed by nuclear bomb 14C. *FASEB J* 2013.

516 26. **Kent MJ, Light ND, and Bailey AJ.** Evidence for glucose-mediated covalent cross-linking
517 of collagen after glycosylation in vitro. *Biochem J* 225: 745-752, 1985.

518 27. **Kohn RR, Cerami A, and Monnier VM.** Collagen aging in vitro by nonenzymatic
519 glycosylation and browning. *Diabetes* 33: 57-59, 1984.

520 28. **Kongsgaard M, Kovanen V, Aagaard P, Doessing S, Hansen P, Laursen AH, Kaldau**
521 **NC, Kjaer M, and Magnusson SP.** Corticosteroid injections, eccentric decline squat training and
522 heavy slow resistance training in patellar tendinopathy. *ScandJMedSciSports* 19: 790-802, 2009.

523 29. **Kongsgaard M, Nielsen CH, Hegnsvad S, Aagaard P, and Magnusson SP.** Mechanical
524 properties of the human Achilles tendon, in vivo. *Clin Biomech (Bristol, Avon)* 26: 772-777, 2011.

525 30. **Kongsgaard M, Qvortrup K, Larsen J, Aagaard P, Doessing S, Hansen P, Kjaer M,**
526 **and Magnusson SP.** Fibril morphology and tendon mechanical properties in patellar tendinopathy:
527 effects of heavy slow resistance training. *AmJSports Med* 38: 749-756, 2010.

528 31. **Lancaster RL, Haut RC, and DeCamp CE.** Changes in the mechanical properties of
529 patellar tendon preparations of spontaneously diabetic dogs under long-term insulin therapy. *J*
530 *Biomech* 27: 1105-1108, 1994.

531 32. **Lenskjold A, Kongsgaard M, Larsen JO, Nielsen RH, Kovanen V, Aagaard P, Kjaer**
532 **M, and Magnusson SP.** The influence of physical activity during youth on structural and
533 functional properties of the Achilles tendon. *Scand J Med Sci Sports* 2013.

534 33. **Li Y, Fessel G, Georgiadis M, and Snedeker JG.** Advanced glycation end-products
535 diminish tendon collagen fiber sliding. *Matrix Biol* 2013.

536 34. **Lyons TJ, Bailie KE, Dyer DG, Dunn JA, and Baynes JW.** Decrease in skin collagen
537 glycation with improved glycemic control in patients with insulin-dependent diabetes mellitus. *J*
538 *Clin Invest* 87: 1910-1915, 1991.

539 35. **McLennan S, Yue DK, Marsh M, Swanson B, Delbridge L, Reeve T, and Turtle JR.**
540 The prevention and reversibility of tissue non-enzymatic glycosylation in diabetes. *Diabet Med* 3:
541 141-146, 1986.

- 542 36. **Meerwaldt R, Graaff R, Oomen PH, Links TP, Jager JJ, Alderson NL, Thorpe SR,**
543 **Baynes JW, Gans RO, and Smit AJ.** Simple non-invasive assessment of advanced glycation
544 endproduct accumulation. *Diabetologia* 47: 1324-1330, 2004.
- 545 37. **Monnier VM, Bautista O, Kenny D, Sell DR, Fogarty J, Dahms W, Cleary PA, Lachin**
546 **J, and Genuth S.** Skin collagen glycation, glycooxidation, and crosslinking are lower in subjects
547 with long-term intensive versus conventional therapy of type 1 diabetes: relevance of glycated
548 collagen products versus HbA1c as markers of diabetic complications. DCCT Skin Collagen
549 Ancillary Study Group. Diabetes Control and Complications Trial. *Diabetes* 48: 870-880, 1999.
- 550 38. **Monnier VM, Kohn RR, and Cerami A.** Accelerated age-related browning of human
551 collagen in diabetes mellitus. *Proc Natl Acad Sci U S A* 81: 583-587, 1984.
- 552 39. **Monnier VM, Sell DR, and Genuth S.** Glycation products as markers and predictors of the
553 progression of diabetic complications. *AnnNYAcad Sci* 1043: 567-581, 2005.
- 554 40. **Mosca A, Carpinelli A, and Bonini P.** Automated determination of glycated hemoglobins
555 with a new high-performance liquid chromatography analyzer. *Clin Chem* 32: 202-203, 1986.
- 556 41. **Mueller MJ, Diamond JE, Delitto A, and Sinacore DR.** Insensitivity, limited joint
557 mobility, and plantar ulcers in patients with diabetes mellitus. *Phys Ther* 69: 453-459; discussion
558 459-462, 1989.
- 559 42. **Mustata GT, Rosca M, Biemel KM, Reihl O, Smith MA, Viswanathan A, Strauch C,**
560 **Du Y, Tang J, Kern TS, Lederer MO, Brownlee M, Weiss MF, and Monnier VM.** Paradoxical
561 effects of green tea (*Camellia sinensis*) and antioxidant vitamins in diabetic rats: improved
562 retinopathy and renal mitochondrial defects but deterioration of collagen matrix glycooxidation and
563 cross-linking. *Diabetes* 54: 517-526, 2005.
- 564 43. **Odetti P, Aragno I, Rolandi R, Garibaldi S, Valentini S, Cosso L, Traverso N,**
565 **Cottalasso D, Pronzato MA, and Marinari UM.** Scanning force microscopy reveals structural
566 alterations in diabetic rat collagen fibrils: role of protein glycation. *Diabetes Metab Res Rev* 16: 74-
567 81, 2000.
- 568 44. **Pedrigi RM, David G, Dziezyc J, and Humphrey JD.** Regional mechanical properties and
569 stress analysis of the human anterior lens capsule. *Vision Res* 47: 1781-1789, 2007.
- 570 45. **Reddy GK, Stehno-Bittel L, and Enwemeka CS.** Glycation-induced matrix stability in the
571 rabbit achilles tendon. *ArchBiochemBiophys* 399: 174-180, 2002.
- 572 46. **Schnider SL, and Kohn RR.** Glucosylation of human collagen in aging and diabetes
573 mellitus. *J Clin Invest* 66: 1179-1181, 1980.
- 574 47. **Sell DR, Biemel KM, Reihl O, Lederer MO, Strauch CM, and Monnier VM.**
575 Glucosepane is a major protein cross-link of the senescent human extracellular matrix. Relationship
576 with diabetes. *J Biol Chem* 280: 12310-12315, 2005.
- 577 48. **Sell DR, Lapolla A, Odetti P, Fogarty J, and Monnier VM.** Pentosidine formation in skin
578 correlates with severity of complications in individuals with long-standing IDDM. *Diabetes* 41:
579 1286-1292, 1992.
- 580 49. **Sims TJ, Rasmussen LM, Oxlund H, and Bailey AJ.** The role of glycation cross-links in
581 diabetic vascular stiffening. *Diabetologia* 39: 946-951, 1996.
- 582 50. **Verzijl N, DeGroot J, Thorpe SR, Bank RA, Shaw JN, Lyons TJ, Bijlsma JW, Lafeber**
583 **FP, Baynes JW, and TeKoppele JM.** Effect of collagen turnover on the accumulation of advanced
584 glycation end products. *J BiolChem* 275: 39027-39031, 2000.
- 585 51. **Veves A, Murray HJ, Young MJ, and Boulton AJ.** The risk of foot ulceration in diabetic
586 patients with high foot pressure: a prospective study. *Diabetologia* 35: 660-663, 1992.
- 587 52. **Vogt BW, Schleicher ED, and Wieland OH.** epsilon-Amino-lysine-bound glucose in
588 human tissues obtained at autopsy. Increase in diabetes mellitus. *Diabetes* 31: 1123-1127, 1982.

589 53. **Yue DK, McLennan S, Delbridge L, Handelsman DJ, Reeve T, and Turtle JR.** The
590 thermal stability of collagen in diabetic rats: correlation with severity of diabetes and non-
591 enzymatic glycosylation. *Diabetologia* 24: 282-285, 1983.
592

593

594 **Tables**

595

	WCD	PCD	DB	CON
Number of participants	22	22	44	11
Age (yrs)	60 ± 7	58 ± 7	59 ± 7	58 ± 5
Height (cm)	177 ± 5	180 ± 6	178 ± 6	177 ± 5
Mass (kg)	91 ± 13	96 ± 10	93 ± 12**	83 ± 8
BMI (kg·m ⁻²)	29 ± 4	30 ± 4	29 ± 6	27 ± 3
Diabetes duration (yr)	12 ± 6	15 ± 8	13 ± 7	-
HbA _{1c} 2yr average (%)	6.9 ± 0.5	9.4 ± 1.4###	8.1 ± 0.3	-
(mmol·mol ⁻¹)	51 ± 6	79 ± 16	65 ± 3	-
HbA _{1c} present (%)	7.2 ± 0.9**	8.9 ± 1.7###, **	8.0 ± 0.3**	5.5 ± 0.3
(mmol·mol ⁻¹)	61 ± 9	73 ± 18	64 ± 3	36 ± 4
Triglyceride (mmol·l ⁻¹)	1.7 ± 0.3	1.7 ± 0.2	1.7 ± 0.1	1.7 ± 0.4
Total cholesterol (mmol·l ⁻¹)	4.6 ± 0.3	4.9 ± 0.2	4.7 ± 0.2	5.6 ± 0.4
HDL Cholesterol (mmol·l ⁻¹)	1.30 ± 0.09	1.21 ± 0.10	1.25 ± 0.07	1.37 ± 0.13
LDL Cholesterol (mmol·l ⁻¹)	2.4 ± 0.2	2.9 ± 0.1	2.7 ± 0.2*	3.6 ± 0.3
IPAQ (MET Score)	2300 ± 1800	1700 ± 1800	2000 ± 300	1400 ± 900

596 **Table 1 - Subject characteristics. WCD = well-controlled diabetic patients, PCD = poorly-controlled diabetic**
597 **patients, DB = merged diabetic patients (WCD + PCD), CON = healthy, age -matched controls.** Data are given as
598 mean ± SD). Different from WCD, #*P* < 0.05, ###*P* < 0.01*Different from CON, **P* < 0.05. ***P* < 0.01.
599

600
601
602
603
604
605

606
607

	WCD	PCD	DB	CON
Tendon Composition				
Number of participants	21	21	42	10
Collagen (mg·mg ⁻¹ dry wt)	0.73 ± 0.03	0.70 ± 0.02	0.72 ± 0.03	0.75 ± 0.03
Hydroxylysyl pyridinoline (HP, mmol·mol ⁻¹ collagen)	1230 ± 80	1340 ± 70	1250 ± 50	1220 ± 80
Lysyl pyridinoline (LP, mmol·mol ⁻¹ collagen)	52 ± 3	53 ± 3	52 ± 2 (*) P = 0.101	43 ± 5
Pentosidine, (mmol·mol ⁻¹ collagen)	33 ± 2	30 ± 3	31 ± 2	28 ± 2
Skin Composition				
Number of Participants	21	20	41	9
Collagen (mg·mg ⁻¹ dry wt)	0.62 ± 0.01	0.64 ± 0.02	0.63 ± 0.01	0.65 ± 0.02
Hydroxylysyl pyridinoline (HP, mmol·mol ⁻¹ collagen)	35 ± 10	54 ± 10	45 ± 6**	19 ± 4
Lysyl pyridinoline (LP, mmol·mol ⁻¹ collagen)	8 ± 1	9 ± 2	9 ± 1**	5 ± 1
Pentosidine, (mmol·mol ⁻¹ collagen)	13 ± 2	16 ± 2	14 ± 1*	9 ± 2
Tendon Fibril Morphology				
Number of Participants	18	22	40	10
Volume fraction (%)	53 ± 2	54 ± 1	53 ± 1	57 ± 2
Density (#fibril·μm ⁻²)	132 ± 10	130 ± 11	131 ± 7*	105 ± 8
Mean fibril diameter (nm)	64 ± 4	65 ± 3	64 ± 2(*) P = 0.096	73 ± 14
Mean fibril area (nm ²)	4300 ± 500	4400 ± 400	4400 ± 300	5500 ± 600

Table 2. Tendon collagen cross-link and fibril composition. Data are given as mean ± SE. Different from CON, *P < 0.05, ** P < 0.01. Compared with CON (*).

608
609
610
611

	WCD	PCD	DB	CON
Achilles Tendon Mechanics				
(At maximum force)				
Number of participants	20	21	41	9
Deformation (mm)	1.80 ± 0.2	1.9 ± 0.1	1.9 ± 0.1	2.3 ± 0.3
Max force (N)	2600 ± 200	2400 ± 200	2500 ± 200	2800 ± 200
Stiffness (kN·mm ⁻¹)	3.4 ± 0.3	3.4 ± 0.1	3.4 ± 0.3	3.1 ± 0.5
Stress (MPa)	41 ± 5	36 ± 3	39 ± 3	40 ± 3
Strain (%)	2.8 ± 0.3	2.7 ± 0.2	2.8 ± 0.2(*) <i>P</i> = 0.075	3.7 ± 0.4
Modulus (GPa)	3.1 ± 0.2	3.2 ± 0.4	3.1 ± 0.3	2.5 ± 0.3
Achilles Tendon Mechanics				
(At common force)				
Number of Participants	17	17	34	9
Deformation (mm)	1.6 ± 0.3	1.6 ± 0.2	1.6 ± 0.2	1.9 ± 0.3
Stiffness (kN·mm ⁻¹)	2.7 ± 0.3	2.7 ± 0.3	2.7 ± 0.2	2.0 ± 0.4
Stress (MPa)	28 ± 2	25 ± 1	27 ± 1	26 ± 1
Strain (%)	2.5 ± 0.4	2.5 ± 0.2	2.5 ± 0.2	3.2 ± 0.4
Modulus (GPa)	2.5 ± 0.2	2.5 ± 0.3	2.5 ± 0.2**	1.7 ± 0.1
Foot pressure mapping				
Number of Participants	21	22	43	10
Peak Plantar Pressure (PPP) (kPa)	650 ± 40	620 ± 40	640 ± 30	580 ± 50
Forefoot PPP (kPa)	630 ± 40	600 ± 40	620 ± 30	530 ± 60
Rearfoot PPP (kPa)	410 ± 20	440 ± 30	42 ± 20	450 ± 30
Forefoot/rearfoot PPP-Ratio	1.7 ± 0.2	1.5 ± 0.1	1.6 ± 0.1*	1.2 ± 0.1

Table 3. Achilles tendon mechanics and foot pressure mapping. Data are given as mean \pm SE. Different from CON, * $P < 0.05$, ** $P < 0.01$.
Achilles tendon mechanical properties determined at maximum and highest common force of 1815 N. Note: Modulus is based on average Achilles tendon CSA.

614
615

616

617 **Figure legends**

618

619 **Figure 1**

620 The Achilles tendon stress-strain relationship based on largest common tendon force observed for

621 merged Diabetic patients (DB) and age-matched healthy controls. Data are given as mean \pm SE. DB

622 showed higher Achilles tendon modulus than controls at highest common tendon force ($P < 0.001$).

623 A = Ultrasound Transducer, B = Strain Gauge.

624

625
626

627

628

629

630

631

632

633

