

Cranial vault trauma and selective mortality in medieval to early modern Denmark

Boldsen, Jesper L; Milner, George R; Weise, Svenja

Published in:

Proceedings of the National Academy of Sciences of the United States of America

DOI:

10.1073/pnas.1412511112

Publication date:

2015

Document version:

Final published version

Citation for pulished version (APA):

Boldsen, J. L., Milner, G. R., & Weise, S. (2015). Cranial vault trauma and selective mortality in medieval to early modern Denmark. *Proceedings of the National Academy of Sciences of the United States of America*, 112(6), 1721-1726. <https://doi.org/10.1073/pnas.1412511112>

Go to publication entry in University of Southern Denmark's Research Portal

Terms of use

This work is brought to you by the University of Southern Denmark.

Unless otherwise specified it has been shared according to the terms for self-archiving.

If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim. Please direct all enquiries to puresupport@bib.sdu.dk

Cranial vault trauma and selective mortality in medieval to early modern Denmark

Jesper L. Boldsen^{a,1}, George R. Milner^{b,1}, and Svenja Weise^c

^aDepartment of Anthropology, Institute of Forensic Medicine, University of Southern Denmark, DK 5260 Odense S, Denmark; ^bDepartment of Anthropology, Pennsylvania State University, University Park, PA 16802; ^cMax Planck Institute for Demographic Research, 18057 Rostock, Germany

Edited by Clark Spencer Larsen, The Ohio State University, Columbus, OH, and accepted by the Editorial Board December 24, 2014 (received for review July 2, 2014)

To date, no estimates of the long-term effect of cranial vault fractures on the risk of dying have been generated from historical or prehistoric skeletons. Excess mortality provides a perspective on the efficacy of modern treatment, as well as the human cost of cranial injuries largely related to interpersonal violence in past populations. Three medieval to early modern Danish skeletal samples are used to estimate the effect of selective mortality on males with cranial vault injuries who survived long enough for bones to heal. The risk of dying for these men was 6.2 times higher than it was for their uninjured counterparts, estimated through a simulation study based on skeletal observations. That is about twice the increased risk of dying experienced by modern people with traumatic brain injuries. The mortality data indicate the initial trauma was probably often accompanied by brain injury. Although the latter cannot be directly observed in skeletal remains, it can be inferred through the relative risks of dying. The ability to identify the effects of selective mortality in this skeletal sample indicates it must be taken into account in paleopathological research. The problem is analogous to extrapolating from death register data to modern communities, so epidemiological studies based on mortality data have the same inherent possibility of biases as analyses of ancient skeletons.

selective mortality | cranial trauma survival | violence | premodern Denmark

The aftereffects of head trauma are a major concern in modern medicine (1). However, for historical and prehistoric populations, there are no quantitative assessments of the long-term effects of such injuries, with the risk of dying estimated from skeletal samples being the most obvious way of doing so. Skeletons serve as a measure of the extent to which modern treatment and subsequent care have improved the outlook for survivors of trauma severe enough to fracture the cranial vault. Much of the cranial trauma in the distant past resulted from interpersonal violence, so these fractures provide a unique perspective on the human cost of conflicts that occur within communities up to the outright warfare that takes place between societies (2, 3).

The effect of selective mortality acting on people who experienced differential exposure or susceptibility to life-threatening diseases or trauma was brought to the attention of paleopathologists a quarter century ago (4). Debates continue over whether it is of any real significance when characterizing the health of past populations (5–16), although it has been shown that various conditions had an effect on mortality (17–23). That is largely because what comes to the paleopathologist's attention often indicates survival well beyond the original traumatic or disease episode, including cranial vault trauma with its potentially debilitating neurological consequences (24, 25).

Through a simulation study, healed cranial vault fractures are used to show that selective mortality has a measurable effect in archaeological skeletal samples. The simulation, based on skeletal observations, is a means of estimating the magnitude of the increased (or decreased) risk of dying associated with healed

cranial vault trauma, even if only slightly different from what everyone else experienced. Of primary importance are the age distributions of people—in this instance, men—with and without evidence of cranial vault injuries. Although the focus is on past societies, the problem is analogous to extrapolating from death register data to modern communities. Epidemiological studies based on mortality data have the same inherent possibility of biases as analyses of ancient skeletons.

Skeletal Samples

The skeletons came from three medieval to early modern Danish cemeteries dating to the late 12th to early 17th centuries. The graveyards held the remains of the inhabitants of rural and urban communities, so they collectively represent a cross-section of the population at that time. Two sites, Tirup (ca. AD 1150–1350) and St. Mikkel (ca. AD 1100–1529), were located in Jutland, and the third, Sortebrødre (ca. AD 1240–1610), was on the adjacent large island of Fyn. Odense, where the Sortebrødre churchyard was located, was inhabited by several thousand people during the period when most of the burials took place following the 1530s Reformation (26). St. Mikkel was located on the outskirts of a similarly sized town (27). Tirup, a small village, was likely occupied by less than a hundred people, and by the mid-14th century it had disappeared (28).

There were altogether 822 complete or partial skeletons, with Sortebrødre ($n = 451$) contributing the most individuals, followed by Tirup ($n = 217$) and St. Mikkel ($n = 154$). Of these skeletons, 236 adult males (>15 y) had enough of the cranium present in good condition for purposes of analysis (Table S1).

Significance

Neurocranial fractures and their aftermath took a toll on people in premodern societies, much like today. Archaeological information on skeletal trauma, however, typically consists of mere tallies of injuries, much like other disease-related lesions. We quantify the increased risk of dying for men with healed cranial vault fractures, an approach that can be adapted to any pathological condition. In medieval to early modern Denmark, head-injured men experienced a relative risk of dying about double that of modern people, probably in large part because of differences in medical care and social support. This approach provides a means of measuring the extent, hence consequences, of excess injury and disease-related mortality across the full range of human societies extending into the distant past.

Author contributions: J.L.B. and G.R.M. designed research; J.L.B., G.R.M., and S.W. performed research; J.L.B. and G.R.M. analyzed data; and J.L.B. and G.R.M. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. C.S.L. is a guest editor invited by the Editorial Board.

¹To whom correspondence may be addressed. Email: ost@psu.edu or jboldsen@health.sdu.dk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1412511112/-DCSupplemental.

Sex was estimated by G.R.M. for the entire sample. The great majority of the assessments are likely to be accurate because in blind trials in other work he correctly classified 98.5% of 813 modern known-sex skeletons from three continents. Only males were included in this study because females had too few cranial fractures for analytical purposes. Male and female mortality patterns were different, which prevented the sexes being combined.

Ages were estimated by G.R.M. for two samples, Sortebrødre and St. Mikkel; S.W. and J.L.B. estimated those for Tirup. Adult ages separately estimated by members of the research team are similar in examinations of European and American known-age skeletons, and there is a high correlation between estimated and actual ages (29, 30). The men who were not injured ranged in age from 15.5 to 72.5 y, and those with fractures from 17.5 to 65.0 y. The ages are midpoints of age intervals assigned to individual skeletons.

Bone fracture information was mostly collected by G.R.M., assisted at times by the late Ulla Freund under J.L.B.'s direction. Only healed fractures identified through visual examinations of skeletons are included here, although a few perimortem fractures are also present in the collections. In all cases, healed fractures showed considerable evidence of bone remodeling (Figs. 1 and 2). Many were simple depressions in the external table of the cranial vault, although some were indicative of more severe injuries, including deep cuts with heavy sharp-edged weapons such as swords and axes. It is usually impossible to identify what caused specific injuries, but it is likely that much of the trauma was from interpersonal violence. That differentiates historical populations from modern ones where traffic accidents are a common cause of head injuries. In all cases, the people in this analysis lived for a prolonged period following the traumatic incidents, to judge from the appearance of the bones.

As in all archaeological samples, the Danish skeletons were often incomplete. To be included here, the frontal and both parietal bones had to be present. That facilitated the recognition of healed fractures because the three bones had to be intact, hence well preserved. Ten frontal bones had fractures, and the



Fig. 1. A shallow depression fracture is present on the right side of the frontal bone in this Sortebrødre cranium. It is typical of the minor injuries seen on cranial vaults.



Fig. 2. A massive depression fracture can be seen on the left parietal of a Sortebrødre cranium, and a radiating fracture extends to the occipital bone. It is one of the most severe injuries that had healed.

left and right parietals had seven apiece. Only one person had a healed fracture elsewhere on the neurocranium. It was a radiating fracture on an occipital bone that originated in a massive depression fracture on a parietal. This individual, in fact, had two separate injuries, one on each parietal bone. One other individual had been hit twice, once on the frontal bone and again on a parietal. So the frontal and parietal bones serve as a reasonable estimate of the people who suffered severe cranial vault trauma, but afterward survived for a lengthy period.

Combining skeletons from separate sites is often necessary to secure large samples for rigorous quantitative analyses. That is true even of conditions, including cranial vault fractures, that occur relatively often in archaeological collections.

Having three sites representing a town, a community immediately outside of a town, and a rural village means that the skeletons provide an overall picture of what Danes of the 12th to early 17th centuries experienced. Major sociopolitical, economic, and demographic changes occurred during this half millennium, but they did not have the same impact in every community and on all segments of society. In the absence of precise temporal controls on individual burials and specific knowledge of local events, it is not possible to tease apart the exact circumstances that impinged on various segments of the three communities. So what these skeletons provide is a general depiction of life during the early part of Denmark's existence as a unified state when it was still a largely agrarian society dominated by small settlements. This period preceded the great growth of cities, commerce, and population associated with industrialization that only began in earnest in the 19th century.

Turning to the skeletons, trauma frequencies in the three samples are not significantly different from one another (χ^2 test, $P > 0.05$). The same is true when cemetery-of-origin and age-at-death are explanatory variables in a logistic regression analysis.

To characterize the effect of selective mortality on the age structure of people with and without cranial vault fractures, the ratio between the frequencies of people with fractures in the >15- to ≤35-y age group and those in the >35 group was calculated for both the simulated and observed individuals. That is, the frequency of young men with fractures (young men with fractures/all young men) was divided by the corresponding figures for old individuals. Ages were collapsed into two intervals, up to 35 y and above 35, because of the initial sample size. The number of skeletons was large by archaeological standards, but small for quantitative purposes. The age structures of the observed and expected categories were similar because the latter were modeled using parameter estimates derived from the former (Eqs. 2 and 4).

Results

The 236 males with complete frontal and parietal bones, 21 of which survived fractures, lived a total of 10,033 y. The mean age at death was 42.5 y. Injured men had a mean age of death of 41.0 y, whereas the corresponding figure for those who were not injured was 42.7 (Table S2).

In Table 1, the frequency of cranial vault injuries in the >35-y age group was lower than it was in younger men, consistent with healed fractures being associated with an increased risk of dying. That difference, however, is not significantly different (χ^2 test, $P > 0.05$), nor was it expected to have been significant because the increased risk of dying experienced by trauma survivors was presumed at the outset to have been too low to be detectable in a small sample using conventional statistical tests based on data aggregated into “young” versus “old.” The simulation provides a means of quantifying the *RR* implied in the Table 1 observed percentages and ratio, the latter the frequency of young men with healed cranial fractures relative to older men.

The Expected (Exp) column under *RR* = 1 in Table 1 shows the number of fractures for the young and old age groups when there is no selectivity. The best-fitting relative risk conforming to the observed fractures in the skeletal sample is *RR* = 6.2 (Fig. 3). The significance of the difference between the observed and expected fractures is characterized by the χ^2 values for *df* = 1 of 15.76 for *RR* = 1, and 0.00 for *RR* = 6.2.

In Fig. 3, the expected young to old fracture frequency ratio, generated from the simulation, is plotted relative to the observed fracture ratio. The *x* axis is the relative risk of dying on a log scale, with *RR* = 1 indicating that injured people experienced no increased risk. The curve depicting the ratio between young and old fracture frequencies is not perfectly smooth because it shows simulation results. The intersection at *RR* = 6.2 of the expected line with the observed values is the increase in the risk of dying for men with healed frontal or parietal bone fractures.

There remains the issue of estimating a confidence interval for the relative risk of dying associated with having a healed cranial vault fracture. The upper limit approaches infinity where the




Fig. 3. The ratio of fracture frequencies among young (>15 to ≤35 y) relative to old (>35 y) men for different values of relative risk (*RR*) (solid curve) is compared with the observed value (*RR* = 6.2) (dashed line). Relative risk refers to the number of times greater the risk of dying was for men with healed skull fractures than it was for men who lacked those injuries.

expected age pattern does not differ significantly from the observed age pattern. As explained above, the two age distributions would be identical if everyone died without bone fractures showing signs of healing. Although that cannot be excluded on statistical grounds, the skeletons clearly indicate that many men survived cranial vault fractures. The lower limit is the *RR* value corresponding to the χ^2 value 3.84, indicating $P = 0.05$; in this instance, *RR* = 2.3.

Discussion

The mere number of cranial fractures underscores the fact that interpersonal violence was an ever-present aspect of life for men in medieval to early modern Denmark. Of all males, 8.9% had healed fractures of the cranial vault. Age-corrected, that figure corresponds to around three times the fracture rate in contemporary Denmark [based on *R* (Eq. 1) and modern Danish fracture data provided by the Accident Prevention Group at Odense University Hospital, as described in ref. 31.] Although in most instances it is not possible to identify the origin of the trauma, the fact that it is heavily weighted toward men—males in the skeletal sample were about four times more likely to have cranial injuries than females—suggests that many cranial fractures resulted from interpersonal violence. That is consistent with the argument, based on written sources, that there was generally more violence in societies hundreds of years ago than there is today (32). Accidents, often occupation related, were also common because there was little to protect people from the hazards of physically demanding work (33).

Fracture frequencies, of course, depend on the rate of fracture acquisition and the period of exposure. If rates of acquisition and years of exposure were the only two processes of importance, then one would expect increasing fracture frequencies with advancing age in archaeological (mortality) samples. Quite the opposite seems to have occurred, as shown in the observed columns in Table 1 where men with healed fractures dropped

Table 1. Adult males (*N*), observed healed cranial fractures (Frac), and the percentage (%) injured in each age group

Age interval	Observed			<i>RR</i> = 1		<i>RR</i> = 6.2	
	<i>N</i>	Frac	%	Exp	%	Exp	%
>15 to ≤35	75	9	12.0	2.8	3.7	9.1	12.1
>35	161	12	7.5	18.8	11.7	11.9	7.4
Ratio			1.61		0.32		1.64

Expected (Exp) fractures for males are given for relative risks (*RR*) 1 and 6.2 times higher for men who survived the initial injuries, along with percentages (%). See Fig. 3 for the intersection of observed and expected at 6.2. Ratio is the frequency in the young group divided by the frequency in the old group.

from 12.0% to 7.5% in the young and old groups, respectively. A decline in frequencies could only come about through a process where the cemetery sample at each age is enriched by people who sustained, but survived, cranial vault fractures. Although these men lived long enough for fractures to heal, they were more likely to enter the mortality sample than men without fractures. Archaeological samples, however, are rarely large enough for simple counts of pathological specimens to be useful when assessing subtle differences in relative risk, such as what might be expected if healed bone fractures were associated with higher mortality.

Through simulating the expected life spans of individuals with and without healed cranial vault trauma, it appears that men with injuries sufficient to produce an indelible mark on the frontal or parietal bones experienced a risk of dying six times that of their contemporaries who avoided such fractures. That is much like today where people who suffer head trauma leading to brain injuries also have a higher risk of dying (34–45). Modern figures vary according to injury severity; where and when the data were collected; sample composition, especially the patient age distribution and inclusion criteria; and posttrauma study duration. Nevertheless, today people with traumatic brain injuries experience roughly two to four times the risk of dying than uninjured people (37, 38, 40–45).

The two sets of figures—cranial vault fractures and traumatic brain injuries—are not directly comparable because one can occur in the absence of the other. Nevertheless, it is reasonable to suppose that brain injuries often accompanied trauma sufficient to fracture the frontal and parietal bones. The cranial vault fractures in the Danish sample, therefore, serve as a crude proxy for traumatic brain injuries with lingering consequences. Although modern relative risks can be within the lower confidence limit of the archaeological sample, it is unlikely the two are indeed the same. Relative to today, effective care was lacking in medieval to early modern Denmark. Injured people now benefit from sophisticated medical procedures plus long-term support and rehabilitation that collectively prolong life. It is not surprising that today's medical and rehabilitative treatments and social services reduce considerably the risk of dying from what it was many centuries ago. Estimating the extent of that improvement is another matter altogether. Although skeletons provide only a rough approximation of relative risk and the modern and archaeological data are not strictly comparable, the Danish data suggest the risk of dying experienced by survivors of cranial vault injuries has been about halved in modern times.

Relative risk is a standard measure of the increased risk of dying within a study population. It can be misleading, however, when used in comparisons of samples with widely different baseline mortality rates. In the medieval and early modern Danish sample, the risk of dying around age 35 was over 40 times what Danes experience today (3.63% versus 0.08%). Therefore, a half-millennium ago the increased risk of dying faced by cranial-fracture survivors of any particular age was many times greater than it might appear from the relative risk values alone.

In modern populations, the relative risk of dying for trauma survivors declines noticeably with advancing age (36, 42, 44). That is a result of an age-increasing background mortality and the fact that the healthiest trauma survivors reach high ages. The latter is an example of selective mortality in a group that was disadvantaged at the outset by their injuries.

Mortality immediately following head injuries must have been high a half-millennium ago in Denmark, much like it is today. There were, in fact, several skeletons in the three-site sample with obvious perimortem cranial trauma, including massive fractures and deep cuts from sharp-edged weapons (swords or axes). Survivors, in contrast, typically displayed trauma that was not as severe, although two men survived grievous injuries from sharp-edged weapons. Unfortunately, the difficulty in consistently

identifying examples of perimortem fractures in archaeological bones prohibits unbiased estimates of more or less immediate fatalities from cranial injuries.

Higher mortality for survivors of cranial vault trauma could have come about in two ways. Some men might have had greater exposure to violent encounters or hazardous occupations than others, so the cranial injuries are an archaeologically visible marker of ways of life that likely shortened some people's life spans relative to their peers. Alternatively, disabilities acquired from severe blows that resulted in underlying soft tissue damage increased the survivors' risks of dying from the time of their injuries onward. That might include a loss of function or changed behavior, much like the motor, cognitive, and psychological impairment that can follow traumatic brain injuries today. Because only the outcome is observable in archaeological remains—a greater relative risk of dying—it is not possible to discriminate between the two alternatives, both of which probably contributed to our results.

As for the lingering consequences of severe injuries, anything that led to difficulties adjusting to life must have been a major problem in relatively small and largely self-sufficient communities where everyone was expected to contribute substantively to his or her own livelihood. Anything short of full and productive integration into household and community life must have posed a mortality risk, at the very least during periods when food was short. When times were hard, the poor and otherwise disadvantaged members of medieval communities elsewhere in Europe appear to have been more likely to succumb to death than their neighbors (46). Skeletal and documentary sources indicate that people with physical or mental impairments were on occasion the recipients of institutionalized care (33, 47, 48). However, relative to today, the survivors of debilitating trauma enjoyed few formally organized services that acted as safety nets to keep them healthy, hence alive, and instead they had to rely heavily on the generosity of family and community members, or resort to begging and thievery (33, 46). In mid-16th century Denmark, an effort to create new hospitals proved to be unsuccessful, and soon thereafter market towns were again permitted to maintain their own institutions as they saw fit (49). It took until the early 18th century before national legislation was passed to cover care for the poor, although there were earlier local efforts to regulate begging (50). Estimating the excess mortality associated with having previously suffered trauma moves us closer to an understanding of the efficacy of institutional and improvised care in past societies.

By looking at pathological skeletal features in conjunction with age distributions, it is possible to quantify the disadvantage experienced by individuals who survived earlier illnesses and injuries, or who suffered from chronic conditions. The overall objective is to move components of the heterogeneity in the risk of dying from being essentially hidden, or unknown, to measurable, or known.

Of immediate archaeological significance is the potential for healed skeletal trauma to be used as a marker to subdivide populations according to life experiences, analogous to an epidemiological risk factor. Although what precipitated the death of any single individual can only be established on exceedingly rare occasions, it is possible to identify cranial vault fractures (and other pathological conditions) and to measure the increase in mortality associated with them on a population basis. The intent of this work is to identify recognizable skeletal features associated with a greater risk of dying and to assess their effect on populations in terms of lost years of life. It is not to establish what caused the premature death of particular individuals or the mechanisms behind sickness and death.

Such analyses, however, require many skeletons, and that will often require combining samples. In this instance, doing so provides an overall picture of excess mortality associated with

healed cranial vault fractures for a temporally long and socially diverse sample. Hundreds of additional skeletons from medieval to early modern Denmark must be examined to extend this proof of concept to finer time intervals and particular segments of society, such as those distinguished by social position, occupation, disease experience, and diet.

Conclusion

We will never know what happened to specific individuals who experienced cranial vault trauma. In general, however, the survivors of these injuries lived shorter lives than they would have otherwise. From the perspective of characterizing life in the past, knowing that a healed cranial vault fracture was associated with an increased risk of dying is important. It is likely that many of

these men suffered the lingering effects of traumatic brain injury. They did not die because of what can be observed, the healed bone fracture, but perhaps from circumstances related to what cannot be directly seen, the long-term health consequences of the initial injury. The fact that the effects of selective mortality can be teased out of skeletal samples for healed cranial fractures indicates that it must be taken into consideration in archaeological analyses and, more generally, epidemiological studies based on death data.

ACKNOWLEDGMENTS. The late Ulla Freund's assistance with the skeletal materials is much appreciated. Her insights and familiarity with the collections are sorely missed. This work was partly supported by G.R.M.'s grant from the American-Scandinavian Foundation.

- Masel BE, DeWitt DS (2010) Traumatic brain injury: A disease process, not an event. *J Neurotrauma* 27(8):1529–1540.
- Walker PL (1997) Wife beating, boxing, and broken noses: Skeletal evidence for the cultural patterning of violence. *Troubled Times: Violence and Warfare in the Past*, eds Martin DL, Frayer DW (Gordon and Breach, Amsterdam), pp 145–179.
- Walker PL (2001) A bioarchaeological perspective on the history of violence. *Annu Rev Anthropol* 30:573–596.
- Wood JW, Milner GR, Harpending HC, Weiss KM (1992) The osteological paradox: Problems of inferring prehistoric health from skeletal samples. *Curr Anthropol* 33(4):343–370.
- Cohen MN (1992) Comments. *Curr Anthropol* 33(4):358–359.
- Goodman AH (1993) On the interpretation of health from skeletal remains. *Curr Anthropol* 34(3):281–288.
- Jackes MK (1993) On paradox and osteology. *Curr Anthropol* 34(4):434–439.
- Saunders SR, Hoppa RD (1993) Growth deficit in survivors and non-survivors: Biological mortality bias in subadult skeletal samples. *Yearb Phys Anthropol* 36:127–151.
- Byers SN (1994) On stress and stature in the “osteological paradox.” *Curr Anthropol* 35(3):282–284.
- Cohen MN (1994) The osteological paradox reconsidered. *Curr Anthropol* 35(5):629–631.
- Wood JW, Milner GR (1994) Reply. *Curr Anthropol* 35(5):631–637.
- Cohen MN (1997) Does paleopathology measure community health? A rebuttal of “the osteological paradox” and its implication for world history. *Integrating Archaeological Demography: Multidisciplinary Approaches to Prehistoric Population*, ed Paine RR (Center for Archaeological Investigations, Southern Illinois University, Carbondale, IL), pp 242–260.
- Wright LE, Yoder CJ (2003) Recent progress in bioarchaeology: Approaches to the osteological paradox. *J Archaeol Res* 11(1):43–70.
- Cohen MN, Crane-Kramer GMM (2007) Editors' summation. *Ancient Health*, eds Cohen MN, Crane-Kramer GMM (Univ Press of Florida, Gainesville, FL), pp 320–343.
- Boldsen JL, Milner GR (2012) An epidemiological approach to paleopathology. *A Companion to Paleopathology*, ed Grauer AL (Wiley-Blackwell, Chichester, UK), pp 114–132.
- Milner GR (2013) Trauma in the medieval to early modern Sortebrødre skeletons from Odense, Denmark. *The Dead Tell Tales: Essays in Honor of Jane E. Buikstra*, eds Lozada MC, O'Donnabhain B (Cotsen Institute of Archaeology, University of California, Los Angeles), pp 172–185.
- Goodman AH, Armelagos GJ (1988) Childhood stress and decreased longevity in a prehistoric population. *Am Anthropol* 90(4):936–994.
- Duray SM (1996) Dental indicators of stress and reduced age at death in prehistoric Native Americans. *Am J Phys Anthropol* 99(2):275–286.
- Boldsen JL (1997) Estimating patterns of disease and mortality in a medieval Danish village. *Integrating Archaeological Demography: Multidisciplinary Approaches to Prehistoric Population*, ed Paine RR (Center for Archaeological Investigations, Southern Illinois University, Carbondale, IL), pp 229–241.
- Usher BM (2000) A multistate model of health and mortality for paleodemography: The Tirup cemetery. PhD dissertation (Pennsylvania State University, University Park, PA).
- Palubeckaitė Z, Jankauskas R, Boldsen J (2002) Enamel hypoplasia in Danish and Lithuanian late medieval/early modern samples. A possible reflection of child morbidity and mortality patterns. *Int J Osteoarchaeol* 12(3):189–201.
- Boldsen JL (2007) Early childhood stress and adult age mortality—a study of dental enamel hypoplasia in the medieval Danish village of Tirup. *Am J Phys Anthropol* 132(1):59–66.
- DeWitte SN, Wood JW (2008) Selectivity of black death mortality with respect to preexisting health. *Proc Natl Acad Sci USA* 105(5):1436–1441.
- Weber J, Czarnetzki A (2001) Brief communication: Neurotraumatological aspects of head injuries resulting from sharp and blunt force in the early medieval period of southwestern Germany. *Am J Phys Anthropol* 114(4):352–356.
- Harrod RP, Martin DL (2014) Signatures of captivity and subordination on skeletonized human remains? A bioarchaeological case study from the ancient Southwest. *Bioarchaeological and Forensic Perspectives on Violence*, eds Martin DL, Anderson CP (Cambridge Univ Press, Cambridge, UK), pp 103–119.
- Boldsen JL, Møllerup L (2006) Outside St. Jørgen: Leprosy in the medieval Danish city of Odense. *Am J Phys Anthropol* 130(3):344–351.
- Petersen HC, Boldsen JL, Paine RR (2006) Population relationships in and around medieval Danish towns. *Urbanism in the Preindustrial World: Cross-Cultural Approaches*, ed Storey GR (Univ of Alabama Press, Tuscaloosa, AL), pp 110–120.
- Boldsen JL (2002) Demografisk struktur i landsbyen Tirup. *Hikuin* 27:233–244.
- Milner GR, Ferrell RJ (2011) Conflict and death in a late prehistoric community in the American Midwest. *Anthropol Anz* 68(4):415–436.
- Milner GR, Boldsen JL (2012) Transition analysis: A validation study with known-age modern American skeletons. *Am J Phys Anthropol* 148(1):98–110.
- Faergemann C, Lauritsen JM, Brink O, Skov O (2007) Trends in deliberate interpersonal violence in the Odense Municipality, Denmark 1991–2002. The Odense study on deliberate interpersonal violence. *J Forensic Leg Med* 14(1):20–26.
- Pinker S (2011) *The Better Angels of Our Nature: Why Violence Has Declined* (Viking, New York).
- Metzler I (2013) *A Social History of Disability in the Middle Ages* (Routledge, New York).
- Walker AE, Leuchs HK, Lechtape-Grüter H, Caveness WF, Kretschman C (1971) Life expectancy of head injured men with and without epilepsy. *Arch Neurol* 24(2):95–100.
- Weiss GH, Caveness WF, Einsiedel-Lechtape H, McNeel ML (1982) Life expectancy and causes of death in a group of head-injured veterans of World War I. *Arch Neurol* 39(12):741–743.
- Strauss D, Shavelle R, DeVivo MJ (1999) Life tables for people with traumatic brain injury. *J Insur Med* 31(2):104–105.
- Baguley I, Siewa-Younan S, Lazarus R, Green A (2000) Long-term mortality trends in patients with traumatic brain injury. *Brain Inj* 14(6):505–512.
- Shavelle R, Strauss D (2000) Comparative mortality of adults with traumatic brain injury in California, 1988–97. *J Insur Med* 32(3):163–166.
- Brown AW, et al. (2004) Long-term survival after traumatic brain injury: A population-based analysis. *NeuroRehabilitation* 19(1):37–43.
- Ratcliff G, Colantonio A, Escobar M, Chase S, Vernich L (2005) Long-term survival following traumatic brain injury. *Disabil Rehabil* 27(6):305–314.
- Cameron CM, Purdie DM, Kliwer EV, McClure RJ (2008) Ten-year outcomes following traumatic brain injury: A population-based cohort. *Brain Inj* 22(6):437–449.
- Baguley IJ, et al. (2012) Late mortality after severe traumatic brain injury in New South Wales: A multicentre study. *Med J Aust* 196(1):40–45.
- Harrison-Felix C, et al. (2012) Mortality after surviving traumatic brain injury: Risks based on age groups. *J Head Trauma Rehabil* 27(6):E45–E56.
- Brooks JC, et al. (2013) Long-term disability and survival in traumatic brain injury: Results from the National Institute on Disability and Rehabilitation Research Model Systems. *Arch Phys Med Rehabil* 94(11):2203–2209.
- Fazel S, Wolf A, Pillas D, Lichtenstein P, Långström N (2014) Suicide, fatal injuries, and other causes of premature mortality in patients with traumatic brain injury: A 41-year Swedish population study. *JAMA Psychiatry* 71(3):326–333.
- Hanawalt BA (1986) *The Ties That Bind* (Oxford Univ Press, New York).
- Grauer AL, Roberts CA (1996) Paleoepidemiology, healing, and possible treatment of trauma in the medieval cemetery population of St. Helen-on-the-Walls, York, England. *Am J Phys Anthropol* 100(4):531–544.
- Metzler I (2011) Disability in the Middle Ages: Impairment at the intersection of historical inquiry and disability studies. *Hist Compass* 9(1):45–60.
- Grell OP, Cunningham A (1997) The Reformation and changes in welfare provision in early modern northern Europe. *Health Care and Poor Relief in Protestant Europe 1500–1700*, eds Grell OP, Cunningham A (Routledge, London), pp 1–41.
- Hansen BD (1996) Fattivæsen. *Den Store Danske Encyklopædi*, ed Lund J (Gyldendal, Copenhagen), pp 237–238.