



University of Southern Denmark

## Erratum to

### **The GOOD life: Study protocol for a social norms intervention to reduce alcohol and other drug use among Danish adolescents**

Stock, Christiane; Vallentin-Holbech, Lotte; Rasmussen, Birthe Marie

*Published in:*  
B M C Public Health

*DOI:*  
10.1186/s12889-017-4444-z

*Publication date:*  
2017

*Document version:*  
Final published version

*Document license:*  
CC BY

*Citation for pulished version (APA):*

Stock, C., Vallentin-Holbech, L., & Rasmussen, B. M. (2017). Erratum to: The GOOD life: Study protocol for a social norms intervention to reduce alcohol and other drug use among Danish adolescents. *B M C Public Health*, 17, [501]. <https://doi.org/10.1186/s12889-017-4444-z>

Go to publication entry in University of Southern Denmark's Research Portal

#### **Terms of use**

This work is brought to you by the University of Southern Denmark.  
Unless otherwise specified it has been shared according to the terms for self-archiving.  
If no other license is stated, these terms apply:

- You may download this work for personal use only.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying this open access version

If you believe that this document breaches copyright please contact us providing details and we will investigate your claim.  
Please direct all enquiries to [puresupport@bib.sdu.dk](mailto:puresupport@bib.sdu.dk)

ERRATUM

Open Access



# Erratum to: The GOOD life: Study protocol for a social norms intervention to reduce alcohol and other drug use among Danish adolescents

Christiane Stock\*, Lotte Vallentin-Holbech and Birthe Marie Rasmussen

## Erratum

Following publication of this study design article [1], it has come to our attention that some terminology was inconsistently used in the article. The GOOD life trial uses a definition of binge drinking of “5 or more drinks on one occasion” in accordance with the ESPAD study [2]. Both in the abstract and in the methods section, when describing measures of alcohol use or outcomes, any formulation of “more than 5 drinks” becomes “5 or more drinks”. Also, when respondents were asked to identify the amount of occasions they have been drinking 5 or more drinks in the past 30 days, the binge drinking definition was not according to the MULDD study [3], but according to the ESPAD study [2].

Received: 15 May 2017 Accepted: 17 May 2017  
Published online: 24 May 2017

## References

1. Stock C, Vallentin-Holbech L, Rasmussen BM. The GOOD life: Study protocol for a social norms intervention to reduce alcohol and other drug use among Danish adolescents. *BMC Public Health*. 2016;16:704.
2. Hibell B, Guttormsson U, Ahlström S, Balakireva O, Bjarnason T, Kokkevi A, Kraus L. The 2011 ESPAD report-substance use among students in 36 European countries. Stockholm: The Swedish Council for Information on Alcohol and Other Drugs (CAN); 2012.
3. Nielsen GA, Ringgaard L, Broholm K, Sindballe A, Olsen SF. Unge's livsstil og dagligdag 2000: forbrug af tobak, alkohol og stoffer [Young people's lifestyle and daily life: use of tobacco, alcohol and drugs]. Copenhagen: The Danish Cancer Society and National Board of Health; 2002.

\* Correspondence: cstock@health.sdu.dk  
Unit for Health Promotion Research, Department of Public Health, University of Southern Denmark, Esbjerg, Denmark

